
Intra-guild competition and ecosystem services of mammal scavengers in a new colonized wolf landscape









Intra-guild competition and ecosystem services of mammal scavengers in a new colonized wolf landscape

R. Brogi*1, P. Bongi1, M. Del Frate1, S. Sieni1, A. Cavallera2, & M. Apollonio1


- 1 Department of Veterinary Medicine, University of Sassari, Italy
- 2 Department of Agri-food Production and Environmental Sciences (DISPAA), University of Florence, Italy
- *rbrogi@uniss.it






Wolf return → consequences on the ecosystem level

Low human impact




Wolf return → consequences on the ecosystem level

Low human impact



Kuijper et al. 2024 J Applied Ecology

Human-modified environments



Novel ecosystem functions


"wolves in human-modified ecosystems will engage in an array of novel interactions and potential <u>novel trophic</u> cascades"

Wolf return to human-dominated, prey-rich landscapes




Wolf return to human-dominated, prey-rich landscapes





- Who is going to win the competition for the emerging resource represented by deer carcasses?
- How does scavenger competition for wolf-killed carcasses change over time following wolf return?

Study area








Methods

- 12,000+ km transects since 2017 to 2022
- 103 fresh wolf-killed fallow deer carcasses
- monitored by camera traps for 7-10 days



Only 3 mammal scavengers:









Methods









- Likelihood of finding and using the carcass
- Total time spent feeding on the carcass
- Outcome of direct interactions
- Interspecific spatiotemporal associations

modelled across species
and along the 5 years
and along period
monitoring period

Likelihood of finding and using the carcass ~ species*time + carcass weight + re(carcass id)



Total time spent feeding on the carcass ~ species*time + carcass weight + re(carcass id)



Outcome of direct interactions on the carcasses



Outcome of direct interactions on the carcasses




Outcome of direct interactions on the carcasses (n=44)




Interspecific spatiotemporal association ~ species*time + re(carcass id)

i.e., delay in arriving on a carcass after the competitor species




Interspecific spatiotemporal association ~ species*time + re(carcass id)

i.e., delay in arriving on a carcass after the competitor species



Interspecific spatiotemporal association ~ species*time + re(carcass id)

i.e., delay in arriving on a carcass after the competitor species



Red foxes did not avoid wild boars; No variations along the study period



Mostly indirect competition





- Wolves did not even take part
- Wild boar > red fox


≠ previous studies

Vucetich et al. 2012, Focardi et al. 2017

Selva et al. 2005, Bassi et al. 2018











91 ± 10 fallow deer/km²



 wolves up, deer down?

91 ± 10 fallow deer/km²




 wolves up, deer down?





CONCLUSIONS

 Wild boar competed with red foxes but progressively monopolized the emerging niche



Wolf return + unnaturally high prey density >
 enhanced predictability of carcass availability



 Minimal carcass reutilization by wolves, but what with the increasing wolf population?



Thanks for listening!



Rudy Brogi rbrogi@uniss.it



