

Advance Publication

# The Journal of Veterinary Medical Science

Accepted Date: 7 May 2022

J-STAGE Advance Published Date: 22 February 2024

©2024 The Japanese Society of Veterinary Science

Author manuscripts have been peer reviewed and accepted for publication but have not yet been edited.

1 *Full paper*

2 Wildlife Science

3 **Quantitative analysis of the skull in the Japanese wolf (*Canis lupus***  
4 ***hodophilax*) using CT**

5  
6 Chihiro SUZUKI<sup>1)</sup>, Motoki SASAKI<sup>1)\*</sup>, Nao TSUZUKI<sup>1)</sup>, Mitsunori KAYANO<sup>1)</sup>, Kazutaka  
7 YAMADA<sup>2)</sup>, Naotaka ISHIGURO<sup>3)</sup>, Satoshi SUZUKI<sup>4)</sup>, Hajime TARU<sup>4)</sup>, Wataru MATSUDA<sup>5)</sup>,  
8 Hideki ENDO<sup>6)</sup>, Tomoaki KIKUCHI<sup>7)</sup>, Kaoru KIKUCHI<sup>8)</sup> and Nobuo KITAMURA<sup>1)</sup>

9  
10 <sup>1</sup>Obihiro University of Agriculture and Veterinary Medicine, Obihiro, Hokkaido 080-8555, Japan

11 <sup>2</sup>Azabu University, Sagamihara, Kanagawa 252-5201, Japan

12 <sup>3</sup>The Graduate University for Advanced Studies, Miura, Kanagawa 240-0193, Japan

13 <sup>4</sup>Kanagawa Prefectural Museum of Natural History, Odawara, Kanagawa 250-0031, Japan

14 <sup>5</sup>Oyodo-cho Town Official Board of Education, Yoshino, Nara 638-8501, Japan

15 <sup>6</sup>The University Museum, The University of Tokyo, Tokyo 113-0033, Japan

16 <sup>7</sup>Hokkaido Chuo Agricultural Mutual Aid Association, Fukagawa, Hokkaido 074-0001, Japan

17 <sup>8</sup>Kikuchi Veterinary Clinic, Oshu, Iwate 023-1131, Japan

18  
19 \*Correspondence to: SASAKI, M.: e-mail: sasakim@obihiro.ac.jp

20  
21 **Running head: MORPHOLOGY OF THE JAPANESE WOLF SKULL**

**ABSTRACT**

In this study using computed tomography (CT), the volumes of the internal cranial cavities, such as the braincase, frontal sinus and tympanic cavity, and the ratio of the volume of each cavity to the skull volume in Japanese wolves were quantified, and CT images of the frontal sinus were observed. The results were then compared with those of other wolf subspecies, including Akita, a dog breed, to clarify the characteristics of the internal cranial cavities in Japanese wolves. The present study revealed that the Japanese wolf had a relatively larger braincase volume and a relatively smaller frontal sinus volume than the wolf ssp. (a group of wild wolf subspecies except the Japanese wolf) and Akita. Moreover, the relative and absolute tympanic cavity volumes of the Japanese wolf and Akita were significantly smaller than those of the wolf ssp. In the CT image or macroscopic observations, the frontal sinuses of the wolf ssp. and Akita were relatively well developed to the caudal and dorsal directions, respectively, compared with that of the Japanese wolf, and the tympanic cavity of the wolf ssp. was more largely swelled ventrally and medially than that of other groups.

**KEY WORDS:** braincase, CT, frontal sinus, Japanese wolf, tympanic cavity

## INTRODUCTION

The gray wolf or wolf (*Canis lupus* Linnaeus, 1758), which is the largest canids, inhabits the Eurasian and North American continents and the North Pole [32, 36]. Dogs (*C. l. familiaris*), classified as subspecies of wolves, were domesticated from wild wolves by humans 15,000 years ago [41, 46]; however, the place of domestication remains controversial. A study on the genetic diversity with mitochondrial DNA (mtDNA) information suggested a Southeast Asian origin for domestication [41], whereas the single nucleotide polymorphism (SNP) genotyping array analysis suggested a simultaneous origin in several areas worldwide [53].

The Japanese wolf (*C. l. hodophilax*) is one of the subspecies of gray wolves and inhabited Honshu, Shikoku and Kyushu islands. Japanese wolves have a distinct mtDNA haplotype and are able to be genetically distinguished from other gray wolves [20, 22-24, 34, 35]. A recent phylogenetic study using the nuclear DNA (nDNA) extracted from nine Japanese wolves suggested that Japanese wolves are genetically separated from other wolves, including ancient wolves [14]. It is also possible to distinguish Japanese wolves from other subspecies by common morphological characters. For example, most Japanese wolves have a rostral alar foramen separated by a bony septum and a horizontal plate of the palatine bone with an anterior notch (Fig. 1) [15, 17, 18, 37-39, 51]. Regarding skull size, that of Japanese wolves is smaller than that of most other subspecies [17].

In the Meiji era (1868-1912), infectious diseases, such as rabies, introduced from overseas spread among Japanese wolves. In addition, with the development of the livestock industry in Japan, the extermination of Japanese wolves was recommended as they attacked livestock [10, 15, 54]. As a result, the population of the Japanese wolf decreased and a male wolf caught at Higashi Yoshino village in Nara prefecture on January 23, 1905 (38th year of Meiji) became the last Japanese wolf; therefore, this subspecies is believed to be extinct [10, 15, 44, 54].

In Japan, as taxidermy and preservation skills were underdeveloped before the extinction of Japanese wolves, there are few skin and skeletal specimens of Japanese wolves available. Among morphological studies on the skull of Japanese wolves, the external morphology was actively examined because of easy access to the target parts of the skull, whereas studies on the internal structure are difficult without destroying the skull. Recently, non-destructive analysis of the internal cranial structures in many species, including extinct species, has become possible by the development and widespread use of computed tomography (CT) and related software. There is an increasing number of reports on the volume and morphology of the cranial cavity, which houses the brain, the frontal sinus, which is a part of the paranasal sinuses, and the tympanic cavity, which forms a part of the middle ear [2, 3, 7, 9, 13, 28, 47]. In the examination of the internal cranial structure of several mammals, the classification of fossil species and the functions of the

internal morphology have been reported [27, 31, 33, 43, 48, 50]. In this manner, non-destructive analyses of the internal cranial structure using CT have provided new findings in many animals.

In the gray wolf, comparative morphological studies of internal cranial structures have been reported with several other species [6, 30], and between wild populations and domestic dogs [45].

On the other hand, there are only two non-destructive studies of the internal cranial structure of the Japanese wolf [11, 12]. However, these studies did not quantitatively or three-dimensionally examine the internal structures of the skull. The internal cranial cavities, such as the braincase, frontal sinus and tympanic cavity are clear areas surrounded by bones, and it is important to examine these internal cranial structures with external one for a comprehensive understanding of the skull. The aim of this study was to clarify the characteristics of these internal cranial cavities in Japanese wolves by examining the structures non-destructively, quantitatively and three-dimensionally, and then comparing the results with those of other extant gray wolves.

## **MATERIALS AND METHODS**

### **Specimens**

Specimens used in this study are shown in Tables 1-3. In this study, the specimens with the mtDNA haplotype of the Japanese wolf by genetic analysis [22] and the external cranial characteristics of the Japanese wolf were defined as Japanese wolf, and other specimens of wild

wolves were defined as wolf ssp. To compare wild wolves with domestic dogs, the skulls of the Akita breed, a native Japanese dog that resembles Japanese wolves in cranial size [11, 56], were used and defined as Akita. In this study, a total of 24 samples of wolves including dogs (4 Japanese wolf; 16 wolf ssp.; 4 Akita) were analyzed. As most samples had no information of sex, the analyses were performed without consideration of sex. Although there was no information of age for all samples, we estimated that all wolves were adults because of the complete closure of cranial sutures.

#### **CT scanning**

All specimens were scanned by a CT scanner (Aquilion LD, Canon, Tokyo, Japan). Scanning conditions were 120 kV, 150 mA and 0.5-mm slice thickness, and scanning data were preserved in DICOM format. A private specimen of the Japanese wolf (ID, Shimane) was covered with dry soft tissue and scanned under different scanning conditions (135 kV and 250 mA). Moreover, KPM-NF1 2913 had a metallic bolt in the occipital bone and was scanned at 135 kV and 300 mA.

#### **Measurement of volume**

Using the medical workstation for image processing (Virtual Place Fujin, AZE, Tokyo, Japan), the braincase, bilateral frontal sinus and tympanic cavity were three-dimensionally reconstructed from DICOM data (Fig. 2), and each volume was measured. In the frontal sinus and tympanic cavity, both sides were summed and used for statistical analysis. To compare the ratio of the volume of each cavity to the skull volume, the volume of each skull was measured except tooth volume. M1185 was missing a part of the right zygomatic arch; therefore, the volume of the left half of the skull was doubled. Similarly, M25794 and KPM-NF 2014 had a damaged tympanic cavity, the volume of the skull was calculated by doubling the intact side [25, 40], and for the broken tympanic cavity, the same value of the intact cavity was used (Table 4). M26696 and M25796 were omitted from the analysis of the volume ratio of each cavity to the skull volume because of breakage of the occipital part across both sides.

## **Statistical analysis**

The median and mean volumes of the braincase, frontal sinus and tympanic cavity, and the volume ratio of each cavity to the skull volume were calculated in each group. To compare the values among three groups, the Steel-Dwass method was performed with R version 4.0.3 (significant difference  $P<0.05$ ).



## RESULTS

### Volume of the skull

The volume of each cavity and the ratio of each cavity volume to the skull volume in individuals are shown in Table 4. Regarding the skull volume, the medians (means $\pm$ standard error, SE) of the Japanese wolf, wolf ssp. and Akita were 407.0 ml (402.5 $\pm$ 33.0 ml), 571.1 ml (568.0 $\pm$ 24.6 ml) and 445.7 ml (439.4 $\pm$ 45.9 ml), respectively. The wolf ssp. had the largest skull volume among the three groups, followed by the Akita and the Japanese wolf (Fig. 3). In the statistical analysis, the skull volume of the Japanese wolf was significantly smaller than that of the wolf ssp. ( $P<0.05$ ; Fig. 3). The skull volume of the Akita tended to be smaller than that of the wolf ssp. ( $P=0.090$ ), although there was no significant difference between the Akita and wolf ssp.

### Braincase

The medians (means) of the braincase volume of the Japanese wolf, wolf ssp. and Akita were 126.4 ml (123.1 $\pm$ 8.3 ml), 149.3 ml (150.4 $\pm$ 3.3 ml) and 103.2 ml (103.0 $\pm$ 3.0 ml), respectively. Based on the statistical analysis, the braincase volume of the Japanese wolf and Akita was significantly smaller than that of the wolf ssp. ( $P<0.05$ ; Fig. 4A). The medians (means) of the braincase volume to the skull volume in the Japanese wolf, wolf ssp. and Akita were 32.0% (32.7 $\pm$ 1.6%), 26.4% (26.8 $\pm$ 0.7%) and 24.6% (24.2 $\pm$ 2.3%), respectively. The statistical analysis

revealed that the ratio to the braincase volume in the Japanese wolf was significantly larger than that in the wolf ssp. ( $P<0.05$ ). The ratio in the Japanese wolf tended to be larger than that in the Akita ( $P=0.086$ ), although there was no significant difference between the Japanese wolf and Akita (Fig. 4B). There was no significant difference between the median and mean braincase volume and volume ratio.

#### **Frontal sinus**

The medians (means) of the frontal sinus volume in the Japanese wolf, wolf ssp. and Akita were 9.8 ml ( $10.5\pm0.8$  ml), 26.3 ml ( $25.9\pm1.8$  ml) and 29.7 ml ( $28.1\pm5.8$  ml), respectively. According to the statistical analysis, the volume of the frontal sinus in the Japanese wolf was significantly smaller than that in the wolf ssp. ( $P<0.05$ ). The volume in the Japanese wolf tended to be smaller than that in the Akita ( $P=0.055$ ), although there was no significant difference between the Japanese wolf and Akita (Fig. 5A). The medians (means) of the frontal sinus volume ratio to the skull volume in the Japanese wolf, wolf ssp. and Akita were 2.8% ( $2.6\pm0.5\%$ ), 4.8% ( $4.6\pm0.3\%$ ) and 6.5% ( $6.2\pm0.8\%$ ), respectively. The statistical analysis revealed that the frontal sinus volume ratio did not significantly differ among the three groups. However, the ratio in the Japanese wolf was tended to be smaller than that in the other groups (wolf ssp.,  $P=0.054$ ; Akita,

$P=0.086$ ). There was no significant difference between the median and mean braincase volume and volume ratio.

In the morphological observation, in the Akita, the stop, which is a depression (slope) made by an altered angle between the nasal and frontal bones [5], was the deepest among the three groups (Fig. 6A). The frontal bone of the Akita was dorsally directed and well developed with a developed stop, thus the frontal sinus was also well developed dorsally with the development of the frontal bone (Fig. 6 A and C). The frontal sinus of the Japanese wolf did not extend posteriorly to the skull compared with the wolf ssp. and Akita (Figs. 2B and 6B).

#### **Tympanic cavity**

The medians (means) of the tympanic cavity in the Japanese wolf, wolf ssp. and Akita were 4.7 ml ( $4.8\pm0.3$  ml), 10.8 ml ( $10.9\pm0.4$  ml) and 5.4 ml ( $5.5\pm0.4$  ml), respectively. Based on the statistical analysis, the volumes of the tympanic cavity in the Japanese wolf and Akita were significantly smaller than that in the wolf ssp. ( $P < 0.01$ ; Fig. 7A). The medians (means) of the tympanic cavity volume ratio to the skull volume in the Japanese wolf, wolf ssp. and Akita were 1.2% ( $1.2\pm0.0\%$ ), 2.0% ( $2.0\pm0.1\%$ ) and 1.2% ( $1.3\pm0.1\%$ ), respectively. The statistical analysis demonstrated that the ratio of the tympanic cavity volume did not significantly differ between the Japanese wolf and Akita, but the ratio in the wolf ssp. was significantly larger than that in the

other two groups ( $P<0.05$ ; Fig. 7B). There was no significant difference between the median and mean braincase volume and volume ratio.

Furthermore, the tympanic bulla of the wolf ssp. swelled more ventrally than that of the Japanese wolf and Akita (Figs. 1 and 8).

## DISCUSSION

### The volume of the braincase

The braincase is the space that contains the brain, and its shape and size reflect the shape and size of the brain in most mammals [26]. There are many studies of the braincase and its volume in gray wolves. The volume of the braincase was reported to be 115 ml to 175 ml in wild gray wolves [8, 28, 29, 31, 49]. In this study, the medians of the braincase volume of the Japanese wolf and wolf ssp. were 126.4 ml and 149.3 ml, respectively, and the braincase volume was significantly smaller in the Japanese wolf than in the wolf ssp. Compared with previous studies, Japanese wolves comprise a group with a small braincase volume among groups of wild gray wolves.

The braincase volume was examined in several dog breeds and reported as 52.5 ml (toy foxterria) to 166 ml (bull dog) [8, 29, 49]. There is a large difference in body size among dog breeds due to human selection; therefore, a broader range of braincase volumes was observed in

the dog than in the wild wolf. The median braincase volume in the Akita (103.2 ml; range, 96.4–109.4 ml) examined in this study was within the range reported in the above studies. Although there was no significant difference between the Japanese wolf and Akita, the braincase volume of the Japanese wolf (126.4 ml) tended to be larger than that of the Akita (103.2 ml). Compared with other dogs, the braincase volume of the Japanese wolf is considered to be slightly larger than average between maximum and minimum (109.3 ml), although it is small among wild gray wolves.

### **The volume of the frontal sinus**

The frontal sinus, which is one of the paranasal sinuses, is formed by the pneumatization of frontal bones. The frontal sinus is generally thought to play roles in protecting the cranium from mechanical loads and in increasing the attachment sites of the temporal muscles to the cranium without increasing the weight of the cranium [48, 50]. The frontal sinus of the gray wolf is more developed than that of other canids such as the red fox (*Vulpes vulpes*) and raccoon dog (*Nyctereutes procyonoides*) [6]. In gray wolves, the frontal sinus is located widely from the rostral side of the frontal bone to the coronal suture, the boundary between the frontal and parietal bones (Fig. 6B). Curtis and VanValkenburgh [6] reported that the average total frontal sinus volume of both sides was 16 ml (n=4) in wolves. In this study, however, the median frontal sinus volume in

wolf ssp. was 26.3 ml (n=16), being larger than that reported by Curtis and VanValkenburgh [6]. In their study, two of the four wild wolves used in the analysis were Mexican wolves (*C. l. baileyi*), which are said to be the smallest subspecies of wolf that lived in North America. Therefore, this inclusion of small wolves may have caused the frontal sinus volume to be smaller than that of the wolf ssp.

The frontal sinus volume of the Japanese wolf (med. 9.8 ml) was significantly smaller than that of wolf ssp. Moreover, the volume of the Japanese wolf was smaller than that reported by Curtis and VanValkenburgh [6]. As the basal length of the cranium in Japanese wolves is shorter than that in Mexican wolves, the frontal sinus volume of Japanese wolves may have been smaller in absolute value [6, 17, 19].

The median of the frontal sinus volume of the Akita was 29.7 ml, and there was no significant difference between that of the Akita and wolf ssp. The basal cranial length of continental wild wolves is longer than that of Japanese wolves [17, 19], and the total skull length of the Akita is almost the same as that of Japanese wolves [11, 56]. Therefore, the similarity in frontal sinus volume between the Akita and wolf ssp. suggests that the relative frontal sinus volume of the Akita is larger than that of wild wolves, including Japanese wolves, compared with the cranial length.

## **Ratio of braincase and frontal sinus volumes to skull volume**

The relative size of the braincase and frontal sinus varies in relation to the skull size. In wild canids, the braincase is relatively smaller in species with larger skulls, and the species with larger skulls form relatively larger frontal sinuses due to the development of the frontal sinus backwards [6]. Among the vombatiform marsupials, species with larger skulls have a relatively smaller braincase, resulting in relatively larger frontal sinuses. Conversely, species with smaller skulls have a relatively larger cranial cavity, resulting in a smaller posterior extension of the frontal sinus and relatively smaller frontal sinuses [47]. In this study, the ratio of the braincase volume to the skull volume in the Japanese wolf was significantly larger than that in wolf ssp., and the ratio of the frontal sinus volume to the skull volume in Japanese wolves tended to be smaller than that in wolf ssp. The relatively large cranial cavity and small frontal sinus of the Japanese wolf are likely to be features caused by differences in the skull volume between the Japanese wolf and wolf ssp. because the skull volume of the Japanese wolf was significantly smaller than that of wolf ssp. (Figs. 3 and 6).

On the other hand, the Akita, which has a cranial size similar to that of the Japanese wolf, had a smaller braincase volume and larger frontal sinus volume than the Japanese wolf. In general, domestic dogs have several skull features that distinguish them from wolves in wild populations. For example, the frontal sinus of dogs becomes larger than that of wild wolves because the frontal

bone of dogs swells up in the dorsal direction [11]. It is known that the stop between the nasal and frontal bones exhibits a large slope with the swelling of the frontal sinus in the dorsal direction [5, 51] (Fig. 6A and C). In previous studies on the morphology of the skull in the Japanese wolf, the frontal bone did not largely protrude to the dorsal direction and the slope of the stop was gentle [17, 21, 37, 39, 44] (Fig. 6A). The two-dimensional and non-quantitative morphological observations of the frontal sinus in the Japanese wolf and Akita using CT revealed that the Akita develops the frontal sinus dorsally compared with the Japanese wolf [11, 12]. In our three-dimensional analysis, the same pattern was noted (Fig. 6A). In addition, the brain weight to body weight and the braincase volume are smaller in domestic dogs than in wild wolves [29, 49]. Therefore, due to domestication, the Akita developed a relatively small braincase and dorsalized formation of frontal bones with a developed stop, allowing the frontal sinus to extend caudally and dorsally, respectively (Fig. 6), causing the difference in the braincase and frontal sinus volume ratios to the skull between the Japanese wolf and Akita.

### **Tympanic cavity volume and tympanic cavity volume ratio to skull volume**

The tympanic cavity covers most of the middle ear from the ventral side of the skull and fills the inside with air [4, 55]. Gray wolves have a pair of swelled tympanic cavities on the ventral side of the skull. However, domestic dogs have smaller tympanic cavities than other wild



subspecies, and this characteristic can distinguish dogs from wolves [5, 16]. It is therefore suggested that the significant differences in tympanic cavity volume and tympanic cavity volume to skull volume ratio between the wolf ssp. and Akita were caused by domestication.

On the other hand, in this study, the tympanic cavity volume of the Japanese wolf was significantly smaller than that of the wolf ssp. and as small as that of the Akita. The comparison of the tympanic cavity volume to the skull volume ratio between the Japanese wolf and wolf ssp. yielded the same result. Regarding the morphological characteristics of the tympanic cavity in the Japanese wolf, it has been reported that it is shorter in diameter and width than that in Korean wolves [44], and Japanese wolves have ventrally flat tympanic cavities compared with other wild wolves (Fig. 8) [1, 42, 51]. Our quantitative examination of the tympanic cavity volume and morphological observation in the Japanese wolves were consistent with previous studies [1, 15, 18, 19, 37, 39, 44, 51] .

Regarding the tympanic bulla size, studies of crossbreeding between wild wolves and dogs reported that the size and swelling of the tympanic bulla follow the Mendelian laws of inheritance [16]. Regarding the degree of swelling of the tympanic bulla, the swelling of the tympanic bulla at F1 showed intermediate level between wild wolves and dogs, and in the swelling level at F2, the wild wolf, dog and intermediate types were observed. For the reason why the tympanic bulla is small in the Japanese wolf, the hybridization between wild wolves and dogs has been suggested

[16]. However, a recent study that analyzed nDNA of nine Japanese wolves reported that ancient genomic introgression occurred from the ancestors of Japanese wolves to the East Eurasian lineage dogs, and Japanese wolves are genetically closest to dogs [14]. Therefore, the small size of the tympanic bulla in the Japanese wolf may not be due to hybridization with dogs but due to an ancestral trait. Thus, it is possible that the small size of the tympanic bulla in dogs was inherited from the ancestral population of Japanese wolves. However, the specific common morphological features of the Japanese wolf, such as the separated rostral alar foramen and the palatine bone with an anterior notch, are not inherited into dogs. Therefore, the ancestor of Japanese wolves may have acquired the morphological characteristics after diverging East Eurasian lineage dogs.

In conclusion, the present study revealed that the Japanese wolf has a relatively large cranial cavity and small frontal sinus compared with the wolf ssp. and Akita, and that the tympanic cavity volume ratio to the skull volume in the Japanese wolf and Akita was significantly smaller than that in the wolf ssp. In this study, we clarified the characteristics of the internal cranial structures in the Japanese wolf three-dimensionally and quantitatively.

## **CONFLICT OF INTEREST**

The authors have nothing to disclose.

313   **ACKNOWLEDGMENTS**

314   The author would like to thank Dr. Shinichiro Kawada, National Museum of Natural History and  
315   Science, Tokyo, Japan for preparing the valuable Japanese wolf, *Wolf* ssp. and Akita specimens,  
316   and for his cooperation and contributions. This study was financially supported in part by a Grant-  
317   in-Aid for Scientific Research (21K00985) from the Ministry of Education, Culture, Sports,  
318   Science and Technology of Japan.

319

320

321

322

323

324

325

326

327

328

329

330

## REFERENCES

1. Abe Y. 1930. On the Corean and Japanese wolves. *J Sci Hiroshima Univ Ser B Div 1*: 33–37.
2. Badlangana NL, Adams JW, Manger PR. 2011. A comparative assessment of the size of the frontal air sinus in the Giraffe (*Giraffa camelopardalis*). *Anat Rec* **294**: 931–940.
3. Bruner E, Manzi G, Arsuaga JL. 2003. Encephalization and allometric trajectories in the genus Homo: evidence from the Neandertal and modern lineages. *Proc Natl Acad Sci USA* **100**: 15335–15340.
4. Cole LK. 2009. Anatomy and physiology of the canine ear. *Vet Dermatol* **21**: 221–231.
5. Culluton-Brock J. 1995. Origins of the dog: domestication and early history. pp. 7–20. In: The Domestic Dog, Its Evolution, Behavior and Interaction with People (Serpell J. ed.), Cambridge University Press, Cambridge.
6. Curtis AA, Van Valkenburgh B. 2014. Beyond the sniffer: frontal sinuses in Carnivora. *Anat Rec* **297**: 2047–2064.
7. Curtis AA, Orke M, Tetradis S, Van Valkenburgh B. 2017. Diet-related differences in craniodental morphology between captive-reared and wild coyotes, *Canis latrans* (Carnivora: Canidae). *Biol J Linn Soc* **123**: 677–693.
8. Czeibert K, Sommesse A, Petneházy Ö, Csörgö T, Kubinyi E. 2020. Digital endocasting in comparative canine brain morphology. *Front. Vet Sci* **7**: 565315.

- 349 9. Defalque VE, Rosenstein DS, Rosser Jr EJ. 2005. Measurement of normal middle ear cavity  
350 volume in mesaticephalic dogs. *Vet Radiol Ultrasound* **46**: 490–493.
- 351 10. Endo K. 2018. Nihon no ookami no saigo: ookamisake, ookamigari, ookamimatsuri no  
352 hakken [The last time of Japanese wolf: wolf sake, wolf hunting, discovery of wolf festa].  
353 Yama-kei Publishers Co., Ltd., Tokyo (in Japanese).
- 354 11. Endo H, Obara I, Yoshida T, Kuroumaru M, Hayashi Y, Suzuki N. 1997. Osteometrical and  
355 CT examination of the Japanese wolf skull. *J Vet Med Sci* **59**: 531–538.
- 356 12. Endo H, Sakai T, Ito T, Koie Y, Kimura J. 2004. Osteological examination and image analysis  
357 of a cranium of the Japanese wolf found at private house in Yamanashi Prefecture. *J Zoo*  
358 *Wildl Med* **9**: 109–114 (in Japanese with English abstract).
- 359 13. Evans DC. 2005. New evidence on brain-endocranial cavity relationships in ornithischian  
360 dinosaurs. *Acta Palaeontol Pol* **50**: 617–622.
- 361 14. Gojobori J, Arakawa N, Xiayire X, Matsumoto Y, Matsumura S, Hongo H, Ishiguro N, Terai,  
362 Y. 2024. The Japanese wolves are most closely related dogs and share DNA with East  
363 Eurasian dogs. *Nat Commun* (In Press)
- 364 15. Hiraiwa Y. 1992. Ookami– sono seitai to rekishi– [The Wolf: The Ecology and History].  
365 Tsukiji Shokan Publishing Co., Ltd., Tokyo (in Japanese).
- 366 16. Iljin NA. 1941. Wolf-dog genetics. *J Gene* **42**: 359–414.

- 367 17. Imaizumi Y. 1970. Systematic status of the extinct Japanese wolf, *Canis hodophilax*: 1.  
368 Identification of specimens. *J Mammal Soc Jpn* **5**: 27–32 (in Japanese with English abstract).
- 369 18. Imaizumi Y. 1970. Systematic status of the extinct Japanese wolf, *Canis hodophilax*: 2.  
370 Similarity relationship of *hodophilax* among species of the genus *Canis*. *J Mammal Soc Jpn*  
371 **5**: 62–66 (in Japanese with English abstract).
- 372 19. Imaizumi Y. 1988. Honyuudoubutsu sinkaron: honyuurui no shu to shubunka [The theory of  
373 mammal evolution]. Newton Press Publishing Co., Ltd., Tokyo (in Japanese).
- 374 20. Ishiguro, N. 2012. Phylogenetic analysis of extinct wolves in Japan. *J Jpn Vet Med Assoc* **65**:  
375 225–231 (in Japanese).
- 376 21. Ishiguro N, Inoshima Y, Sasaki M. 2017. Computed tomography examination and  
377 mitochondrial DNA analysis of Japanese wolf skull covered with skin. *J Vet Med Sci* **79**: 14–  
378 17.
- 379 22. Ishiguro N, Inoshima Y, Shigehara N. 2009. Mitochondrial DNA analysis of the Japanese  
380 wolf (*Canis lupus hodophilax* Temminck, 1819) and comparison with representative wolf and  
381 domestic dog haplotypes. *Zool Sci* **26**: 765–770.
- 382 23. Ishiguro N, Inoshima Y, Yanai T, Sasaki M, Matsui A, Kikuchi H, Maruyama M, Hongo H,  
383 Vostretsov YE, Gasilin V, Kosintsev PA, Quanjia C, Chunxue W. 2016. Japanese wolves are  
384 genetically divided into two groups based on an 8-nucleotide insertion/deletion within the

385 mtDNA control region. *Zool Sci* **33**: 44–49.

386 24. Ishiguro N, Matsumura S, Terai Y, Hongo H. 2021. The mystery of Japanese wolves called  
387 ookami or yamainu in the Siebold collection. *J Jpn Vet Med Assoc* **74**: 389–395 (in Japanese).

388 25. Iurino DA, Mecozzi B, Iannucci A, Moscarella A, Strani F, Bone F, Gaeta M, Sardella R.  
389 2022. A Middle Pleistocene wolf from central Italy provides insights on the first  
390 occurrence of *Canis lupus* in Europe. *Sci Rep* **12**: 2882.

391 26. Kawabe S. 2015. Fossil research using computed tomography scanner: Introduction to  
392 paleoneurology. *Earth Sci* **69**: 241–248 (in Japanese).

393 27. Kawabe S, Ando T, Endo H. 2014. Enigmatic affinity in the brain morphology between  
394 plotopterids and penguins, with a comprehensive comparison among water birds. *Zool J Linn*  
395 *Soc* **170**: 467–493.

396 28. Kawabe S, Shimokawa T, Miki H, Okamoto T, Matsuda S, Itou T, Koie H, Kitagawa M,  
397 Sakai T, Hosojima M, Endo H. 2013. Relationship between brain volume and brain width in  
398 mammals and birds. *Paleontol Res* **17**: 282–293.

399 29. Kruska DC. 2005. On the evolutionary significance of encephalization in some eutherian  
400 mammals: effects of adaptive radiation, domestication, and feralization. *Brain Behav Evol*  
401 **65**: 73–108.

- 402 30. Lyras GA, Van der Geer AA, Dermitzakis MD. 2001. Evolution of the brain of Plio-  
403 Pleistocene wolves. *Cranium* **18**: 30–40.
- 404 31. Lyras GA. 2009. The evolution of the brain in Canidae (Mammalia: Carnivora). *Scripta Geol*  
405 **139**: 1–93.
- 406 32. Macdonald DW, Sillero-Zubri C. 2004. Dramatis personae: wild canids-an introduction and  
407 dramatis personae. pp 3–36, In: Biology and Conservation of Wild Canids (Macdonald DW,  
408 Sillero-Zubri C eds.), Oxford University Press, Oxford.
- 409 33. Malkemper EP, Mason MJ, Burda H. 2020. Functional anatomy of the middle and inner ears  
410 of the red fox, in comparison to domestic dogs and cats. *J Anat* **236**: 980–995.
- 411 34. Matsumura S, Inoshima Y, Ishiguro N. 2014. Reconstructing the colonization history of lost  
412 wolf lineages by the analysis of the mitochondrial genome. *Mol Phylogenet Evol* **80**: 105–  
413 112.
- 414 35. Matsumura S, Terai Y, Hongo H, Ishiguro N. 2021. Analysis of the mitochondrial genome of  
415 Japanese wolf specimens in the Siebold collection, Leiden. *Zool Sci* **38**: 60–66.
- 416 36. Mech LD. 1974. *Canis lupus*. *Mammalian Species* **37**: 1–6.
- 417 37. Miyamoto F, Maki I. 1982. On the repaired specimen of Japanese wolf (*Canis lupus*  
418 *hodophilax* TEMMINCK) and its skull newly taken out. *Bull Fac Ed Wakayama Univ Nat Sci*  
419 **32**: 9–16 (in Japanese with English abstract).



- 420 38. Naora N. 1965. On Wolves in Japan. Azakura-shobo, Tokyo (in Japanese with English  
421 abstract).
- 422 39. Obara I. 1990. Skull of Japanese wolf, *Canis hodophilax*, preserved as old private houses in  
423 Atsugi—shi and Kiyokawa—mura, Kanagawa Prefecture. *Nat Hist Rep Kanagawa* **11**: 53–  
424 65 (in Japanese).
- 425 40. O'Keefe FR, Binder WJ, Frost SR, Sadlier RW, Van Valkenburgh B. 2014. Cranial  
426 morphometrics of the dire wolf, *Canis dirus*, at Rancho La Brea: temporal variability and its  
427 links to nutrient stress and climate. *Palaeontol Electronica* **17**: 17A.
- 428 41. Pang J, Kluetsch C, Zou X, Zhang A, Luo L, Angleby H, Ardalan A, Ekström C, Skölleremo  
429 A, Lundeberg J, Matsumura S, Leitner T, Zhang Y, Savolainen P. 2009. mtDNA data indicate  
430 a single origin for dogs south of Yangtze River, less than 16,300 years ago, from numerous  
431 wolves. *Mol Biol Evol* **26**: 2849–2864.
- 432 42. Pocock RI. 1935. The races of *Canis lupus*. *J Zool* **105**: 658–689.
- 433 43. Rosowski JJ, Ravicz ME, Teoh SW, Flandermeyer D. 1999. Measurements of middle-ear  
434 function in the Mongolian Gerbil, a specialized mammalian ear. *Audiol Neurotol* **4**: 129–  
435 136.
- 436 44. Saito H. 1938. Tokyo Kagaku Hakubutsukan Souko ni Hakken Seraretaru Yamainu no  
437 Zenshin Kokkaku narabini Sonohoka no Doushiryou ni tsuite [Skeletal specimen of yamainu

438 and other data discovered at National Museum of Natural History and Science, Tokyo].

439 *Hakubutuan Kenkyu* **11**: 2–7 (in Japanese).

440 45. Sakai ST, White B, Arsznov BM, Lundrigan BL. 2018. Endocranial development in the

441 Coyote (*Canis latrans*) and Gray Wolf (*Canis lupus*): a computed tomographic study. *Brain*

442 *Behave Evol* **91**: 65–81.

443 46. Savolainen P, Zhang Y -P, Luo J, Lundeberg J, Leitner T. 2002. Genetic evidence for an East

444 Asian origin of domestic dogs. *Science* **298**: 1610–1613.

445 47. Sharp AC. 2016. A quantitative comparative analysis of the size of the frontoparietal sinuses

446 and brain in vombatiform marsupials. *Mem Mus Vic* **74**: 331–342.

447 48. Sharp AC, Rich TH. 2016. Cranial biomechanics, bite force and function of the endocranial

448 sinuses in *Diprotodon optatum*, the largest known marsupial. *J Anat* **228**: 984–995.

449 49. Smith BP, Lucas TA, Norris M, Henneberg M. 2018. Brain size/body weight in the dingo

450 (*Canis dingo*): comparisons with domestic and wild canids. *Aust J Zool* **65**: 292–301.

451 50. Tanner JB, Dumont ER, Sakai ST, Lundrigan BL, Holekamp KE. 2008. Of arcs and vaults:

452 the biomechanics of bone-cracking in spotted hyenas (*Crocuta crocuta*). *Biol J Linn Soc* **95**:

453 246–255.

454 51. Tasumi M. 1991. Reality of Japanese wolf as investigated from the skull remains. *The Bone*

455 **6**: 119–128 (in Japanese).

52. von den Driesch A. 1976. A Guide to the Measurement of Animal Bones from Archaeological Sites. Harvard University press, Massachusetts.
53. Vonholdt BM, Pollinger JP, Lohmueller KE, Han E, Parker HG, Quinon P, Degenhardt JD, Boyko AR, Earl DA, Auton A, Reynolds A, Bryc K, Brisbin A, Knowles JC, Mosher DS, Spady TC, Elkahouloun A, Geffen E, Pilot M, Jedrzejewski W, Greco C, Randi E, Bannasch D, Wilton A, Shearman J, Musiani M, Cargill M, Jones PG, Qian Z, Huang W, Ding ZL, Zhang YP, Bustamante CD, Ostrander EA, Novembre J, Wayne RK. 2010. Genome-wide SNP and haplotype analyses reveal a rich history underlying dog domestication. *Nature* **464**: 898–902.
54. Walker BL. 2005. The Lost Wolves of Japan. University of Washington Press, Washington.
55. Wang X, Tedford RH. 2008. Dogs: Their Fossil Relatives and Evolutionary History, Columbia University Press, New York.
56. Yoshida T, Endo H, Kurohmaru M, Hayashi Y. 1999. Identification of the Japanese wolf skull in the 3D morphometrical method. *Honyurui Kagaku* **39**: 239–246 (in Japanese with English abstract).

## FIGURE LEGENDS AND TABLE CAPTIONS

**Fig. 1.** Characteristics of the skull of Japanese wolves different from the other subspecies of gray wolves. A, Left lateral view of skull; B, Ventral view of skull; 1, Japanese wolf (M1185); 2, continental wild gray wolf (M31475). Japanese wolves have two rostral alar foramens (A-1), whereas the other wolves, including domestic dogs, have only one foramen (A-2). The mid-caudal end of the horizontal plate of palatine bone is indented rostrally in Japanese wolves (B-1), whereas that of the other wolves protrudes caudally or flat (B-2).

**Fig. 2.** CT images of the braincase (A), frontal sinus (B) and tympanic cavity (C) within the skull. The skull of the Japanese wolf (Nara) is used in this figure. A and B, dorsal view; C, ventral view.

**Fig. 3.** The representative values of the skull volume in three groups of gray wolves. Asterisk indicates a significant difference. ( $P < 0.05$ , Steel-Dwass test).

**Fig. 4.** The representative values of the braincase volume (A) and the ratio of the braincase volume to the skull volume (B) in three groups of gray wolves. Asterisks indicate a significant difference (\*  $P < 0.05$ , Steel-Dwass test).

**Fig. 5.** The representative values of the frontal sinus volume (A) and the ratio of the frontal sinus volume to the skull volume (B) in three groups of gray wolves. Asterisks indicate a significant difference (\*  $P < 0.05$ , Steel-Dwass test).

**Fig. 6.** Comparison of the morphology of frontal sinuses. Left lateral (A; Frontal sinus, braincase and tympanic cavity are superposed.), dorsal (B; Frontal sinus and braincase are superposed.) and rostral (C; Only frontal sinus) views of CT images of the frontal sinus within the skull in the Japanese wolf (M26696), Wolf ssp. (M31475) and Akita (M18303). White arrows indicate significant differences macroscopically from the Japanese wolf. A red arrow shows the stop. The size is matched by condylobasal length.

**Fig. 7.** The representative values of the tympanic cavity volume (A) and the ratio of the tympanic cavity volume to the skull volume (B) in three groups of gray wolves. Asterisks indicate a significant difference (\*  $P < 0.05$ , \*\*  $P < 0.01$ , Steel-Dwass test).

**Fig. 8.** Comparison of the morphology of the tympanic cavity in gray wolves. Left lateral (A) and ventral (B) views of the tympanic cavity in the Japanese wolf (Nara), Wolf ssp.

(M25791) and Akita (M19301). The size of all specimens is matched by condylobasal length.

The upper red dotted line runs through the ventral end of the external acoustic pore in all

wolf groups, and a lower line is drawn on the ventral end of the tympanic cavity in Wolf ssp.

**Table 1.** Specimens used in this study

ID, specimen number; UKN, unknown.

Shimane and Nara are shown as prefecture names where these specimens are preserved

because there are no specimen numbers.

**Table 2.** Descriptions of cranial measurement points

\* Skulls were measured by the methods of von den Driesch [52].

**Table 3.** Linear measurements of each skull with a caliper

ID, specimen number; NA, not analyzed. Measurement points are shown by abbreviations.

The relationship between measurement points and abbreviations is described in Table 2.

\* Shimane was covered with dry soft tissue, so this specimen was measured on the medical

workstation for image processing (ziocube, Tokyo, Japan).

528     **Table 4.** The volume of each cavity and the ratio of each cavity volume to the skull volume

529             ID, specimen number; NA, not analyzed. L, left side; R, right side.

Table 1. Specimens used in this study

ID	Subspecies	Group name	SEX	Facilities of preservation
M1185	<i>Canis lupus hodophilax</i>	Japanese wolf	UKN	National museum of nature and science, Tokyo
M26696	<i>Canis lupus hodophilax</i>	Japanese wolf	UKN	National museum of nature and science, Tokyo
Shimane	<i>Canis lupus hodophilax</i>	Japanese wolf	UKN	Private
Nara	<i>Canis lupus hodophilax</i>	Japanese wolf	UKN	Ooyodo-cho town official board of education
M17278	<i>Canis lupus</i> ssp.	Wolf ssp.	Male	National museum of nature and science, Tokyo
M18304	<i>Canis lupus</i> ssp.	Wolf ssp.	Male	National museum of nature and science, Tokyo
M31474	<i>Canis lupus</i> ssp.	Wolf ssp.	Male	National museum of nature and science, Tokyo
M31475	<i>Canis lupus</i> ssp.	Wolf ssp.	Male	National museum of nature and science, Tokyo
M31476	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	National museum of nature and science, Tokyo
M25791	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	National museum of nature and science, Tokyo
M25792	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	National museum of nature and science, Tokyo
M25793	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	National museum of nature and science, Tokyo
M25794	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	National museum of nature and science, Tokyo
M25795	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	National museum of nature and science, Tokyo
M25796	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	National museum of nature and science, Tokyo
M43287	<i>Canis lupus</i> ssp.	Wolf ssp.	Female	National museum of nature and science, Tokyo
KPM-NFM 2014	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	Kanagawa prefectural museum of natural science
KPM-NFM 2015	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	Kanagawa prefectural museum of natural science
KPM-NFM 2303	<i>Canis lupus</i> ssp.	Wolf ssp.	Male	Kanagawa prefectural museum of natural science
KPM-NFM 2913	<i>Canis lupus</i> ssp.	Wolf ssp.	UKN	Kanagawa prefectural museum of natural science
M3879	<i>Canis lupus familiaris</i>	Akita	Male	National museum of nature and science, Tokyo
M12921	<i>Canis lupus familiaris</i>	Akita	Male	National museum of nature and science, Tokyo
M18303	<i>Canis lupus familiaris</i>	Akita	Male	National museum of nature and science, Tokyo
M19301	<i>Canis lupus familiaris</i>	Akita	Male	National museum of nature and science, Tokyo

ID, specimen number; UKN, unknown.

Shimane and Nara are shown as prefecture names where these specimens are preserved because there are no specimen numbers.



Table 2. Measurement points in skulls

Measurments *	Abbreviations
Total length: Akrokranium - Prosthion	TL
Basal length: Basion - Synsphenion	BL
Zygomatic breadth: Zygion - Zygion	ZB
Skull height. The two pointers of the slide gauge are placed basally on the basis of the skull (on the basioccipital) and dorsally on the highest elevation of the sagittal crest.	SH

\* Skulls were measured by the methods of von den Driesch [52].

Table 3. Linear measurements of each skull with a caliper

ID	Group name	Measurement points			
		TL	BL	ZB	SH
M1185	Japanese wolf	204.3	187.2	NA	64.0
M26696	Japanese wolf	202.5	176.7	116.9	62.5
Shimane*	Japanese wolf	226.9	197.2	128.0	NA
Nara	Japanese wolf	212.5	188.4	116.9	63.7
M17278	Wolf ssp.	221.2	193.2	120.6	66.8
M18304	Wolf ssp.	251.2	226.7	133.7	76.3
M31474	Wolf ssp.	245.8	217.0	134.0	71.3
M31475	Wolf ssp.	247.6	214.4	124.9	72.1
M31476	Wolf ssp.	254.7	221.4	136.4	75.0
M25791	Wolf ssp.	260.3	227.2	141.6	74.3
M25792	Wolf ssp.	246.7	220.3	130.1	71.2
M25793	Wolf ssp.	246.5	NA	135.9	NA
M25794	Wolf ssp.	229.1	210.9	117.8	58.3
M25795	Wolf ssp.	234.4	206.2	131.0	63.4
M25796	Wolf ssp.	258.2	225.9	129.5	72.2
M43287	Wolf ssp.	239.4	212.1	134.0	62.6
KPM-NF1 2014	Wolf ssp.	266.6	235.9	143.0	76.2
KPM-NF1 2015	Wolf ssp.	271.6	235.2	146.0	78.0
KPM-NF1 2303	Wolf ssp.	245.9	217.7	127.3	73.9
KPM-NF1 2913	Wolf ssp.	246.3	221.2	133.8	79.6
M3879	Akita	224.3	197.2	133.1	71.9
M12921	Akita	223.2	200.6	125.0	70.7
M18303	Akita	214.0	193.8	120.6	67.3
M19301	Akita	201.9	184.3	116.3	62.8

ID, specimen number; NA, not analyzed.

Measurement points are shown by abbreviations (see Table 2).

The relationship between measurement points and abbreviations is described in Table 2.


\*: Shimane was covered with dry soft tissue, so this specimen was measured on the medical workstation for image processing (ziocube, Tokyo, Japan) .

Table 4. The volume of each cavity and the ratio of each cavity volume to the skull volume


ID	Group name	Skull	Braincase		Frontal sinus				Tympanic cavity			
		Volume (ml)	Volume (ml)	Ratio (%)	Volume (L) (ml)	Volume (R) (ml)	Volume (L+R) (ml)	Ratio (L+R) (%)	Volume (L) (ml)	Volume (R) (ml)	Volume (L+R) (ml)	Ratio (L+R) (%)
M1185	Japanese wolf	343.1	122.4	35.7	4.8	4.7	9.5	2.8	1.97	2.29	4.26	1.24
M26696	Japanese wolf	NA	100.5	NA	4.1	6.0	10.1	NA	2.35	2.33	4.68	NA
Shimane	Japanese wolf	457.3	139.0	30.4	6.0	7.0	13.0	2.8	2.69	2.82	5.51	1.20
Nara	Japanese wolf	407.0	130.4	32.0	5.0	4.5	9.5	2.3	2.27	2.39	4.66	1.14
M17278	Wolf ssp.	425.8	130.3	30.6	13.1	13.7	26.8	6.3	4.54	4.47	9.01	2.12
M18304	Wolf ssp.	623.3	164.4	26.4	19.1	19.0	38.2	6.1	4.92	5.80	10.72	1.72
M31474	Wolf ssp.	433.7	134.1	30.9	7.7	7.2	14.8	3.4	4.40	4.46	8.86	2.04
M31475	Wolf ssp.	471.6	136.4	28.9	14.7	14.6	29.3	6.2	4.76	4.49	9.25	1.96
M31476	Wolf ssp.	633.2	159.2	25.1	12.7	12.8	25.5	4.0	6.59	6.59	13.18	2.08
M25791	Wolf ssp.	479.5	144.2	30.1	13.1	12.6	25.7	5.4	5.15	5.03	10.18	2.12
M25792	Wolf ssp.	571.1	149.1	26.1	11.6	11.1	22.7	4.0	6.14	5.94	12.08	2.12
M25793	Wolf ssp.	589.1	164.7	28.0	6.4	6.6	13.0	2.2	6.12	6.46	12.58	2.14
M25794	Wolf ssp.	583.4	143.1	24.5	17.6	19.1	36.7	6.3	6.52	6.52	13.04	2.24
M25795	Wolf ssp.	555.9	148.0	26.6	12.0	14.9	26.9	4.8	5.33	5.54	10.87	1.96
M25796	Wolf ssp.	NA	157.0	NA	10.2	10.1	20.3	NA	4.84	4.73	9.57	NA
M43287	Wolf ssp.	505.7	132.3	26.2	9.1	9.0	18.1	3.6	4.02	3.95	7.97	1.58
KPM-NFM 2014	Wolf ssp.	723.6	163.7	22.6	15.7	14.6	30.2	4.2	5.54	5.54	11.08	1.53
KPM-NFM 2015	Wolf ssp.	740.8	175.5	23.7	11.8	14.0	25.8	3.5	4.99	5.17	10.16	1.37
KPM-NFM 2303	Wolf ssp.	542.6	149.4	27.5	13.6	13.4	27.0	5.0	7.43	6.57	14.00	2.58
KPM-NFM 2913	Wolf ssp.	641.0	154.6	24.1	16.7	17.0	33.7	5.3	5.35	5.75	11.10	1.73
M3879	Akita	501.1	109.4	21.8	19.2	19.6	38.8	7.7	3.11	3.43	6.54	1.31
M12921	Akita	530.6	99.5	18.8	15.4	16.5	31.9	6.0	2.89	2.91	5.80	1.09
M18303	Akita	390.3	106.8	27.4	13.1	14.4	27.4	7.0	2.10	2.44	4.54	1.16
M19301	Akita	335.7	96.4	28.7	6.8	7.4	14.2	4.2	2.59	2.48	5.07	1.51

ID, specimen number; NA, not analyzed.


L, left side; R, right side.




**Fig. 1.** Characteristics of the skull of Japanese wolves different from the other subspecies of gray wolves. A, Left lateral view of skull; B, Ventral view of skull; 1, Japanese wolf (M1185); 2, continental wild gray wolf (M31475). Japanese wolves have two rostral alar foramina (A-1), whereas the other wolves, including domestic dogs, have only one foramen (A-2). The mid-caudal end of the horizontal plate of palatine bone is indented rostrally in Japanese wolves (B-1), whereas that of the other wolves protrudes caudally or flat (B-2).




**Fig. 2.** CT images of the braincase (A), frontal sinus (B) and tympanic cavity (C) within the skull. The skull of the Japanese wolf (Nara) is used in this figure. A and B, dorsal view; C, ventral view.



**Fig. 3.** The representative values of the skull volume in three groups of gray wolves. Asterisk indicates a significant difference. ( $P < 0.05$ , Steel-Dwass test).




**Fig. 4.** The representative values of the braincase volume (A) and the ratio of the braincase volume to the skull volume (B) in three groups of gray wolves. Asterisks indicate a significant difference (\*  $P < 0.05$ , Steel-Dwass test).




**Fig. 5.** The representative values of the frontal sinus volume (A) and the ratio of the frontal sinus volume to the skull volume (B) in three groups of gray wolves. Asterisks indicate a significant difference (\*  $P < 0.05$ , Steel-Dwass test).






**Fig. 6.** Comparison of the morphology of frontal sinuses. Left lateral (A; Frontal sinus, braincase and tympanic cavity are superposed.), dorsal (B; Frontal sinus and braincase are superposed.) and rostral (C; Only frontal sinus) views of CT images of the frontal sinus within the skull in the Japanese wolf (M26696), Wolf ssp. (M31475) and Akita (M18303). White arrows indicate significant differences macroscopically from the Japanese wolf. A red arrow shows the stop. The size is matched by condylobasal length.



**Fig. 7.** The representative values of the tympanic cavity volume (A) and the ratio of the tympanic cavity volume to the skull volume (B) in three groups of gray wolves. Asterisks indicate a significant difference (\*  $P < 0.05$ , \*\*  $P < 0.01$ , Steel-Dwass test).



**Fig. 8.** Comparison of the morphology of the tympanic bulla in gray wolves. Left lateral (A) and ventral (B) views of the tympanic bulla in the Japanese wolf (Nara), Wolf ssp. (M25791) and Akita (M19301). The size of all specimens is matched by condylobasal length. The upper red dotted line runs through the ventral end of the external acoustic pore in all wolf groups, and a lower line is drawn on the ventral end of the tympanic bulla in Wolf ssp.