

RESEARCH

Open Access



# First report of *Angiostrongylus vasorum* in an African golden wolf (*Canis lupaster*) in Algeria

Noureddine Mechouck<sup>1†</sup>, Georgiana Deak<sup>1\*†</sup>, Angela Monica Ionică<sup>1,2</sup>, Corina Toma<sup>3</sup>, Andrada Gabriela Negoescu<sup>3</sup>, Marian Taulescu<sup>3</sup>, Zihad Bouslama<sup>4</sup> and Andrei Daniel Mihalca<sup>1</sup>

## Abstract

**Background** *Angiostrongylus vasorum*, commonly known as the “French heartworm,” is a nematode belonging to the Metastrongyloidea superfamily. This parasite was first identified in Toulouse, France in 1853 infecting the pulmonary arteries and the right side of the heart of a Pointer dog. Angiostrongylosis is an important infection due to its severe clinical signs and potential for causing high morbidity and mortality in domestic dogs. This nematode has not been studied in Algeria. The aim of this study was to investigate the presence of lungworms among different mammal species in a number of Algerian regions.

**Methods** Between February 2022 and September 2023, 47 road-killed animals were collected from six administrative units (departments) in Algeria. All carcasses underwent a full parasitological necropsy, and lung tissues were preserved in 10% buffered formalin and concentrated ethanol for further study. All collected samples were subjected to histological and PCR (cytochrome c oxidase subunit 1 gene) analyses for lungworm identification.

**Results** Histological examination revealed the presence of nematode eggs and larvae in the alveolar space and chronic obstructive vascular changes were detected in a single golden African wolf (*Canis lupaster*) collected from the department of Constantine. First-stage larvae were collected and morphologically identified as *Angiostrongylus* spp. The molecular identification confirmed the presence of *A. vasorum*. All other animals tested were negative for lungworms.

**Conclusions** To the best of our knowledge, this is the first report of *A. vasorum* infection in an African golden wolf (*Canis lupaster*). We report a new host association, highlighting the importance of further studies to update the geographical distribution of *A. vasorum* and its epidemiology across Algeria.

**Keywords** *Angiostrongylus vasorum*, Golden African wolf (*Canis lupaster*), Algeria

<sup>†</sup>Noureddine Mechouck and Georgiana Deak contributed equally to this work and share first authorship.

\*Correspondence:

Georgiana Deak  
georgiana.deak@usamvcluj.ro

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

*Angiostrongylus vasorum*, commonly known as the “French heartworm”, was first identified in Toulouse, France by Serres in 1853 infecting the pulmonary artery and the right side of the heart of a Pointer dog [1]. Since then, the parasite has been described across a wide range of countries, in Europe, the Americas and Africa [2]. In the last two decades, the parasite gained much interest among researchers due to the severe clinical features and acute course of the infection in domestic dogs and its rapid expansion in Europe [3]. *Angiostrongylus vasorum* is the causative agent of angiostrongylosis in canids, which can present as various clinical signs, ranging from subclinical to lethal, with the most commonly observed symptoms including breathing difficulty, heart insufficiency, coagulation disorder, anorexia and decreased physical endurance. Clinical neurological signs are also common and can be manifested as ataxia, paresis, paralysis or seizures [4–6]. Unusual localizations of adult nematodes have been reported in the anterior chamber of the eye [7, 8], femoral artery [9], pericardial sac and urinary bladder [10], and larval stages have been identified in the brain [11], diaphragm, pancreas, liver, and skin [10].

This nematode has an indirect life-cycle using gastropods (snails and slugs) as intermediate hosts, while small vertebrates can act as paratenic hosts [6, 12–14]. The domestic chicken *Gallus domesticus* was shown to be a potentially suitable paratenic host in studies involving artificial infections [15]. The primary transmission route of *A. vasorum* is through the ingestion of third-stage larvae (L3) by the intermediate host (intentionally or accidentally) and/or a paratenic host. Infection can occur when the final host consumes food contaminated with infective intermediate hosts [4, 16] or water contaminated with L3 [17]. There is also the possibility of infection through the consumption of free L3 released in snail slime [17, 18].

Among wild animals, wild foxes (*Vulpes vulpes*) commonly serve as the typical definitive hosts for *A. vasorum* [19], although there have been multiple reports of other wild canid definitive host species, such as the crab-eating fox (*Cerdocyon thous*), hoary fox (*Lycalopex vetulus*), coyote (*Canis latrans*), golden jackal (*Canis aureus*), gray wolf (*Canis lupus*) and raccoon dog (*Nyctereutes procyonoides*) [4, 12, 20–24]. Non-canid wild captive animals like the red panda (*Ailurus fulgens fulgens*) [25] and meerkats (*Suricata suricatta*) [26] have also been confirmed as definitive hosts. All of these hosts have the potential to act as reservoirs of *A. vasorum* for the infection for domestic dogs (*Canis familiaris*) [4, 27] and cats (*Felis catus*) [28].

In Africa, the first *A. vasorum* infection was documented in Uganda in 1972, in five necropsied domestic dogs [29]; more recently, a potential autochthonous case in a 6-month-old asymptomatic dog was reported in Morocco based on the morphology of first-stage larvae (L1) [30].

There are 100 mammalian species in Algeria, belonging to 37 families and 11 orders, among which carnivores account for 21 species and seven families [31]. The diets of these species are mainly obtained by scavenging and/or predation, making these animals susceptible to food-borne parasitic infections [32]. Considering the abundant wild fauna in Algeria and the occurrence of the parasite in other northern African countries, the aim of this study was to investigate the presence of *A. vasorum* in animal hosts in Algeria.

## Methods

Between February 2022 and September 2023, the carcasses of 47 road-killed animals belonging to the orders Carnivora (11 *Canis familiaris*, 9 *Canis lupaster*, 6 *Felis catus*, 1 *Genetta genetta*, 3 *Herpestes ichneumon*, 3 *Vulpes vulpes* and 13 *Vulpes zerda*) and Rodentia (1 *Hystrix cristata*) were collected from 11 localities and six administrative units (departments) in Algeria (Table 1; Fig. 1) and transported to the National Research Center (CRE) in Annaba according to national governmental provisions. Prior to examination, the carcasses were entirely enclosed in a plastic bag (with identification) for health safety reasons and stored at - 20 °C. An extensive parasitological necropsy was performed on each carcass, and comprehensive data on each animal's age and sexual maturity were collected according to [33, 34]. The entire cardio-respiratory system was removed from all animals, and the trachea, bronchi and bronchioles as well as the pulmonary arteries and the heart chambers were longitudinally opened and carefully examined for parasites using a stereomicroscope as described in [35]. During necropsy, macroscopic pictures of the lungs were taken where lesions were observed, samples of lung tissue were collected from each animal and preserved in both 10% buffered-formalin and concentrated ethanol separately. The collected samples were then legally transported to the Faculty of Veterinary Medicine (University of Agriculture and Veterinary Medicine) of Cluj-Napoca for molecular and histological analyses.

The formalin-fixed lung tissue samples were trimmed and embedded in paraffin wax, according to standard protocols, following which 2- to 3-µm-thick sections were obtained and stained with hematoxylin and eosin (H&E). The histological assessment was performed using a light Olympus BX-41 microscope equipped with an Olympus SP 350 digital camera (Olympus Corp.,


**Table 1** Prevalence of lungworms in various animal species across different departments and localities

Department	Locality	Animal species	Number of examined animals	Animals positive for lungworms (N, %)
El Tarf	Ain kerma	<i>Canis familiaris</i>	6 (3 M, 3 F)	0
		<i>Canis lupaster</i>	1 M	0
		<i>Herpestes ichneumon</i>	1 M	0
	Asfour	<i>Canis lupaster</i>	1 F	0
	Bouhadjar	<i>Hystrix cristata</i>	1 M	0
		<i>Canis lupaster</i>	1 M	0
		<i>Vulpes vulpes</i>	2 (1 M, 1 F)	0
	Drean	<i>Felis catus</i>	1 M	0
		<i>Herpestes ichneumon</i>	1 M	0
		<i>Canis fammiliaris</i>	1 F	0
	Zitouna	<i>Genetta genetta</i>	1 F	0
		<i>Canis lupaster</i>	1 F	0
Annaba	Annaba	<i>Canis familiaris</i>	1 F	0
		<i>Canis lupaster</i>	2 (1 M, 1 F)	0
		<i>Felis catus</i>	4 (1 M, 3 F)	0
		<i>Herpestes ichneumon</i>	1 M	0
		<i>Canis Lupaster</i>	1 M	0
Constantine	Seraidi	<i>Canis Lupaster</i>	1 M	0
	Constantine	<i>Canis lupaster</i>	1 M	<i>Angios-trongylus vasorum</i> (1, 2.13%)
Skikda	Skikda	<i>Vulpes vulpes</i>	1 F	0
		<i>Felis catus</i>	1 M	0
		<i>Canis lupaster</i>	1 M	0
Algiers	Staoueli	<i>Canis familiaris</i>	2 (1 M, 1F)	0
Oued Souf	Oued souf	<i>Vulpes zerda</i>	13 (7 M, 6F)	0
Total			<b>47 (24 M, 23F)</b>	<b>1, 2.13%</b>

F Female, M male

Tokyo, Japan). The photomicrographs were taken using the Stream Basic imaging software (Olympus Corp.). When the histological examination showed a parasitic infection, the entire quantity of formalin in which lungs were conserved was placed in a 15-ml conical centrifuge tube and centrifuged (13,000g); the obtained sediment was examined under a stereomicroscope. One drop of Lugol solution was added directly onto the slide for a better view of larvae while taking pictures. Genomic DNA was isolated from small lung sections conserved in concentrated ethanol from all animals using a commercial kit (Isolate II Genomic DNA kit; Meridian Bioscience, London, UK) according to the manufacturer's instructions. DNA was also isolated from a pool of larvae collected from the ethanol solution in which the African wolf lung tissues were conserved. A fragment of the mitochondrial cytochrome *c* oxidase

subunit 1 (cox1, approx. 700 bp) gene was then amplified by conventional PCR using the universal primers LCO1490/HCO2198, according to protocols reported in the literature [36]. The obtained PCR products were visualized following electrophoresis in 2% agarose gels stained with EcoSafe nucleic acid staining solution (Pacific Image Electronics, Taiwan), and their molecular weight was assessed by comparison to a molecular marker (HyperLadder™ 100 bp, meridian Bioscience, UK). The obtained band was excised from the gel, purified on a silica membrane spin column (Gel/PCR DNA Fragments Kit; Geneaid Biotech, New Taipei City, Taiwan), and sequenced bidirectionally using an external service (performed by Macrogen Europe B.V., Amsterdam, the Netherlands). The sequences were assembled using Geneious software (Biomatters Ltd, Auckland, New Zealand) and compared to other sequences available in the NCBI GenBank® database by Basic Local Alignment Search Tool (BLAST) analysis.




**Fig. 1** Map showing the geographical locations (departments) of the tested animals and the *Angiostrongylus vasorum*-positive animal

## Results

At the time of the necropsy, adult lungworms were not detected in any of the examined animals. In one single African golden wolf (*Canis lupaster*) collected from Constantine department (36°21'N 6°36'E), the pulmonary parenchyma showed moderate congestion and edema, multifocal dark-red and variably-sized

areas of densification, compatible with foci of ischemic necrosis, severe hemorrhage and interstitial verminous pneumonia (Fig. 2a, b). The lesions were mainly located at the periphery of the lung parenchyma. No adult parasites were observed within the right ventricle and pulmonary arteries during the macroscopical and microscopical examination. The lungs of all other



**Fig. 2** Gross evaluation of the lungs of an African golden wolf (*Canis lupaster*) with suspected *Angiostrongylus vasorum* infection. **a, b** Pulmonary parenchyma showing multifocal to coalescing dark-red areas (arrows) consistent with verminous pneumonia and severe alteration of blood vessels




examined animals had no serious macroscopic lesions. In the African golden wolf with pulmonary lesions, histological examination revealed numerous transverse to longitudinal sections of nematode larvae and free eggs in the alveolar spaces and interstitium, associated with a mixed inflammatory reaction composed mainly of macrophages, including binucleated and multinucleated cells, plasma cells, small lymphocytes and a few eosinophils. The larvae were approximately 20–30  $\mu\text{m}$  in width with a thin basophilic cuticle and numerous deep basophilic, internal nuclei. The eggs were round to ovoid, thin-walled and approximately 50–60  $\mu\text{m}$  in diameter; they contained either a larva or a morula. Additional findings included arterial thrombosis, proliferative endarteritis with recanalization, pulmonary necrosis, hemorrhage, edema, interstitial fibrosis, type II pneumocyte hyperplasia and small aggregates of hemosiderin-laden macrophages (Fig. 3a–d). No histological changes related to lungworms were noted in the pulmonary parenchyma collected from the other animals. Larvae recovered from the sediment of the centrifuged solutions were morphologically consistent

with *Angiostrongylus* spp. larvae, having a terminal oral opening and a kinked tail with a spine and a notch (Fig. 4).



**Fig. 4** First-stage larva of *Angiostrongylus vasorum* collected from the lungs. Note the median cranial oral opening and the kinked tail with a dorsal spine and a notch



**Fig. 3** Microscopical findings of pulmonary angiostrongylosis in an African golden wolf (*Canis lupaster*) with suspected *Angiostrongylus vasorum* infection. **a–d** The pulmonary interstitium is moderately expanded by numerous inflammatory nodules centered on parasitic eggs and larvae (arrows)

The BLAST analysis revealed 99.17–100% nucleotide identity to numerous other *A. vasorum* isolates from dogs and foxes registered in Europe (e.g. GenBank Accession Numbers: OQ210698, GQ982791, GQ982874). The sequence was deposited in GenBank under Accession Number PP872515.

All other animals were also negative for PCR as well.

## Discussion

In the present study, we demonstrated the presence of *A. vasorum* in Algeria, provided the first molecular confirmation of this species in Africa, and report a new host-parasite association. The finding of *A. vasorum* may indicate the continuous geographical expansion of this parasite to new areas that were previously not considered endemic, as recorded also in Morocco [30]. Interestingly, although the first report of *A. vasorum* in Africa dates back to > 50 years ago, when it was identified in domestic dogs from Uganda [29], the lack of more recent reports could be attributed to the lack of awareness or lack of interest among veterinarians and dog owners regarding this potentially fatal disease, as noted in a previous study [4], or, more likely, to the sporadic occurrence of *A. vasorum* in Africa. Another hypothesis would be that *A. vasorum* was introduced into northern Africa more recently. In the more recent report from Morocco, *A. vasorum* was identified in a symptomatic domestic dog (*C. familiaris*) from Rabat city using morphological identification of L1 collected from feces. The occurrence of *A. vasorum* is related to several factors, both biotic (humidity, temperature) and abiotic (intermediate, paratenic and final hosts). Numerous gastropod species have been reported in the study area from Algeria where the *A. vasorum*-positive carcass was collected [37]. However, the specificity of *A. vasorum* for the intermediate host is broad, and many snails and slugs can host the larvae for their development [27]. In endemic areas in Europe, the emergence of this parasite could be linked to various factors, such as climate change, urbanization of the red fox and dog movement/transport [38].

The present finding underlines the importance of complementary diagnostic methods for the detection and identification of lungworms, as has been previously recommended [35]. Once again, histology proved to be a key method for the detection of *A. vasorum* infection [39]. Explanations for not finding adult nematodes during the necropsy likely include a low number of worms or the lack of use of additional methods, such as the artificial digestion technique. Examination of feces using the larval concentration method may also have revealed some larvae although the carcass was frozen [40]. The absence of infection in the carcasses of the other animal species examined might be correlated to the low number of

examined animals, their food habits, their unsuitability as hosts, the recent introduction of the parasite to Algeria or the patchy distribution of the parasite.

There are many stray and free-roaming dogs in Algeria [41] that share the same lifestyle (scavenging) as other canids, such as the African golden wolf, and these could be easily infected with *A. vasorum*. Therefore, further studies are required for a better understanding of the epidemiological scenario of *A. vasorum* and to determine the potential life-cycle pattern of *A. vasorum* in Algeria.

## Conclusions

This new host and geographical record of *A. vasorum* expands current knowledge on this clinically important parasite, highlighting the importance of studies in wild carnivores from areas where the investigations on their parasitic fauna have been historically limited. Based on these results, we also advocate the use of complementary diagnostic techniques when examining dead animals. Additionally, due to the possible severe evolution of the infection in domestic dogs, awareness among local veterinarians should be raised for including infections by *A. vasorum* on differential diagnosis, especially in symptomatic animals.

## Acknowledgements

We are thankful to Alla Eddine Bouguerra, Khalil Ferhat and Hamza Djelab for the carcass collections and the local authorities for their collaboration in this study. The authors thank the association "Algeria BILLY pour la compassion, l'humanité et l'empathie" for its collaboration.

## Author contributions

NM collected the carcasses, performed necropsies and wrote the first draft. GD performed necropsies and the PCRs and revised the manuscript. AMI performed the PCRs and revised the manuscript. CGT, AGN, MT performed the histopathological examinations and revised the manuscript. ZB and ADM coordinated the study, revised and approved the final version of the manuscript. All authors read and approved the final manuscript.

## Funding

This study is a part of a PhD program for NM provided by the Romanian MAE (0713/III/4/CMJ/01.07.2020).

## Availability of data and materials

No datasets were generated or analyzed during the current study.

## Declarations

### Ethics approval and consent to participate

The National Environmental Research Center Annaba Ethics Committee granted ethical permission for this study. Algerian legislation (Ordinance No. 06–05 of 19 Jomada Ethania 1427, corresponding to July 15, 2006) was followed, regarding the collection and examination of the animals. The gathering of carcasses was reported to the local authorities, who also verbally approved the field activities.

### Consent for publication

Not applicable.

### Competing interests

Andrei D. Mihalca is an editor at *Parasites and Vectors*.



# Author details

<sup>1</sup>Department of Parasitology and Parasitic Diseases, University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, 3-5 Calea Mănătur, 400372 Cluj-Napoca, Romania. <sup>2</sup>Molecular Diagnosis Laboratory, Clinical Hospital of Infectious Diseases of Cluj-Napoca, 23 Iuliu Moldovan, 400348 Cluj-Napoca, Romania. <sup>3</sup>Department of Anatomic Pathology, University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, 3-5 Calea Mănătur, 400372 Cluj-Napoca, Romania. <sup>4</sup>National Environmental Research Center, Sidi Amar Campus, BP 2024, 23005 Annaba, Algeria.

Received: 13 June 2024 Accepted: 11 October 2024

Published online: 28 October 2024

# References

- Serres E. Entozoaires trouvés dans l'oreille droite, le ventricule correspondant et l'artère pulmonaire d'un chien. *J Vét Midi*. 1854;7:70.
- Uribe M, Segeritz L, Schnyder M, Taubert A, Hermosilla C, López-Osorio S, et al. Nationwide seroprevalence survey of *Angiostrongylus vasorum*-derived antigens and specific antibodies in dogs from Colombia. *Microorganisms*. 2022;10:1565. <https://doi.org/10.3390/microorganisms10081565>.
- Morgan ER, Modry D, Paredes-Esquivel C, Foronda P, Traversa D. Angiostrongylosis in animals and humans in Europe. *Pathogens*. 2021;10:1236. <https://doi.org/10.3390/pathogens10101236>.
- Bolt G, Monrad J, Koch J, Jensen AL. Canine angiostrongylosis: a review. *Vet Rec*. 1994;135:447–52.
- Chapman PS, Boag AK, Guitian J, Boswood A. *Angiostrongylus vasorum* infection in 23 dogs (1999–2002). *J Small Anim Pract*. 2004;45:435–40.
- Koch J, Willesen J. Canine pulmonary angiostrongylosis: an update. *Vet J*. 2009;179:348–59. <https://doi.org/10.1016/j.tvjl.2007.11.014>.
- King MCA, Grose RMR, Startup PG. *Angiostrongylus vasorum* in the anterior chamber of a dog's eye. *J Small Anim Pract*. 1994;35:326–8.
- Rosenlund P, Boserup F, Monrad J. *Angiostrongylus vasorum* I forreste øjenkammer hos hund [*Angiostrongylus vasorum* in the anterior chamber of the eye of a dog]. *Dansk-Vet tidsskr*. 1991;74:451–3 (Abstract in English).
- Cury MC, Lima WS. Rupture of femoral artery in a dog infected with *Angiostrongylus vasorum*. *Vet Parasitol*. 1996;65:313–5.
- Oliveira-Júnior SD, Barçante JMP, Barçante TA, Ribeiro VM, Lima WS. Ectopic location of adult worms and first-stage larvae of *Angiostrongylus vasorum* in an infected dog. *Vet Parasitol*. 2004;121:293–6. <https://doi.org/10.1016/j.vetpar.2004.02.018>.
- Taulescu MA, Negoescu A, Ungur A, Toma C, Ioniță AM, Gal C, et al. Is the *Angiostrongylus vasorum* infection in domestic dogs underestimated or misdiagnosed? A comprehensive presentation of four lethal cases. *Front Vet Sci*. 2023;10:1146713. <https://doi.org/10.3389/fvets.2023.1146713>.
- Rosen L, Ash LR, Wallace GD. Life history of the canine lungworm *Angiostrongylus vasorum* (Baillet). *Am J Vet Res*. 1970;31:131–42.
- Guilhonn J, Cens B. *Angiostrongylus vasorum* (Baillet, 1866) Etude biologique et morphologique. [*Angiostrongylus vasorum* (Baillet, 1866) Morphological and biological study]. *Ann Parasit Hum Comp*. 1973;48:567–96.
- Bolt G, Monrad J, Frandsen F, Henriksen P, Dietz HH. The common frog (*Rana temporaria*) as a potential paratenic and intermediate host for *Angiostrongylus vasorum*. *Parasitol Res*. 1993;79:428–30.
- Mozzer LR, Lima WS. *Gallus gallus domesticus*: paratenic host of *Angiostrongylus vasorum*. *Vet Parasitol*. 2015;207:81–4.
- Morgan RE, Shaw ES, Brenan F, De Wall DT. *Angiostrongylus vasorum*: a real heartbreaker. *Trends Parasitol*. 2005;21:49–51. <https://doi.org/10.1016/j.pt.2004.11.006>.
- Barçante TA, Barçante JMDP, Dias SRC, Kamensky, 1905: emergence of third-stage larvae from infected *Biomphalaria glabrata* snails. *Parasitol Res*. 2003;91:471–5 <https://doi.org/10.1007/s00436-003-1000-9>.
- Martin MWS, Neal C. Distribution of angiostrongylosis in Cornwall. *J Small Anim Pract*. 1992;33:327–30.
- Morgan ER, Jefferies R, Krajewski M, Ward P, Shaw SE. Canine pulmonary angiostrongylosis: the influence of climate on parasite distribution. *Parasitol Int*. 2009;59:406–10. <https://doi.org/10.1016/j.parint.2009.08.003>.
- Bourque A, Whitney H, Conboy G. *Angiostrongylus vasorum* infection in a coyote (*Canis latrans*) from Newfoundland and Labrador, Canada. *J Wildl Dis*. 2005;41:816–9. <https://doi.org/10.7589/0090-3558-41.4.816>.
- Lima WS, Guimaraes MP, Lemos IS. Occurrence of *Angiostrongylus vasorum* in the lungs of the Brazilian fox *Dusicyon vetulus*. *J Helminthol*. 1994;68:87.
- Takács A, Szabó L, Juhász L, Takács A, Lanszki J, Takács P, et al. Data on the parasitological status of golden jackal (*Canis aureus* L., 1758) in Hungary. *Acta Vet Hung*. 2014;2:33–41. <https://doi.org/10.1556/AVet.2013.058>.
- Gavrilović P, Marinković D, Vidanović D, Dobrosavljević I, Gavrilović A. Are golden jackals (*Canis aureus*) definitive hosts for *Angiostrongylus vasorum*? *Transbound Emerg Dis*. 2019. <https://doi.org/10.1111/tbed.13284>.
- Orioles, M, Fabbri D, Miani G, Pesaro S, Dorigo L, Bregoli M, et al. (2024). Double trouble: co-infection of *Angiostrongylus vasorum* and *Dirofilaria immitis* in golden jackal (*Canis aureus*) in Friuli Venezia Giulia, Italy. *Int J Parasitol Parasites Wildl*. 24:1009692024; <https://doi.org/10.1016/j.ijppaw.2024.100969>.
- PPatterson-Kane JC, Gibbons LM, Jefferies R, Morgan ER, Wenzlow N, Redrobe SP. Pneumonia from *Angiostrongylus vasorum* infection in a red panda (*Ailurus fulgens fulgens*). *J Vet Diagn Invest*. 2009; 21(2): 270–273. <https://doi.org/10.1177/104063870902100219>.
- Gillis-Germitsch N, Manser MB, Hilbe M, Schnyder M. Meerkats (*Suricata suricatta*), a new definitive host of the canid nematode *Angiostrongylus vasorum*. *Int J Parasitol Parasites Wildl*. 2017;6(3): 349–353. <https://doi.org/10.1016/j.ijppaw.2017.10.002>.
- Conboy GA. Canine angiostrongylosis: The French heartworm: an emerging threat in North America. *Vet Parasitol*. 2011;176(4):382–389. <https://doi.org/10.1016/j.vetpar.2011.01.025>.
- Guelndner EK, Gilli U, Strube C, Schnyder M. Seroprevalence, biogeographic distribution and risk factors for *Aelurostrongylus abstrusus* infections in Swiss cats. *Vet Parasitol*. 2019;266:27–33. <https://doi.org/10.1016/j.vetpar.2018.12.013>.
- Bwangamoi O. *Angiostrongylus vasorum* and other worms in dogs in Uganda. *Vet Rec*. 1972;91:267.
- Idrissi H, Khatat SEH, Duchateau L, Kachani M, Azrib R, Sahibi H. Canine cardiopulmonary nematodes in Morocco: Prevalence of *Dirofilaria immitis* and report of the first autochthonous infection with *Angiostrongylus vasorum*. *Mor J Agric Sci*. 2022;3(2):117–123. <https://doi.org/10.5281/ZENODO.8030663>.
- Ahmim M. Les mammifères sauvages d'Algérie Répartition et Biologie de la conservation. Les Editions du Net, St St Ouen; 2019.
- Gutema TM, Atickem A, Tsegaye D, Bekele A, Sillero-Zubiri C, Marino J, et al. Foraging ecology of African wolves (*Canis lupaster*) and its implications for the conservation of Ethiopian wolves (*Canis simensis*). *R Soc Open Sci*. 2019;6:190772. <https://doi.org/10.1098/rsos.190772>.
- Klevezal' GA, Kleinenberg SE. Age determination of mammals by layered structure in teeth and bone. Fisheries Research Board of Canada; 1967.
- Roulichova J, Andera M. Simple method of age determination in red fox, *Vulpes vulpes*. *Folia Zool*. 2007;56:440.
- Rojas A, Germitsch N, Oren S, Szazmand A, Deak G. Wildlife parasitology: sample collection and processing, diagnostic constraints, and methodological challenges in terrestrial carnivores. *Parasites Vectors*. 2024;17(1): 127. <https://doi.org/10.1186/s13071-024-06226-4>.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol*. 1994;3:294–9.
- Belhouani H, El-Okki MEH, Afri-Mehennaoui FZ, Sahli L. Terrestrial gastropod diversity, distribution and abundance in areas with and without anthropogenic disturbances, Northeast Algeria. *Biodiversitas J Biol Divers*. 2019;20:243–249. <https://doi.org/10.13057/biodiv/d200128>.
- Montlaur L. Prévalence de l'angiostrongylose canine dans le sud-ouest de la France. Thesis. Ecole nationale vétérinaire de Toulouse, Toulouse; 2015.
- De Liberato C, Grifoni G, Lorenzetti R, Meoli R, Cucumelli C, Mastromattei A, et al. *Angiostrongylus vasorum* in wolves in Italy: prevalence and pathological findings. *Parasit Vectors*. 2017;10:386. <https://doi.org/10.1186/s13071-017-2307-1>.
- Jeffery RA, Lankester MW, McGrath MJ, Whitney HG. *Angiostrongylus vasorum* and *Crenosoma vulpis* in red foxes (*Vulpes vulpes*) in Newfoundland. *Canada Can J Zool*. 2004;82:66–74. <https://doi.org/10.1139/z03-211>.
- Saadi F. Bilan épidémiologique de la rage en Algérie. Thesis. Université de Montpellier 2, Montpellier; 2008.

# Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

[onlineservice@springernature.com](mailto:onlineservice@springernature.com)