



# Identification of small circular DNA viruses in coyote fecal samples from Arizona (USA)

Savage C. Hess<sup>1</sup> · Katherine C. B. Weiss<sup>1</sup> · Joy M. Custer<sup>2</sup> · Jesse S. Lewis<sup>3</sup> · Simona Kraberger<sup>2</sup> · Arvind Varsani<sup>1,2,4,5</sup> 

Received: 26 October 2023 / Accepted: 22 November 2023 / Published online: 28 December 2023  
© The Author(s), under exclusive licence to Springer-Verlag GmbH Austria, part of Springer Nature 2023

## Abstract

Coyotes (*Canis latrans*) have a broad geographic distribution across North and Central America. Despite their widespread presence in urban environments in the USA, there is limited information regarding viruses associated with coyotes in the USA and in particular the state of Arizona. To explore viruses associated with coyotes, particularly small DNA viruses, 44 scat samples were collected (April–June 2021 and November 2021–January 2022) along the Salt River near Phoenix, Arizona (USA), along 43 transects (500 m). From these samples, we identified 11 viral genomes: two novel circoviruses, six unclassified cressdnaviruses, and two anelloviruses. One of the circoviruses is most closely related to a circovirus sequence identified from an aerosolized dust sample in Arizona, USA. The second circovirus is most closely related to a rodent-associated circovirus and canine circovirus. Of the unclassified cressdnaviruses, three encode replication-associated proteins that are similar to those found in protists (*Histomonas meleagridis* and *Monocercomonoides exilis*), implying an evolutionary relationship with or a connection to similar unidentified protist hosts. The two anelloviruses are most closely related to those found in rodents, and this suggests a diet-related identification.

## Introduction

Coyotes (*Canis latrans*) are mid-sized mesocarnivores in the family Canidae that have an expansive range across the gradient of urbanization, ranging from rural to urban areas throughout North America and much of Central America [1, 2]. They are born into small packs consisting of their parents and siblings [3]. Notably, these animals are physiologically

similar to domestic dogs (*Canis lupus familiaris*) [4]. In general, coyotes tend to eat small animals but have been known to kill larger animals such as white-tailed deer (*Odocoileus virginianus*) [5]. In addition, they also consume plant matter, such as fruits and seeds [5, 6]. Their presence in human-occupied environments has led to a long history of human-coyote conflict due to coyote predation of domesticated animals, such as sheep and pets [5–7]. Due to their generalist habitat requirements, social interactions, and omnivorous diet, coyotes are exposed to a suite of pathogens, which also can be transmitted to other wildlife and humans. However, the suite of pathogens that coyotes carry is unknown in many ecosystems, especially those influenced by urbanization.

Over the last six decades, there have been a significant number of viruses identified infecting coyotes. These include viruses in the families *Adenoviridae*, *Orthoherpesviridae*, and *Papillomaviridae* with double-stranded DNA genomes, *Coronaviridae*, *Flaviviridae*, *Picornaviridae*, and *Togaviridae* with single-stranded, positive-sense RNA genomes, and *Orthomyxoviridae*, *Paramyxoviridae*, *Peribunyaviridae*, *Phenuiviridae*, and *Rhabdoviridae* with single-stranded, negative-sense RNA genomes. Due to concerns of viral spillover between coyotes and domestic animals, there has been a significant amount of surveillance work associated with coyotes infected with or exposed to rabies virus [8–13],

Handling Editor: Sheela Ramamoorthy.

✉ Arvind Varsani  
arvind.varsani@asu.edu

- <sup>1</sup> The School of Life Sciences, Arizona State University, 427 E Tyler Mall, Tempe, AZ 85281, USA
- <sup>2</sup> The Biodesign Center for Fundamental and Applied Microbiomics, Arizona State University, 1001 S. McAllister Ave, Tempe, AZ 85287, USA
- <sup>3</sup> College of Integrative Sciences and Arts, Arizona State University, Polytechnic Campus, 6073 South Backus Mall, Mesa, AZ 85212, USA
- <sup>4</sup> Center of Evolution and Medicine, Arizona State University, 427 E Tyler Mall, Tempe, AZ 85281, USA
- <sup>5</sup> Structural Biology Research Unit, Department of Integrative Biomedical Sciences, University of Cape Town, Cape Town 7925, South Africa

canine morbillivirus [14–16], and canine parvovirus [14, 17–20]. This is not surprising, due to the close encounters of coyotes with humans and other canids (both wild and domestic) [21]. Nonetheless, the majority of the work on identification of virus infections in coyotes has been done through serological assays. To date, 151 viral sequences in GenBank associated with coyotes belong to members of the following species: *Varicellovirus canidalpha1* (family *Orthoherpesviridae*;  $n=2$ ), *Morbillivirus canis* (family *Paramyxoviridae*;  $n=20$ ), *Chaphamaparvovirus carnivoran1* (family *Parvoviridae*;  $n=4$ ), *Protoparvovirus carnivoran1* (family *Parvoviridae*;  $n=34$ ), and *Lyssavirus rabies* (family *Rhabdoviridae*;  $n=93$ ). Of these, there are only seven complete genome sequences: two from *Chaphamaparvovirus carnivoran1*, three from *Protoparvovirus carnivoran1*, and two from *Lyssavirus rabies* (Table 1).

Given the limited knowledge of circular small DNA viruses associated with coyotes in urbanized areas, we undertook a pilot project to see if we can detect these in a non-invasive manner via scat sampling. We sampled scat along the Salt River near Phoenix, Arizona, United States (USA), and amplified circular DNA viruses from these samples using a viral metagenomics protocol coupled with PCR amplification and cloning of viral genomes, resulting in our discovery of 11 novel viruses. Here, we describe our identification of two new anelloviruses, three new circoviruses, and six new viruses in the phylum *Cressdnaviricota* that cannot be assigned to currently established families.

## Materials and methods

### Sample collection

Coyote scat samples were collected along the Salt River within the Phoenix Metropolitan Area and Tonto National Forest, Arizona, USA, between April and June 2021 and between November 2021 and January 2022. Scats were collected monthly during the study duration along 43 (500 m) transects, which were recorded using the Gaia GPS phone application (<https://gaiagps.com/>). All scats were cleared from each transect one month prior to the first collection date at each site. Sites were then visited monthly for two to three months to collect coyote scats. Scats were identified morphologically [22] and collected in plastic bags with associated spatial locations and confidence in species identification (50%–100%). Scats were then stored in a  $-80^{\circ}\text{C}$  freezer. Scats with less than 100% confidence in species ID were later swabbed for remnant epithelial DNA and sent for microsatellite marker genetic testing to confirm they were deposited by a coyote (<https://www.wildlifegenetics.ca/>). Any scats not deposited by a coyote were excluded from analysis, as well as any scats unable to be swabbed that had

less than 100% confidence in species ID. A random subset of coyote scats with 100% confidence in morphological ID were also swabbed for DNA to confirm the success of our methodology. A subset of the collected scats ( $n=44$ ) was then used for viral analysis. Scats were selected based on a representative sample of location and season (fall–winter or spring–summer) collected, and  $\sim 2$  g of the fecal material was used downstream for viral work (Fig. 1).


For diet assessments, we conducted hard-parts analysis of prey remains in the coyote scats. We first sanitized the scats by baking them in an oven for 24 h at  $60^{\circ}\text{C}$  [22]. We then soaked scats in soapy water for 24–48 h and rinsed out any remaining fecal material using cheesecloth to isolate the dietary remains left in the scats for identification. Scat contents were then left to air-dry for a subsequent 24–48 h before storing in envelopes. We classified dietary items as mammal (to the lowest taxonomic classification possible), bird, snake, fish, arthropod (including insects, scorpions, and crayfish), vegetation (to family or genus, when possible), trash, egg-shell, and unidentifiable (including within each mammal category and overall). To identify mammals, five random samples of hair per scat were examined under a microscope (AmScope B120, United Scope, LLC, Irvine, CA) [23] and photographed to compare with (1) reference slides of hair using specimens from Arizona State University's Natural History Collections, (2) identification keys [24–26], and expert solicitation. We also identified teeth remains found in scats using reference images of teeth from specimens at Arizona State University's Natural History Collections, as well as expert solicitation. In addition to hair, domestic cat (*Felis catus*) remains were also identified via the presence of claws. Domestic cats were further distinguished from bobcats (*Lynx rufus*) via hair color and texture, as well as location, when compared with where bobcats were detected on camera traps. Similarly, domestic dogs (*Canis familiaris*) were distinguished from coyotes based on hair color and texture, as well as expert review. Lagomorph hairs were classified as desert cottontail rabbit (*Sylvilagus audubonii*) or unidentifiable lagomorph based on the number of rows of medulla within a sample being either greater than four (i.e., cottontail rabbit) or less than or equal to four (i.e., to distinguish between cottontail rabbit and black-tailed jack-rabbit, *Lepus californicus* [24]). Any sample with at least one lagomorph hair greater than four medulla cells wide was identified as being cottontail rabbit—and often corroborated by lagomorph teeth of a smaller size more likely representative of a cottontail rabbit—while lagomorph samples with only thinner hairs in terms of medullar width were identified as unidentifiable lagomorph. Neotominae species (i.e., *Neotoma* or *Peromyscus* spp.) were grouped together due to the similarity in their hair structure. Bird remains were identified via feathers or hollow bones present in the scats, while reptiles and fish were identified via scales and

**Table 1** Summary of sequences of viruses associated with coyotes available in the GenBank database prior to this study

Genome type	Family	Genus	Species	Virus	Accession no.	Country	Genome sequences		References
							Partial	Complete	
dsDNA	Orthoherpesviridae	Varicellovirus	<i>Varicellovirus canidalphal</i>	Canid alphaherpesvirus 1	KX062143, KX062144	Canada	2	-	[93]
ssRNA(-)	Paramyxoviridae	Morbillivirus	<i>Morbillivirus canis</i>	Canine distemper virus	OK666919, OK666920, OK666921, OK666922, OK666923, OK666924, OK666925, OK666926, OK666927, OK666928, OK666929, OK666930, OK666931, OK666932, OK666933, OK666934, OK666935, OK666936, OK666937, KT387222	Canada, Mexico	20	-	[94]
ssDNA(+/-)	Parvoviridae	Chaphamaparvovirus	<i>Chaphamaparvovirus carnivoran1</i>	Cachavirus	OM640108, OM640109, OM640110, OM640111	Canada (Newfoundland)	2	2	[18]
ssDNA(+/-)	Parvoviridae	Protoparvovirus	<i>Protoparvovirus carnivoran1</i>	Canine parvovirus	MH480648, MG425949, MG425950, KJ813866, KJ813867, KJ813868, KJ813872, KJ813874, KJ813875, KJ813876, KJ813883, KJ813884, KJ813885, KJ813886, KJ813887, KJ813888, KJ813889, KJ813891, KJ813892, JX475238, JX475239, JX475269, JX475273, JX475274, JX475275, JX475276, JX475277, JX475278	Canada (Newfoundland), Mexico, USA (Alabama, Alaska, Arkansas, Georgia, Montana, Nebraska, North Dakota, Oklahoma)	31	3	[17–19, 95]

Table 1 (continued)

Genome type	Family	Genus	Species	Virus	Accession no.	Country	Genome sequences		References
							Partial	Complete	
ssRNA(-)	<i>Rhabdoviridae</i>	<i>Lyssavirus</i>	<i>Lyssavirus rabies</i>	Rabies virus	OM203023, OM203045, MW566991, KU963491, KU963495, KJ174683, KC791818, KC791823, KC791961, KC792102, JQ685917, JQ685973, JQ685994, FJ228520, FJ228526, FJ228529, FJ228530, FJ228536, FJ228551, FJ228552, FJ228557, FJ228563, FJ228564, FJ228567, FJ228568, FJ228569, FJ228571, FJ228575, FJ228576, FJ228577, FJ228580, FJ228584, FJ228585, FJ228587, FJ228588, FJ228589, FJ228591, FJ228592, FJ228594, FJ228604, FJ228605, FJ228606, FJ228607, FJ228608, FJ228612, FJ228613, FJ228614, FJ228616, FJ228617, FJ228618, FJ228619, FJ228620, FJ228622, FJ228623, FJ228624, FJ228625, FJ228626, FJ228627, FJ228629, FJ228632, FJ228633, FJ228634, FJ228636, FJ228637, FJ228639, FJ228640, FJ228641, FJ228644, FJ228645, FJ228648, FJ228650, FJ228651, FJ228652, FJ228653, FJ228654, FJ228655, FJ228656, FJ228658, FJ228660, FJ228662, FJ228663, FJ228664, FJ228665, FJ228670, FJ228674, FJ228676, FJ228683, FJ228685, DQ118248, AY561764, AF369289, AF369308, AF369337	Canada (Ontario), Mexico (Coahuila, Texas), US-Mexico border, US (Arizona, California, New Mexico, Texas)	91	2	[13, 96–103]



**Fig. 1** Map of study sites along the Salt River Corridor in the Phoenix Metropolitan Area, Arizona, USA. Black circles indicate study sites where scats were collected for analysis but were not used due to an inability to detect, amplify, and clone the viruses from these sam-

ples. Sites in yellow indicate areas where coyote scats were analyzed for viruses, including (from left to right) samples V16\_S08, N10\_S01, N07\_S03, N06\_S02, and T05\_2F02.

confirmed via expert review. Due to desiccation, crayfish remains were only identifiable when washing the scats prior to drying. Vegetation, particularly seeds, was identified via expert solicitation, and eggshells and trash were identified by sight. Diet remains that contributed to <1% of the volume of a dried scat were not considered in analyses, as these items were expected to be incidental.

### Sample processing and viral metagenomics

The 44 scat samples were resuspended separately and homogenized in 3 ml of SM buffer. The homogenate was centrifuged at 10,000×g and passed through a 0.22-μm syringe filter. Two hundred microliters of the filtrate was used for DNA extraction using a High Pure Viral Nucleic Acid Kit (Roche Diagnostics, USA). A TempliPhi 2000 kit (GE Healthcare, USA) was used to amplify circular DNA in the viral extracts, using rolling-circle amplification (RCA). The RCA products were combined into three pools, and these were used to generate Illumina sequencing libraries using Illumina DNA Prep, (M) Tagmentation (96 Samples, IPB) Kit (Illumina, USA) and sequenced on an Illumina

HiSeq2500 instrument at Psomagen Inc. (Maryland, USA). The raw reads (2 × 150 nt) were trimmed using Trimmomatic v0.39 [27] and assembled *de novo* using MEGAHIT v1.2.9 [28]. Contigs of >1000 nt were compared, using BLASTx [29], to a viral RefSeq protein sequence database (release 207) for viral-like sequences.

### Recovery of circular viral genomes

Based on the *de novo*-assembled contigs with similarities to small circular DNA viruses, abutting primer pairs were designed and used to amplify complete viral genomes (Supplementary Table S1). The primers were used in a polymerase chain reaction (PCR) with KAPA HiFi HotStart DNA Polymerase (Roche Diagnostics, USA), using the following thermal cycling conditions to screen and amplify the genome with the specific primers and the RCA product from the 44 scat samples as templates: initial denaturation at 98°C for 3 min, followed by 25 cycles of 98°C for 15 s, 60°C for 15 s, and 72°C for 3 min, and a final extension at 72°C for 3 min. The PCR amplicons were resolved on a 0.7% agarose gel, and bands corresponding to the expected nucleotide length



were excised and gel-purified using a MEGAquick-spin Plus Fragment DNA Purification Kit (iNtRON Biotechnology, South Korea). The purified product was ligated into the pJet 1.2 plasmid vector (Thermo Fisher Scientific, USA), and the construct was used to transform competent *E. coli* DH5 cells. Bacterial colonies were grown overnight, and recombinant plasmids were extracted using a Fast DNA-spin Plasmid DNA Purification Kit (iNtRON Biotechnology, South Korea) and then sequenced at Macrogen Inc. (South Korea) by primer walking. The Sanger sequences were assembled using Geneious Prime (Biomatters Ltd., New Zealand). The genomes were annotated using ORFfinder (ncbi.nlm.nih.gov/orffinder/) and manually checked.

### Viral genome analysis

For the cressdnaviruses, a dataset of replication-associated protein (Rep) sequences was assembled from ~ 4000 sequences, which were extracted from genome sequences of representative members of established virus families as well as currently unclassified viruses with sequences available in the GenBank database, and those from this study. This Rep protein sequence dataset was used to cluster new sequences at the family level based on sequence similarity network (SSN) analysis, using EST-EFI [30] with a threshold of 60. The SSN was visualized using Cytoscape v3.8.2 [31] with an organic layout. The clusters containing the Rep sequences from this study were extracted for downstream analysis. Representative classified cressdnavirus and CRESSV1-6 group Rep sequences were aligned with the Rep sequences from this study, together with the other sequences from their clusters (clusters 1–4), using MAFFT [32]. The alignment was trimmed using TrimAL [33], and this was then used to construct a maximum-likelihood phylogenetic tree using IQ-Tree 2 [34] with the Q.pfam+F+G4 substitution model, and the tree was visualized in iTOL v6 [35]. For each cluster,

outgroup Rep sequences were included and aligned using MAFFT [32].

For the anelloviruses, ORF1 is the most highly conserved coding region. Therefore, a representative dataset of ORF1 protein sequences from GenBank was compiled, and the sequences were aligned with those from this study, using MAFFT [32]. The coyote anelloviruses clustered with members of the genus *Wawtorquevirus*. Therefore, a maximum-likelihood phylogenetic tree based on ORF1 sequences of viruses of the genus *Wawtorquevirus* was constructed, using ORF1 sequences of thetatorqueviruses as an outgroup.

The best amino acid substitution model for each phylogeny was determined using ProtTest 3 [36]. The phylogenetic trees were inferred using PhyML 3 [37] with aLRT branch support. Branches with <0.8 posterior support were collapsed using TreeGraph2 [38]. All pairwise identities (genome-wide and amino acid) were determined using SDT v1.2 [39].

### Results and discussion

Among the *de novo* assembled contigs, we identified 192 viral contigs (>1000 nt). Of these, six were identified to be related to anelloviruses, 39 to cressdnaviruses, and 147 to bacteriophages (microviruses and caudoviruses). We selected the largest contigs (>1500 nt) that had similarities to anelloviruses ( $n = 2$ ) and cressdnaviruses ( $n = 8$ ) and designed abutting primer pairs (Table 2) to screen and amplify the genomes from the 44 scat samples.

We were able to detect, amplify, and clone the viral genomes from five of the 44 samples, i.e., sample ID T05\_2F02 ( $n = 6$ ), N10\_S01 ( $n = 1$ ), N07\_S03 ( $n = 1$ ), N06\_S02 ( $n = 1$ ), V16\_S08 ( $n = 2$ ) (see Table 3 for summary and Fig. 1). Of the 11 viral genomes that were detected, nine were from members of the phylum *Cressdnaviricota* (three from the family *Circoviridae* and six from unclassified

**Table 2** List of viruses identified with their GenBank accession numbers and the coyote scat sample information

Scat sample ID	Latitude	Longitude	Sampling date	Family/group	Virus name	GenBank accession no.
T05_2F02	33.54753	− 111.602	17 Dec 2021	<i>Anelloviridae</i>	Banfec anellovirus 1	OQ599920
				<i>Anelloviridae</i>	Banfec anellovirus 2	OQ599921
				Cluster 2	Banfec virus 2	OQ599926
				Cluster 3	Banfec virus 1	OQ599925
				Cluster 4	Banfec virus 4	OQ599928
				Cluster 4	Banfec virus 5	OQ599929
N10_S01	33.40255	− 112.178	24 Apr 2021	<i>Circoviridae</i>	Banfec circovirus 1	OQ599922
N07_S03	33.41004	− 112.142	23 Apr 2021	<i>Circoviridae</i>	Banfec circovirus 1	OQ599923
N06_S02	33.41114	− 112.129	23 Apr 2021	<i>Circoviridae</i>	Banfec circovirus 2	OQ599924
V16_S08	33.38929	− 112.245	26 Apr 2021	CRESSV1	Banfec virus 3	OQ599927
				Cluster 1	Banfec virus 6	OQ599930

**Table 3** Summary of the rolling-circle replication (RCR) endonuclease motifs (motifs I, II, and III) and the superfamily 3 (SF3) helicase motifs (Walker A, Walker B, and Motif C) of the cressdnaviruses

Family/group	Virus	Accession no.	Motif I	Motif II	Motif III	Walker A	Walker B	Motif C
<i>Circoviridae</i>	Banfec circovirus 1	OQ599922	CFTVNN	PHLQG	QNQAYCQK	GPPGVGKTRYAV	ILDDF	ITSN
	Banfec circovirus 1	OQ599923	CFTVNN	PHLQG	QNQAYCQK	GPPGVGKTRYAV	ILDDF	ITSN
	Banfec circovirus 2	OQ599924	CFTINN	PHLQG	SNRDYCSK	GPPGCGKSRLCF	ILDDF	ITSN
Unclassified CRESSV1	Banfec virus 3	OQ599927	CVTLFK	IHWHL	EAIAYAKK	GAAGSGKSHHCW	WFDEF	ISTV
Unclassified cluster 1	Banfec virus 6	OQ599930	CFTLNN	PHLQG	QASNYCKK	GKSGVGKTRFAV	IIDDF	ITCE
Unclassified cluster 2	Banfec virus 2	OQ599926	VITWNN	HHIQG	DCILYCRK	GRPGTGKSRWAR	IIDDF	VTSN
Unclassified cluster 3	Banfec virus 1	OQ599925	QFTLNQ	KHMH	QNIAYIEK	GDTGCGKTYNAY	VIEEF	ICSI
Unclassified cluster 4	Banfec virus 4	OQ599928	AFTLNN	PHLQG	QNKKYCSK	GPARKGKTNKAM	VFDDF	FTSI
	Banfec virus 5	OQ599929	AFTVNN	AHLQG	QNKKYCTK	GPARKGKTRRAM	IFDDF	FTSI

cressdnaviruses), and two were from members of the family *Anelloviridae*. It is noteworthy that we amplified two circovirus isolates sharing 97.4% identity using the same primer pair from two samples (N10\_S01 and N07\_S03) collected a day apart.

We were able to use the primer pairs on only a subset of the samples because some of the scat samples had been degraded due to prolonged exposure to the dry and hot climate in Phoenix prior to sampling. Notably, six viruses were identified from sample ID T05\_2F02, which was noted for being particularly wet at the time of collection compared to the other scat samples used for analysis, suggesting that recoverability of viral signatures may be a function of the time since deposition, or at least the moisture of the sample. A diet analysis showed that sample T05\_2F02 contained evidence of Neotominae species (likely *Neotoma* or *Peromyscus*), N10\_S01 had desert cottontail rabbit (*Sylvilagus audubonii*), Neotominae, and avian species, N07\_S03 had desert cottontail rabbit and *Chaetodipus* species, and N06\_S02 and V16\_S08 primarily had desert cottontail rabbit.

The viral sequences identified here have been deposited in the GenBank database under the accession numbers OQ599920–OQ599930, and their names are derived from the term “ban” for the O’odham word for coyote and “fec” as an abbreviation for “feces” (Table 2).

### Cressdnaviricota

Cressdnaviruses (phylum *Cressdnaviricota*) are circular Rep-encoding single-stranded DNA viruses that infect a broad range of eukaryotic species [40]. The Rep protein has a HUH endonuclease domain and a superfamily 3 helicase domain, which are essential for initiating rolling-circle replication [41–44]. Within the phylum *Cressdnaviricota*, there are 12 established families (*Amesuviridae*, *Bacilladnaviridae*, *Circoviridae*, *Geminiviridae*, *Genomoviridae*, *Metaxoviridae*, *Nanoviridae*, *Naryaviridae*, *Nenyaviridae*, *Redondoviridae*, *Smacoviridae*, and *Vilyaviridae*)

with six additional informal groupings of viruses labeled CRESSV1–6 [40, 45, 46] (Fig. 2). Many cressdnaviruses have been identified but not yet classified [47, 48].


Because the Rep protein is the most conserved protein among cressdnaviruses, SSN analysis has been used in the past to roughly group Rep sequences of new viruses at the family level with a network threshold of 60 [47, 49–51]. Based on SSN analysis using EFI [30], the nine Rep sequences of the cressdnaviruses identified in this study fall into clusters with those of members of the family *Circoviridae* ( $n=3$ ) and the clusters CRESSV1 ( $n=1$ ), cluster 1 ( $n=1$ ), cluster 2 ( $n=1$ ), cluster 3 ( $n=1$ ), and cluster 4 ( $n=2$ ) (Fig. 2). The CRESSV1 group has been described previously and mentioned in literature but is currently not classified [40, 45, 46]. In the Rep sequences of these nine cressdnaviruses, we identified the three HUH endonuclease domains and the three SF3 helicase domains (Table 3).

### Circoviridae

Circoviruses (family *Circoviridae*) are classified into two genera: *Circovirus* and *Cyclovirus* [52, 53]. Members of the genus *Circovirus* have been found to be associated with various diseases in avian and mammal species, e.g., psittacine beak and feather disease caused by beak and feather disease virus [54] and postweaning multisystemic wasting syndrome caused by porcine circovirus 2 [55]. Approximately a decade ago, canine circovirus 1 was identified and associated with hemorrhagic enteritis in canines, and it appears to have become widespread globally [56–61]. In general, circoviruses have small circular ssDNA genomes of ~1.7–2.2 kb encoding a capsid protein (CP) and a Rep protein [52]. The species demarcation threshold for circoviruses is 80% pairwise identity [43].

From the scat (sample IDs N10\_S01, N07\_S03, and N06\_S02), we identified three circovirus genomes (two banfec circovirus 1 genomes [OQ599922 and OQ599923] and one banfec circovirus 2 genome [OQ599924]). The two

**Fig. 2** Maximum-likelihood phylogenetic tree (inferred using the Q.pfam+F+G4 substitution model) based on Rep sequences of representative classified cressdnaviruses, CRESSV1-6, the four clusters of unclassified Rep, and the sequences from this study. The Rep sequences of the cressdnaviruses from this study that fall within the family level clusters are indicated in bold blue font.



banfec circovirus 1 genomes (1680 nt) share 97.4% pairwise nucleotide sequence identity. When compared to other circoviruses, banfec circovirus 1 shares 97.1–97.8% pairwise nucleotide sequence identity with the virus D1\_1542 (MW678871), which was identified in aerosolized dust particles in Arizona, USA [62], and 71.3–71.9% with calfel virus LSF45\_cir359 (ON596197), which was identified in a fecal sample from a bobcat (*Lynx rufus*) in California, USA [63].

On the other hand, the genome of banfec circovirus 2 (2079 nt) shares 70.8% pairwise nucleotide sequence identity with dipodfec virus UA04Rod\_4537 (OM869597)


identified in a Merriam's kangaroo rat (*Dipodomys merriami*) fecal sample from Arizona, USA [51], and 69.1% with a canine circovirus (KC241982) identified in the liver of a domestic dog (*Canis lupus familiaris*) [58].

Banfec circoviruses 1 and 2 represent two new species based on the 80% species demarcation criteria outlined for circoviruses [52]. It is also noteworthy that the circoviruses identified in this study are most closely related to other circoviruses from the Southwest of the USA. However, this may be due to a gap in our knowledge of circovirus diversity.




Phylogenetic analysis of the Rep sequences of the circoviruses (Fig. 3) revealed that banfec circovirus 1 forms a clade with viruses from samples of bobcat (*Lynx rufus*) feces collected in California, USA [63] (ON596197), a South China field mouse (*Apodemus draco*) from China (KY370037) [64], a tawny owl (*Strix aluco*) from Italy (OL411978) [65], and a rufous horseshoe bat (*Rhinolophus sinicus*) and an intermediate horseshoe bat (*Rhinolophus affinis*) from China

(KJ641723-KJ641724) [66] (Fig. 3). The Rep sequence of banfec circovirus 2 clusters with those of circoviruses identified in samples from a dog (KC241982) [58] in the USA, a bobcat (MT610105) in Mexico [67], and various rodents, i.e., a Merriam's kangaroo rat (OM869597) from Arizona, USA [51], a hoary bamboo rat (*Rhizomys pruinosus*, MF497827) from China [68], and a Siberian jerboa (*Allactaga sibirica*, KY370029), a Chevrier's field mouse



**Fig. 3** Maximum-likelihood phylogenetic tree (inferred using the Q.pfam+F+G4 substitution model) based on Rep sequences based on (one per species) of circoviruses and rooted cyclovirus Rep sequences. The sequences from this study are indicated in bold blue

font, and the genome organizations of the circoviruses identified in this study are shown to the right of the phylogenetic tree. Common names of the sources/hosts of the viruses are shown in parentheses next to the virus names.



**Fig. 4** Maximum-likelihood trees (inferred using the rtREV+G+I substitution model) based on Rep sequences of members of unclassified cressdnavirus cluster CRESSV1 (A) and cluster 1 (B) rooted with Reps of vilyaviruses. The sequences from this study are indicated in bold blue font, and the genome organizations of the circoviruses identified in this study are shown to the right of the phylogenetic tree. Common names of the sources/hosts of the viruses are shown in parentheses next to the virus names.

(*Apodemus chevrieri*, KY370042) and a Clarke's vole (*Neodon clarkei*, KY370034, KY370039), all from China [64] (Fig. 1).

The similarity of banfec circovirus 1 to the virus D1\_1542 (MW678871) from Arizona suggests that this virus is circulating in animals in Arizona and was inadvertently sampled, likely due to aerosolization of animal fecal matter, by Finn et al. [62]. Furthermore, the similarity to calfel virus LSF45\_cir359 (ON596197) from a bobcat [63] provides some support that this may be a virus that is associated with canids, at least in the Southwest of the USA. Given that banfec circovirus 2 is distantly related to known circoviruses, although it clusters with primarily rodent-derived circoviruses, it is likely that this is a rodent circovirus that was present in the prey of the coyote.

## Unclassified cressdnaviruses

### CRESSV1

The Rep sequence of banfec virus 3 from scat V16\_S08 (Table 3) clustered with those of viruses in CRESSV1. Banfec virus 3 (3047 nt; OQ599927) encodes a Rep and a CP (Fig. 4). BLASTn analysis showed that its CP shares 68.51% identity (22% genome coverage; E-value, 2E-61) with the genome of rodent circovirus RtBi-CV-1/FJ2015 (KY370035), derived from a greater bandicoot rat (*Bandidota indica*) in China [64], and its Rep shares 56.9% amino acid sequence identity. The Rep phylogeny showed that it clusters with viruses from human feces from Chile (KT600067) [69], the Netherlands (LC708000) [48], Nicaragua (KT600066) [69], Peru (KT600065) [69], and Venezuela (MG571895, MG571897) [70] and a virus from rhesus monkey feces (*Macaca mulatta*, KU043425) [71] (Fig. 4). Banfec virus 3 would represent a new species in a genus with rodent circovirus RtBi-CV-1/FJ2015 when the viruses in CRESSV1 get assigned to a new family.

### Cluster 1

Banfec virus 6 (1826 nt, OQ599930), which encodes a Rep and CP and was identified in scat V16\_S08 (Table 3), is part of cluster 1 based on Rep amino acid sequences. A BLASTn analysis revealed that it shares 70.96% sequence identity (16% coverage; E-value, 3E-30) with CRESS virus isolate


ctdb70 (MK012454) from turkey (*Meleagris gallopavo*) tissue [47]. The Rep of banfec virus 6 shares <58% pairwise amino acid sequence identity with all other Rep proteins within this cluster. Phylogenetically (Fig. 4), the Rep of banfec virus 6 clusters with those of viruses derived mainly from wastewater in the USA (KY487946, KY487959) [72], water samples from France (MT478490, MT478498, MT478501, MT478503, MT478516, MT478524, MT478544, MT478551, MT478572, MT478575) [73], freshwater invertebrates from Lake Sarah in New Zealand [74], a crucian tissue (*Carassius* sp.) (MK012490) [47], red panda feces in China (*Ailurus fulgens*, MZ556125, MZ556127) [75], and red-crowned crane (*Grus japonensis*, KY3122556) [76].

### Cluster 2

Banfec virus 2 (4332 nt, OQ599926) derived from scat T05\_2F02 (Table 3) contains three large ORFs in its genome in addition to the Rep and CP genes, and it belongs to cluster 2. The genome of banfec virus 2 shares 68.54% sequence identity (4% coverage; E-value, 2E-06) with human fecal virus Jorvi2 (MF118167) [77]. The Rep of banfec virus 2 shares <44% amino acid sequence identity with those of the other members of this cluster. This cluster consists of Rep-like sequences of circular DNA molecules derived from desert tortoise (*Gopherus* sp.) feces from the USA (MK858259, MK858260) [78], viruses from human feces from Finland (MF118167) [77] and the Netherlands (LC708032, LC708033) [48], striped dwarf hamster (*Cricetulus barabensis*) from China (KY370030, KY370031) [64], and Chevrier's field mouse (*Apodemus chevrieri*), also from China (KY370041) [64] (Fig. 5). A BLASTp analysis of the Rep of banfec virus 2 revealed that it shares 40.89–43.75% amino acid sequence identity with Rep-like sequences (KAH0785549, KAH0792676, KAH0797972, KAH0785997) in the genome of a parasitic protozoan, *Histomonas meleagridis*, derived from turkey (*Meleagris gallopavo*) in Austria (KAH0785549) [79]. *Histomonas meleagridis* is believed to be the cause of black histomoniasis (otherwise known as blackhead disease) in galliforms such as chickens (*Gallus*) [80]. Thus, it is likely that banfec virus 2 may infect a closely related protozoan.

### Cluster 3

Based on the Rep analysis, banfec virus 1 (1957 nt, OQ599925) identified in scat ID T05\_2F02 (Table 3) clusters with 11 other cressdnaviruses in cluster 3. This cluster includes viruses identified from oral swabs of chickens (*Gallus gallus*) from Georgia, USA (MN379584, MN379586, MN379589, MN379591) [49], a four-spotted chaser (*Libellula quadrimaculata*) from Arizona, USA (KM598400) [81], feces from a Mexican free-tailed bat (*Tadarida brasiliensis*)



**Fig. 5** Maximum-likelihood trees based on Rep sequences of unclassified cressdnavirus in cluster 2, including *Histomonas meleagridis* Rep-like sequences (inferred using the rtREV+G+I substitution model) rooted with vilyaviruses (**A**), cluster 3 (inferred using the rtREV+G+F substitution model) rooted with Rep sequences of narxaviruses (**B**), and cluster 4, including *Monocercomonoides exilis*

Rep-like sequences (inferred using the VT+G+I substitution model) rooted with Rep sequences of cycloviruses (**C**). The sequences from this study are indicated in bold blue font, and the genome organizations of the circoviruses identified in this study are shown to the right of the phylogenetic tree. Common names of the sources/hosts of the viruses are shown in parentheses next to the virus names.

in Arizona, USA (OM523005) [50], porcine serum from Brazil (MH170057) [82], dust particles from Arizona, USA (MW678880) [62], and a virus identified as a reagent contamination in a dataset of mycoviruses of a pathogenic fungus found in humans in the Netherlands (LC671629) [83] (Fig. 5). The genome of banfec virus 1 shares 69.25% sequence identity (42% coverage; E-value, 2E-52) with the genome of porcine serum associated circular DNA virus 2 - 2B/RS/BR (MH170057) from porcine serum [82]. The Rep of banfec virus 1 has 57% sequence identity to that of MT367278, a virus identified in elk feces in China (Fig. 5).

#### Cluster 4

Cluster 4 is composed of three viruses: banfec virus 4 (2913 nt, OQ599928) and banfec virus 5 (2933 nt, OQ599928), derived from scat T05\_2F02 (Table 4), and sigmofec virus UA08Rod\_11492 (3200 nt, OM869492) from a fecal sample of Arizona cotton rat (*Sigmodon arizonae*) [51]. The genomes of banfec virus 4 and 5 share ~65.7% genome-wide sequence identity, and their CPs appear to have an intron of ~220 nt (Fig. 5). They share 59.3–62.6% sequence identity with the genome of sigmofec virus UA08Rod\_11492. A BLASTp analysis of the Rep proteins of all three viruses in this cluster revealed that they share 37.35–43.38% amino acid sequence identity with Rep-like sequences (KAH7821972, KAH7821973, KAH7817465, KAH7817467) and 33.81–54.79% identity with CP-like sequences (KAH7817464, KAH7817468, KAH7817480, KAH7815554, KAH7815557) from the protozoan *Monocercomonoides exilis*, an obligate parasitic protozoan found in mammal, reptile, and insect digestive tracts [84]. *Monocercomonoides exilis* strain PA203 was isolated from a long-tailed chinchilla (*Chinchilla lanigera*) in the Czech Republic [84].

#### Anelloviridae

*Anelloviridae* is a family of single-stranded DNA viruses comprising 30 genera [85]. Their genomes range in size from 1.6 to 3.9 kb [86, 87]. Anelloviruses infect a wide variety of animals, including primates, rodents, birds, and mammalian carnivores [85]. Anellovirus genomes contain at least three large open reading frames (ORFs), and the largest of these encodes a capsid protein [87, 88]. ORF1 is the most conserved ORF in anelloviruses and thus is used in their classification. The species demarcation threshold is 69% pairwise identity in ORF1, whereas genera are distinguished based on the ORF1 amino acid sequence phylogeny [85].

The genomes of two anelloviruses, banfec anellovirus 1 (2471 nt, OQ599920) and banfec anellovirus 2 (2479 nt, OQ599921) were identified and amplified from scat sample T05\_2F02 and found to share 65.6% nucleotide sequence


identity. These two genomes share sufficient identity within the ORF1 region to be considered members of the genus *Wawtorquevirus*. The ORF1 of banfec anellovirus 1 shares 93.5% nucleotide sequence identity with neotofec virus RodL2\_5 (OM869618), whereas banfec anellovirus 2 shares 74.5% identity with neotofec virus RodL2\_6 (OM869617) (Fig. 6). Both neotofec virus RodL2\_5 and RodL2\_6 were identified in liver samples of white-throated woodrat (*Neotoma albigula*) from Arizona [51]. Based on the species demarcation criteria for anelloviruses [85], banfec anellovirus 1 and neotofec virus RodL2\_5 represent a new species, and banfec anellovirus 2 and neotofec virus RodL2\_6 represent another new species, as their ORF1 nucleotide sequences share less than 69% identity with those of any of the other wawtorqueviruses.

All but one of the wawtorqueviruses have been identified in rodent samples, which is a clear indication that banfec anellovirus 1 and 2 were most likely derived from a rodent meal. The exception within this genus, mosquito VEM anellovirus SDRB A (HQ335084) [89], was likely acquired via a blood meal from a rodent (Fig. 6). The two anelloviruses identified here were from sample T05\_2F02, in which there was evidence of Neotominae species remains, and this provides further support for the assumption that the source of these two viruses was a rodent that was consumed by the coyote.

#### Conclusion

From five fecal samples of coyotes, we identified 11 small circular DNA viruses. Of these, we identified three circoviruses representing two new species. The two banfec circovirus 1 genomes share ~97% identity and are most closely related to a circovirus in aerosolized dust from Arizona, suggesting that closely related viruses are circulating in mammals in Arizona. These three circoviruses cluster with other circoviruses from various mammals and an owl. Banfec circovirus 2 clusters with circoviruses from various rodents and carnivores (sonfela circovirus 1 from a felid and canine circovirus from canines). At this stage, we are unable to ascertain whether banfec circovirus 1 or banfec circovirus 2 actually infects coyotes or is merely associated with their diet. Screening of blood or tissue samples would presumably help to identify the tissues and organisms that are infected by these circoviruses. Banfec viruses 1–6 all represent new species of cressdnaviruses in at least five family-level groupings, highlighting the unexplored diversity of cressdnaviruses in these ecosystems. The two anelloviruses that represent two new species in the genus *Wawtorquevirus* are likely derived from rodent prey of the coyote. Furthermore, three of the viruses identified here (banfec virus 1, banfec virus 4, and banfec virus 5), based on amino acid sequence





**Fig. 6** Maximum-likelihood tree (inferred using the LG+G substitution model) based on ORF1 protein sequences of anelloviruses in the genus *Wawtorquevirus*, rooted with members of the genus *Thetatorquevirus*. The sequences from this study are indicated in bold blue

similarity to predicted proteins of two protists (*Histomonas meleagridis* and *Monocercomonoides exilis*), are likely to infect enteric protists of coyotes or their prey. Endogenized Rep-like sequences have been identified in the genomes of various organisms [90–92], suggesting virus-host interactions in their evolutionary history.

This study focused on viruses identified in coyote scats associated with a major metropolitan area; however, scat samples were collected in regions of relatively low levels of urbanization on the outskirts of a highly urbanized area (i.e., Phoenix and adjacent cities). Coyotes can travel many kilometers on a given day, and thus, the pathogens detected in this study could have originated within either lower or higher levels of urbanization. Further work could include sampling of pathogens in coyote scats across the

entire gradient of urbanization and relate viruses to specific animals with telemetry collars to track their movements and better understand how habitat use and diet relate to pathogens in wildlife.

Although we are unable to determine whether any of the cressdnnaviruses actively infect coyotes, the genome sequences of the two circoviruses can be used to develop molecular probes for screening coyote tissue or blood samples for pathogens, either in the wild by trapping or in natural history collections. Overall, using non-invasive sampling of a native predator found in urban environments, this study provides sequence information on some of the small circular DNA viruses circulating in coyotes within the Greater Phoenix region of Arizona, USA.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00705-023-05937-w>.

**Acknowledgements** We thank Angela Oreshkova and Zach Ziebarth for their assistance in collecting the scat that was used in this research. We would like to also acknowledge the support provided by the Arizona State University Natural History Collection and the Central Arizona Phoenix Long Term Ecological Research Project (grant number DEB-1832016) through the equipment and space used to enable this research. Savage Cree Hess was supported by an Arizona State University Graduate Fellowship Grant. We thank Dakota M. Rowsey (Arizona State University, USA) for the constructive feedback on the manuscript and equipment provided to process the scat for dietary analysis. We acknowledge that this research has taken place on or near the ancestral lands of 22 tribes within Arizona, such as Akimel O'odham.

**Data availability** The sequences described in this study have been deposited in the GenBank database under accession numbers OQ599920–OQ599930.

## Declarations

**Conflict of interest** The authors declare that there are no conflicts of interest.

**Research involving human participants and/or animals** The research did not involve human participants or animals. This study is a non-invasive study sampling feces of coyotes.

## References

- Bateman PW, Fleming PA, Le Comber S (2012) Big city life: carnivores in urban environments. *J Zool* 287:1–23
- Hody JW, Kays R (2018) Mapping the expansion of coyotes (*Canis latrans*) across North and Central America. *Zookeys* 759:81–97
- Snow CJ (1967) Some observations on the behavioral and morphological development of coyote pups. *Am Zool* 7:353–355
- Clark KA, Neill SU, Smith JS, Wilson PJ, Whadford VW, McKirahan GW (1994) Epizootic canine rabies transmitted by coyotes in south Texas. *J Am Vet Med Assoc* 204:536–540
- Morey PS, Gese EM, Gehrt S (2007) Spatial and temporal variation in the diet of coyotes in the Chicago metropolitan area. *Am Midl Nat* 158:147–161
- Poessel SA, Mock EC, Breck SW (2017) Coyote (*Canis latrans*) diet in an urban environment: variation relative to pet conflicts, housing density, and season. *Can J Zool* 95:287–297
- Sacks BN, Neale JCC (2002) Foraging strategy of a generalist predator toward a special prey: coyote predation on sheep. *Ecol Appl* 12:299–306
- Ma X, Bonaparte S, Corbett P, Orciari LA, Gigante CM, Kirby JD, Chipman RB, Fehlner-Gardiner C, Thang C, Cedillo VG (2023) Rabies surveillance in the United States during 2021. *J Am Vet Med Assoc* 261:1045–1053
- Krebs JW, Noll HR, Rupprecht CE, Childs JE (2002) Rabies surveillance in the United States during 2001. *J Am Vet Med Assoc* 221:1690–1701
- Blanton JD, Robertson K, Palmer D, Rupprecht CE (2009) Rabies surveillance in the United States during 2008. *J Am Vet Med Assoc* 235:676–689
- Malmlov A, Breck S, Fry T, Duncan C (2014) Serologic survey for cross-species pathogens in urban coyotes (*Canis latrans*), Colorado, USA. *J Wildl Dis* 50:946–950
- Wang X, Brown CM, Smole S, Werner BG, Han L, Farris M, DeMaria A (2010) Aggression and rabid coyotes, Massachusetts, USA. *Emerg Infect Dis* 16:357–359
- Velasco-Villa A, Orciari LA, Souza V, Juarez-Islas V, Gomez-Sierra M, Castillo A, Flisser A, Rupprecht CE (2005) Molecular epizootiology of rabies associated with terrestrial carnivores in Mexico. *Virus Res* 111:13–27
- Gese EM, Schultz RD, Johnson MR, Williams ES, Crabtree RL, Ruff RL (1997) Serological survey for diseases in free-ranging coyotes (*Canis latrans*) in Yellowstone National Park, Wyoming. *J Wildl Dis* 33:47–56
- Bischof R, Rogers DG (2005) Serologic survey of select infectious diseases in coyotes and raccoons in Nebraska. *J Wildl Dis* 41:787–791
- Guo W, Evermann JF, Foreyt WJ, Knowlton FF, Windberg LA (1986) Canine distemper virus in coyotes: a serologic survey. *J Am Vet Med Assoc* 189:1099–1100
- Allison AB, Kohler DJ, Ortega A, Hoover EA, Grove DM, Holmes EC, Parrish CR (2014) Host-specific parvovirus evolution in nature is recapitulated by in vitro adaptation to different carnivore species. *PLoS Pathog* 10:e1004475
- Canuti M, Mira F, Sorensen RG, Rodrigues B, Bouchard E, Walzthoni N, Hopson M, Gilroy C, Whitney HG, Lang AS (2022) Distribution and diversity of dog parvoviruses in wild, free-roaming and domestic canids of Newfoundland and Labrador, Canada. *Transbound Emerg Dis* 69:e2694–e2705
- Canuti M, Rodrigues B, Whitney HG, Lang AS (2017) Introduction of canine parvovirus 2 into wildlife on the Island of Newfoundland, Canada. *Infect Genet Evol* 55:205–208
- Gese EM, Grothe S (1995) Analysis of coyote predation on deer and elk during winter in Yellowstone National Park, Wyoming. *Am Midl Nat* 133:36–43
- Pluemer M, Dubay S, Drake D, Crimmins S, Veverka T, Hovanec H, Torkelson M, Mueller M (2019) Red foxes (*Vulpes vulpes*) and coyotes (*Canis latrans*) in an urban landscape: prevalence and risk factors for disease. *J Urban Ecol*. <https://doi.org/10.1093/jue/juz022>
- Brown JL, Wold B, Larson RN (2020) LA urban coyote project volunteer training. <https://doi.org/10.17504/protocols.io.933h8qn>
- Morin DJ, Higdon SD, Lonsinger RC, Gosselin EN, Kelly MJ, Waits LP (2019) Comparing methods of estimating carnivore diets with uncertainty and imperfect detection. *Wildl Soc Bull* 43:651–660
- Debelica A, Thies ML (2009) Atlas and key to the hair of terrestrial Texas mammals. Museum of Texas Tech University, Lubbock
- Larson RN, Brown JL, Karels T, Riley SP (2020) Effects of urbanization on resource use and individual specialization in coyotes (*Canis latrans*) in southern California. *PLoS ONE* 15:e0228881
- Moore TD, Spence LE, Dugnolle CE (1997) Identification of the dorsal guard hairs of some mammals of Wyoming. Wyoming Game and Fish Department
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120
- Li D, Luo R, Liu CM, Leung CM, Ting HF, Sadakane K, Yamashita H, Lam TW (2016) MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced methodologies and community practices. *Methods* 102:3–11
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410
- Zallot R, Oberg N, Gerlt JA (2019) The EFI web resource for genomic enzymology tools: leveraging protein, genome, and

- metagenome databases to discover novel enzymes and metabolic pathways. *Biochemistry* 58:4169–4182
31. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 13:2498–2504
  32. Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30:772–780
  33. Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T (2009) trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972–1973
  34. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R (2020) IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Mol Biol Evol* 37:1530–1534
  35. Letunic I, Bork P (2021) Interactive tree of life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res* 49:W293–W296
  36. Darriba D, Taboada GL, Doallo R, Posada D (2011) ProtTest 3: fast selection of best-fit models of protein evolution. *Bioinformatics* 27:1164–1165
  37. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 59:307–321
  38. Stover BC, Muller KF (2010) TreeGraph 2: combining and visualizing evidence from different phylogenetic analyses. *BMC Bioinform* 11:7
  39. Muhire BM, Varsani A, Martin DP (2014) SDT: a virus classification tool based on pairwise sequence alignment and identity calculation. *PLoS ONE* 9:e108277
  40. Krupovic M, Varsani A, Kazlauskas D, Breitbart M, Delwart E, Rosario K, Yutin N, Wolf YI, Harrach B, Zerbini FM, Dolja VV, Kuhn JH, Koonin EV (2020) Cressdnaviricota: a virus phylum unifying seven families of rep-encoding viruses with single-stranded, circular DNA genomes. *J Virol* 94:e00582–00520
  41. Ilyina TV, Koonin EV (1992) Conserved sequence motifs in the initiator proteins for rolling circle DNA replication encoded by diverse replicons from Eubacteria, Eucaryotes and Archaeobacteria. *Nucleic Acids Res* 20:3279–3285
  42. Koonin EV, Ilyina TV (1993) Computer-assisted dissection of rolling circle DNA replication. *Biosystems* 30:241–268
  43. Rosario K, Duffy S, Breitbart M (2012) A field guide to eukaryotic circular single-stranded DNA viruses: insights gained from metagenomics. *Arch Virol* 157:1851–1871
  44. Zhao L, Rosario K, Breitbart M, Duffy S (2019) Eukaryotic circular rep-encoding single-stranded DNA (CRESS DNA) viruses: ubiquitous viruses with small genomes and a diverse host range. *Adv Virus Res* 103:71–133
  45. Kazlauskas D, Varsani A, Krupovic M (2018) Pervasive chimerism in the replication-associated proteins of uncultured single-stranded DNA viruses. *Viruses* 10:187
  46. Kazlauskas D, Varsani A, Koonin EV, Krupovic M (2019) Multiple origins of prokaryotic and eukaryotic single-stranded DNA viruses from bacterial and archaeal plasmids. *Nat Commun* 10:3425
  47. Tisza MJ, Pastrana DV, Welch NL, Stewart B, Peretti A, Starrett GJ, Pang Y-YS, Krishnamurthy SR, Pesavento PA, McDermott DH, Murphy PM, Whited JL, Miller B, Brenchley J, Rosshart SP, Rehmann B, Doorbar J, Ta'ala BA, Pletnikova O, Troncoso JC, Resnick SM, Bolduc B, Sullivan MB, Varsani A, Segall AM, Buck CB (2020) Discovery of several thousand highly diverse circular DNA viruses. *Elife* 9:e51971
  48. Kinsella CM, Deijs M, Becker C, Broekhuizen P, van Gool T, Bart A, Schaefer AS, van der Hoek L (2022) Host prediction for disease-associated gastrointestinal cressdnaviruses. *Virus Evol* 8:vea087
  49. Chrzastek K, Kraberger S, Schmidlin K, Fontenele RS, Kulkarni A, Chappell L, Dufour-Zavala L, Kapczynski DR, Varsani A (2021) Diverse single-stranded DNA viruses identified in chicken buccal swabs. *Microorganisms* 9:2602
  50. Harding C, Larsen BB, Otto HW, Potticary AL, Kraberger K, Custer JM, Suazo C, Upham NS, Worobey M, van Doorslaer K, Varsani A (2022) Diverse DNA virus genomes identified in fecal samples of Mexican free-tailed bats (*Tadarida brasiliensis*) captured in Chiricahua Mountains of southeast Arizona (USA). *Virology* 580:98–111
  51. Lund MC, Larsen BB, Rowsey DM, Otto HW, Gryseels S, Kraberger S, Custer JM, Steger L, Yule KM, Harris RE, Worobey M, Van Doorslaer K, Upham NS, Varsani A (2023) Using archived and biocollection samples towards deciphering the DNA virus diversity associated with rodent species in the families Cricetidae and Heteromyidae. *Virology* 585:42–60
  52. Rosario K, Breitbart M, Harrach B, Segales J, Delwart E, Biagini P, Varsani A (2017) Revisiting the taxonomy of the family Circoviridae: establishment of the genus Cyclovirus and removal of the genus Gyrovirus. *Arch Virol* 162:1447–1463
  53. Breitbart M, Delwart E, Rosario K, Segales J, Varsani A, ICTV Report C (2017) ICTV virus taxonomy profile: Circoviridae. *J Gen Virol* 98:1997–1998
  54. Pass DA, Perry RA (1984) The pathology of psittacine beak and feather disease. *Aust Vet J* 61:69–74
  55. Meng XJ (2013) Porcine circovirus type 2 (PCV2): pathogenesis and interaction with the immune system. *Annu Rev Anim Biosci* 1:43–64
  56. Gomez-Betancur D, Vargas-Bermudez DS, Giraldo-Ramírez S, Jaime J, Ruiz-Saenz J (2023) Canine circovirus: an emerging or an endemic undiagnosed enteritis virus? *Front Vet Sci* 10:1150636
  57. Hsu HS, Lin TH, Wu HY, Lin LS, Chung CS, Chiou MT, Lin CN (2016) High detection rate of dog circovirus in diarrheal dogs. *BMC Vet Res* 12:116
  58. Kapoor A, Dubovi EJ, Henriquez-Rivera JA, Lipkin WI (2012) Complete genome sequence of the first canine circovirus. *J Virol* 86:7018
  59. Li L, McGraw S, Zhu K, Leutenegger CM, Marks SL, Kubiski S, Gaffney P, Dela Cruz FN Jr, Wang C, Delwart E, Pesavento PA (2013) Circovirus in tissues of dogs with vasculitis and hemorrhage. *Emerg Infect Dis* 19:534–541
  60. Piewbang C, Jo WK, Puff C, van der Vries E, Kesdangsakonwut S, Rungsipipat A, Kruppa J, Jung K, Baumgartner W, Techangamsuwan S, Ludlow M, Osterhaus A (2018) Novel canine circovirus strains from Thailand: evidence for genetic recombination. *Sci Rep* 8:7524
  61. Anderson A, Hartmann K, Leutenegger CM, Proksch AL, Mueller RS, Unterer S (2017) Role of canine circovirus in dogs with acute haemorrhagic diarrhoea. *Vet Rec* 180:542
  62. Finn DR, Maldonado J, de Martini F, Yu J, Penton CR, Fontenele RS, Schmidlin K, Kraberger S, Varsani A, Gile GH, Barker B, Kollath DR, Muenich RL, Herckes P, Fraser M, Garcia-Pichel F (2021) Agricultural practices drive biological loads, seasonal patterns and potential pathogens in the aerobiome of a mixed-land-use dryland. *Sci Total Environ* 798:149239
  63. Cerna GM, Serieys LEK, Riley SPD, Richet C, Kraberger S, Varsani A (2023) A circovirus and cycloviruses identified in feces of bobcats (*Lynx rufus*) in California. *Arch Virol* 168:23
  64. Wu Z, Lu L, Du J, Yang L, Ren X, Liu B, Jiang J, Yang J, Dong J, Sun L, Zhu Y, Li Y, Zheng D, Zhang C, Su H, Zheng Y, Zhou H, Zhu G, Li H, Chmura A, Yang F, Daszak P, Wang J, Liu Q, Jin Q (2018) Comparative analysis of rodent and small mammal

- viromes to better understand the wildlife origin of emerging infectious diseases. *Microbiome* 6:178
65. Legnardi M, Grassi L, Franzo G, Menandro ML, Tucciarone CM, Minichino A, Dipineto L, Borrelli L, Fioretti A, Cecchinato M (2022) Detection and molecular characterization of a novel species of circovirus in a tawny owl (*Strix aluco*) in Southern Italy. *Animals (Basel)* 12:135
  66. Wu Z, Yang L, Ren X, He G, Zhang J, Yang J, Qian Z, Dong J, Sun L, Zhu Y, Du J, Yang F, Zhang S, Jin Q (2016) Deciphering the bat virome catalog to better understand the ecological diversity of bat viruses and the bat origin of emerging infectious diseases. *ISME J* 10:609–620
  67. Payne N, Kraberger S, Fontenele RS, Schmidlin K, Bergeman MH, Cassaigne I, Culver M, Varsani A, Van Doorslaer K (2020) novel circoviruses detected in feces of sonoran felids. *Viruses* 12:1027
  68. Tang HB, Chen F, Rao G, Bai A, Jiang J, Du Y, Ren P, Liu J, Qin S, Yang L, Wu J (2017) Characterization of Akabane virus from domestic bamboo rat, Southern China. *Vet Microbiol* 207:280–285
  69. Phan TG, da Costa AC, Del Valle MJ, Bucardo-Rivera F, Nordgren J, O’Ryan M, Deng X, Delwart E (2016) The fecal virome of South and Central American children with diarrhea includes small circular DNA viral genomes of unknown origin. *Arch Virol* 161:959–966
  70. Siqueira JD, Dominguez-Bello MG, Contreras M, Lander O, Caballero-Arias H, Xutao D, Noya-Alarcon O, Delwart E (2018) Complex virome in feces from Amerindian children in isolated Amazonian villages. *Nat Commun* 9:4270
  71. Kapusinszky B, Ardeshtir A, Mulvaney U, Deng X, Delwart E (2017) Case-control comparison of enteric viromes in captive rhesus macaques with acute or idiopathic chronic diarrhea. *J Virol*. <https://doi.org/10.1128/JVI.00952-17>
  72. Pearson VM, Caudle SB, Rokyta DR (2016) Viral recombination blurs taxonomic lines: examination of single-stranded DNA viruses in a wastewater treatment plant. *PeerJ* 4:e2585
  73. de la Higuera I, Kasun GW, Torrance EL, Pratt AA, Maluenda A, Colombet J, Bisseux M, Ravet V, Dayaram A, Stainton D, Kraberger S, Zawar-Reza P, Goldstien S, Briskie JV, White R, Taylor H, Gomez C, Ainley DG, Harding JS, Fontenele RS, Schreck J, Ribeiro SG, Oswald SA, Arnold JM, Enault F, Varsani A, Stedman KM (2020) unveiling crucivirus diversity by mining metagenomic data. *MBio* 11:10–1128
  74. Dayaram A, Galatowitsch ML, Arguello-Astorga GR, van Bysterveldt K, Kraberger S, Stainton D, Harding JS, Roumagnac P, Martin DP, Lefeuvre P, Varsani A (2016) Diverse circular replication-associated protein encoding viruses circulating in invertebrates within a lake ecosystem. *Infect Genet Evol* 39:304–316
  75. Zhao M, Yue C, Yang Z, Li Y, Zhang D, Zhang J, Yang S, Shen Q, Su X, Qi D, Ma R, Xiao Y, Hou R, Yan X, Li L, Zhou Y, Liu J, Wang X, Wu W, Zhang W, Shan T, Liu S (2022) Viral metagenomics unveiled extensive communications of viruses within giant pandas and their associated organisms in the same ecosystem. *Sci Total Environ* 820:153317
  76. Wang Y, Yang S, Liu D, Zhou C, Li W, Lin Y, Wang X, Shen Q, Wang H, Li C, Zong M, Ding Y, Song Q, Deng X, Qi D, Zhang W, Delwart E (2019) The fecal virome of red-crowned cranes. *Arch Virol* 164:3–16
  77. Zhao G, Vatanen T, Droit L, Park A, Kostic AD, Poon TW, Vlamakis H, Siljander H, Harkonen T, Hamalainen AM, Peet A, Tillmann V, Ilonen J, Wang D, Knip M, Xavier RJ, Virgin HW (2017) Intestinal virome changes precede autoimmunity in type I diabetes-susceptible children. *Proc Natl Acad Sci USA* 114:E6166–E6175
  78. Orton JP, Morales M, Fontenele RS, Schmidlin K, Kraberger S, Leavitt DJ, Webster TH, Wilson MA, Kusumi K, Dolby GA, Varsani A (2020) Virus discovery in desert tortoise fecal samples: novel circular single-stranded DNA viruses. *Viruses* 12:13
  79. Palmieri N, de Jesus RM, Hess M, Bilic I (2021) Complete genomes of the eukaryotic poultry parasite *Histomonas meleagridis*: linking sequence analysis with virulence/attenuation. *BMC Genom* 22:753
  80. Beer LC, Petrone-Garcia VM, Graham BD, Hargis BM, Tellez-Isaías G, Vuong CN (2022) Histomonosis in poultry: a comprehensive review. *Front Vet Sci* 9:880738
  81. Dayaram A, Potter KA, Pailes R, Marinov M, Rosenstein DD, Varsani A (2015) Identification of diverse circular single-stranded DNA viruses in adult dragonflies and damselflies (Insecta: Odonata) of Arizona and Oklahoma, USA. *Infect Genet Evol* 30:278–287
  82. Tochetto C, Muterle Varela AP, Alves de Lima D, Loiko MR, Mengue Scheffer C, Pinto Paim W, Cerva C, Schmidt C, Cibulski SP, Cano Ortiz L, Callegari Jacques SM, Franco AC, Quos Mayer F, Roehe PM (2020) Viral DNA genomes in sera of farrowing sows with or without stillbirths. *PLoS ONE* 15:e0230714
  83. Kinsella CM, Deijs M, Gittelbauer HM, van der Hoek L, van Dijk K (2022) Human clinical isolates of pathogenic fungi are host to diverse mycoviruses. *Microbiol Spectr* 10:e0161022
  84. Karnkowska A, Vacek V, Zubacova Z, Treitli SC, Petzelkova R, Eme L, Novak L, Zarsky V, Barlow LD, Herman EK, Soukal P, Hroudova M, Dolezal P, Stairs CW, Roger AJ, Elias M, Dacks JB, Vlcek C, Hampl V (2016) A eukaryote without a mitochondrial organelle. *Curr Biol* 26:1274–1284
  85. Varsani A, Opriessnig T, Celer V, Maggi F, Okamoto H, Blomstrom AL, Cadar D, Harrach B, Biagini P, Kraberger S (2021) Taxonomic update for mammalian anelloviruses (family Anelloviridae). *Arch Virol* 166:2943–2953
  86. Biagini P (2009) Classification of TTV and related viruses (anelloviruses). *Curr Top Microbiol Immunol* 331:21–33
  87. Biagini P, Bendinelli M, Hino S, Kakkola L, Mankertz A, Niel C, Okamoto H, Raidal S, Teo CG, Todd D (2012) Family Anelloviridae. In: King AMQ, Adams EB, Carstens EB, E.J. L (eds) *Virus taxonomy: ninth report of the International Committee on taxonomy of viruses*. Academic press, London, pp 331–341
  88. Butkovic A, Kraberger S, Smelee Z, Martin DP, Schmidlin K, Fontenele RS, Shero MR, Beltran RS, Kirkham AL, Aleamotu’a M, Burns JM, Koonin EV, Varsani A, Krupovic M (2023) Evolution of anelloviruses from a circovirus-like ancestor through gradual augmentation of the jelly-roll capsid protein. *Virus Evol* 9:vead035
  89. Ng TF, Willner DL, Lim YW, Schmieder R, Chau B, Nilsson C, Anthony S, Ruan Y, Rohwer F, Breitbart M (2011) Broad surveys of DNA viral diversity obtained through viral metagenomics of mosquitoes. *PLoS ONE* 6:e20579
  90. Dennis TPW, de Souza WM, Marsile-Medun S, Singer JB, Wilson SJ, Gifford RJ (2019) The evolution, distribution and diversity of endogenous circoviral elements in vertebrate genomes. *Virus Res* 262:15–23
  91. Liu H, Li P, Liu J (2011) Numerical investigation of underlying tunnel heave during a new tunnel construction. *Tunn Undergr Space Technol* 26:276–283
  92. Patterson QM, Kraberger S, Martin DP, Shero MR, Beltran RS, Kirkham AL, Aleamotu’a M, Ainley DG, Kim S, Burns JM, Varsani A (2021) Circoviruses and cycloviruses identified in Weddell seal fecal samples from McMurdo Sound, Antarctica. *Infect Genet Evol* 95:105070
  93. Dalton CS, van de Rakt K, Fahlman A, Ruckstuhl K, Neuhaus P, Popko R, Kutz S, van der Meer F (2017) Discovery of herpesviruses in Canadian wildlife. *Arch Virol* 162:449–456
  94. Giacinti JA, Pearl DL, Ojkic D, Campbell GD, Jardine CM (2022) Genetic characterization of canine distemper virus from

- wild and domestic animal submissions to diagnostic facilities in Canada. *Prev Vet Med* 198:105535
95. Lopez-Perez AM, Moreno K, Chaves A, Ibarra-Cerdena CN, Rubio A, Foley J, List R, Suzan G, Sarmiento RE (2019) Carnivore protoparvovirus 1 at the wild-domestic carnivore interface in Northwestern Mexico. *EcoHealth* 16:502–511
  96. Condori RE, Aragon A, Breckenridge M, Pesko K, Mower K, Ettestad P, Melman S, Velasco-Villa A, Orciari LA, Yager P, Streicker DG, Gigante CM, Morgan C, Wallace R, Li Y (2022) Divergent rabies virus variant of probable bat origin in 2 gray foxes, New Mexico, USA. *Emerg Infect Dis* 28:1137–1145
  97. Pepin KM, Davis AJ, Streicker DG, Fischer JW, VerCauteren KC, Gilbert AT (2017) Predicting spatial spread of rabies in skunk populations using surveillance data reported by the public. *PLoS Negl Trop Dis* 11:e0005822
  98. Wadhwa A, Wilkins K, Gao J, Condori Condori RE, Gigante CM, Zhao H, Ma X, Ellison JA, Greenberg L, Velasco-Villa A, Orciari L, Li Y (2017) A pan-lyssavirus taqman real-time RT-PCR assay for the detection of highly variable rabies virus and other lyssaviruses. *PLoS Negl Trop Dis* 11:e0005258
  99. Kuzmin IV, Shi M, Orciari LA, Yager PA, Velasco-Villa A, Kuzmina NA, Streicker DG, Bergman DL, Rupprecht CE (2012) Molecular inferences suggest multiple host shifts of rabies viruses from bats to mesocarnivores in Arizona during 2001–2009. *PLoS Pathog* 8:e1002786
  100. Velasco-Villa A, Reeder SA, Orciari LA, Yager PA, Franka R, Blanton JD, Zuckero L, Hunt P, Oertli EH, Robinson LE, Rupprecht CE (2008) Enzootic rabies elimination from dogs and reemergence in wild terrestrial carnivores, United States. *Emerg Infect Dis* 14:1849–1854
  101. Real LA, Henderson JC, Biek R, Snaman J, Jack TL, Childs JE, Stahl E, Waller L, Tinline R, Nadin-Davis S (2005) Unifying the spatial population dynamics and molecular evolution of epidemic rabies virus. *Proc Natl Acad Sci USA* 102:12107–12111
  102. Nadin-Davis SA, Abdel-Malik M, Armstrong J, Wandeler AI (2002) Lyssavirus P gene characterisation provides insights into the phylogeny of the genus and identifies structural similarities and diversity within the encoded phosphoprotein. *Virology* 298:286–305
  103. Kuzmina NA, Kuzmin I, Ellison J, Rupprecht CE (2013) Conservation of binding epitopes for monoclonal antibodies on the rabies virus glycoprotein. *J Antivir Antiretrovir* 5:37–43

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.



## Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

[onlineservice@springernature.com](mailto:onlineservice@springernature.com)