


Antler Casting Phenology and Late-Breeding in Nebraska White-Tailed Deer

Poster · February 2021  
DOI: 10.3390/RS13.0206.4953


CITATIONS  
1

5 authors, including:



**Brian C. Peterson**  
University of Nebraska at Kearney  
27 PUBLICATIONS 207 CITATIONS  
[SEE PROFILE](#)

READS  
63



**Felix G. Farrell**  
45 PUBLICATIONS 87 CITATIONS  
[SEE PROFILE](#)





# Antler Casting Phenology and Occurrence of Late-breeding in Nebraska White-tailed Deer

Brian C. Peterson<sup>1</sup>, Patrick Farrell<sup>2</sup>,  
Dave Fehlhafer<sup>3</sup>, Miranda Reinson<sup>1</sup>, Dustin H. Ranglack<sup>1</sup>  
University of Nebraska at Kearney<sup>1</sup> Headwaters Corporation<sup>2</sup> Ecological Solutions<sup>3</sup>



## INTRODUCTION


- Antler casting can be variable, influenced by a variety of factors (e.g., photoperiod, geography, sex ratios, hormone levels and body condition).
- In Nebraska, WTD primarily breed in November, but this season extends into December and January for unfertilized females and healthy fawns reaching appropriate weight.
- Documented historical antler casting timeframes in Nebraska are inconsistent: Dec-Jan, Jan-Feb, Jan-May.....

## OBJECTIVES

- Document observations of early and late antler casting
- Establish a baseline for antler casting phenology
- Describe a seasonally late-breeding observation within the central Platte River valley

## METHODS

- Our study area is best described as the big bend reach of the central Nebraska Platte River Valley (2009-Present).




- We compiled opportunistic and systematic observations to quantify antler casting timeframe for this region.
- Records of cast antlered males and fresh antler casts were used to develop a conservative casting timeframe.
- We investigated regional camera trap captures to determine cast antler phenology between November and April.
- Additionally, we detailed late-breeding and late antler casting by a male WTD in this region.

## RESULTS

- Several cast antlers were observed (harvested or collected) in early January with the earliest cast collected 21 Dec 2020.
- Late antler casting was observed by four males ( $\geq 2.5$ -yr-olds) after 14 Apr, the latest observation occurring 20 Apr 2011.
- 22.3% of freshly cast antlers contained pedicle blood (77/346).
- Average collection date of casts with pedicle blood was 4 days earlier (9 Mar) than non-bloody (13 Mar) pedicle casts ( $P=0.03$ ).



- Based on our camera capture effort, the mean casting timeframe occurred the first fifteen days of March with 81% of males casting both antlers, and 99% of antlers cast by the start of April.



## Late-Breeding Observation (Figures below)

- We observed a mature male mounting and copulating with a young female on 5 April 2019.
- The following morning, we observed the same male scenting checking and chasing a group of young females.
- Video and pictures revealed protracted and enlarged testes.
- This individual retained his antlers until a least April 14<sup>th</sup>.



## DISCUSSION

- We determined that casting timeframe was broad, and later, on average than previously documented for this region.
- We can only speculate on causality of the late-breeding and antler casting observations (estrous female/elevated test.)
- Our observations serve as a baseline for future spatial and temporal comparisons and may assist state biologists and property managers as they assess future regulations.

## ACKNOWLEDGMENTS

We thank the landowners and organizations that permitted access during or study:



The Animal Care and Use Committee at the University of Nebraska at Kearney reviewed and approved for field study exemption (IACUC #090821).