

Structural bioinformatics

NMRe: a web server for NMR protein structure refinement with high-quality structure validation scores

Hyojung Ryu^{1,2}, GyuTae Lim^{1,2}, Bong Hyun Sung³ and Jinhyuk Lee^{1,2,*}

¹Korean Bioinformation Center (KOBIC), Korea Research Institute of Bioscience and Biotechnology, Yuseong-gu, Daejeon 305-806, the Republic of Korea, ²Department of Nanobiotechnology and Bioinformatics, University of Science and Technology, Yuseong-Gu, Daejeon 305-350, the Republic of Korea and ³Bioenergy and Biochemical Research Center, Korea Research Institute of Bioscience and Biotechnology, Yuseong-gu, Daejeon 305-806, the Republic of Korea

*To whom correspondence should be addressed.

Associate Editor: Anna Tramontano

Received on June 9, 2015; revised on September 23, 2015; accepted on October 10, 2015

Abstract

Summary: Protein structure refinement is a necessary step for the study of protein function. In particular, some nuclear magnetic resonance (NMR) structures are of lower quality than X-ray crystallographic structures. Here, we present NMRe, a web-based server for NMR structure refinement. The previously developed knowledge-based energy function STAP (Statistical Torsion Angle Potential) was used for NMRe refinement. With STAP, NMRe provides two refinement protocols using two types of distance restraints. If a user provides NOE (Nuclear Overhauser Effect) data, the refinement is performed with the NOE distance restraints as a conventional NMR structure refinement. Additionally, NMRe generates NOE-like distance restraints based on the inter-hydrogen distances derived from the input structure. The efficiency of NMRe refinement was validated on 20 NMR structures. Most of the quality assessment scores of the refined NMR structures were better than those of the original structures. The refinement results are provided as a three-dimensional structure view, a secondary structure scheme, and numerical and graphical structure validation scores.

Availability and implementation: NMRe is available at <http://psb.kobic.re.kr/nmre/>

Contact: jinhyuk@kribb.re.kr

Supplementary information: [Supplementary data](#) are available at *Bioinformatics* online.

1 Introduction

The determination of three-dimensional (3D) protein structures is an important challenge in structural biology because structure is closely related to function. NMR (Nuclear Magnetic Resonance) spectroscopy is one method of protein structure determination, and it also provides information about the dynamics of proteins. However, protein structures determined by NMR spectroscopy are of lower quality than X-ray structures. Thus, refinement is a necessary step in NMR structure study. Three NMR structure refinement databases have been constructed: DRESS (a Database of REfined Solution NMR Structures; [Nabuurs et al., 2004](#)), RECOORD

(REcalculated COORDinates Database; [Nederveen et al., 2005](#)) and STAP (Statistical Torsion Angle Potential; [Yang et al., 2012](#)). NMR structure refinement using the NOE (Nuclear Overhauser Effect) distance data obtained by NMR is performed in many refinement studies ([Bertini et al., 2011](#); [Chen et al., 2004](#); [Mao et al., 2014](#)). However, NOE distance data exist in various formats, such as CYANA, CNS and XPLOR. Therefore, NOE data manipulation is difficult for general researchers. Another problem in improving NMR structures is the ambiguity and sparseness of NOE data ([Nilges, 1997](#)). The NMRe web server provides simple and fast NMR structure refinement with NOE-like distance data using the

References

- Bertini, I. *et al.* (2011) A grid-enabled web portal for NMR structure refinement with AMBER. *Bioinformatics*, **27**, 2384–2390.
- Brooks, B.R. *et al.* (2009) CHARMM: the biomolecular simulation program. *J. Comput. Chem.*, **30**, 1545–1614.
- Chen, J. *et al.* (2004) Refinement of NMR structures using implicit solvent and advanced sampling techniques. *J. Am. Chem. Soc.*, **126**, 16038–16047.
- Kim, T.R. *et al.* (2013) Statistical torsion angle potential energy functions for protein structure modeling: a bicubic interpolation approach. *Proteins*, **81**, 1156–1165.
- Lazaridis, T. and Karplus, M. (1999) Effective energy function for proteins in solution. *Proteins*, **35**, 133–152.
- Mao, B. *et al.* (2014) Protein NMR structures refined with Rosetta have higher accuracy relative to corresponding X-ray crystal structures. *J. Am. Chem. Soc.*, **136**, 1893–1906.
- Nabuurs, S.B. *et al.* (2004) DRESS: a database of REfined solution NMR structures. *Proteins*, **55**, 483–486.
- Nederveen, A.J. *et al.* (2005) RECOORD: a recalculated coordinate database of 500+ proteins from the PDB using restraints from the BioMagResBank. *Proteins*, **59**, 662–672.
- Nilges, M. (1997) Ambiguous distance data in the calculation of NMR structures. *Fold. Des.*, **2**, S53–S57.
- Ryu, H. *et al.* (2014) Protein NMR structures refined without NOE data. *PLoS One*, **9**, e108888.
- Yang, J.S. *et al.* (2012) STAP refinement of the NMR database: a database of 2405 refined solution NMR structures. *Nucleic Acids Res.*, **40**, D525–D530.