

The Xeno-glycomics database (XDB): a relational database of qualitative and quantitative pig glycome repertoire

Hae-Min Park¹, Ju-Hyeong Park², Yoon-Woo Kim³, Kyoung-Jin Kim³, Hee-Jin Jeong¹, Kyoung-Soon Jang¹, Byung-Gee Kim^{1,4,5,*} and Yun-Gon Kim^{3,*}

¹School of Chemical and Biological Engineering and ²Department of Computer Science and Engineering, Seoul National University, Seoul 151-742, Korea ³Department of Chemical Engineering, Soongsil University, Seoul 156-743, Korea ⁴Institute of Molecular Biology and Genetics and ⁵Institute of Bioengineering, Seoul National University, Seoul 151-742, Korea

Associate Editor: John Hancock

ABSTRACT

Summary: In recent years, the improvement of mass spectrometry-based glycomics techniques (i.e. highly sensitive, quantitative and high-throughput analytical tools) has enabled us to obtain a large dataset of glycans. Here we present a database named Xeno-glycomics database (XDB) that contains cell- or tissue-specific pig glycomes analyzed with mass spectrometry-based techniques, including a comprehensive pig glycan information on chemical structures, mass values, types and relative quantities. It was designed as a user-friendly web-based interface that allows users to query the database according to pig tissue/cell types or glycan masses. This database will contribute in providing qualitative and quantitative information on glycomes characterized from various pig cells/organs in xenotransplantation and might eventually provide new targets in the α 1,3-galactosyltransferase gene-knock out pigs era.

Availability: The database can be accessed on the web at <http://bioinformatics.snu.ac.kr/xdb>.

Contact: byungkim@snu.ac.kr or ygkim@ssu.ac.kr

Supplementary information: Supplementary data are available at *Bioinformatics* online.

Received on April 5, 2013; revised on August 7, 2013; accepted on August 26, 2013

1 INTRODUCTION

Cell surface carbohydrates can act as ligand molecules for diverse biological processes such as cell–cell interaction, signal transduction, metastasis and so forth (Ohtsubo and Marth, 2006; Rudd *et al.*, 2001). Especially in pig-to-human xenotransplantation, the interaction between terminal α -Gal (Gal α 1-3Gal-R) antigens on pig tissues or cells' surfaces and circulating natural human antibodies lead to hyper acute immune rejection (Oriol *et al.*, 1993). Despite the development of α 1,3-galactosyltransferase gene-knock out pigs, unidentified non-Gal antigens remain as possible factors triggering delayed xenograft rejection (Ezzelarab *et al.*, 2005). Thus, the comprehensive identification of α -Gal and non-Gal carbohydrate antigens from pig tissues and cells is essential for successful xenotransplantation.

During the past decades, several databases have been developed to assist the assignment, annotation and interpretation of glycan and glycoconjugates datasets (Campbell, *et al.*, 2011; Campbell *et al.*, 2008; Cooper *et al.*, 2003; Kameyama *et al.*, 2005; Raman *et al.*, 2006; von der Lieth *et al.*, 2011). However, none of them systematically classify and update the pig glycans according to the specific pig tissues and cells. Qualitative and quantitative information on the glycan antigens, which play an important role in immune rejection, is also lacking.

Here, we have built up a relational database [Xeno-glycomics database (XDB)] from specific pathogen-free Chicago Medical School miniature pig (Setcavage and Kim, 1976) glycan profiles, which have been qualitatively and quantitatively characterized by various mass spectrometry-based techniques (Supplementary Data). It includes *N*-glycomes from pig kidney (Kim *et al.*, 2008, 2006), endothelial cells, islets (Kim *et al.*, 2009a), corneal endothelial cells, keratocytes (Kim *et al.*, 2009b), heart and glycosphingolipid-derived glycans from pig endothelial cells and islets (Kim *et al.*, 2009c). The *O*-glycomes from pig kidney, endothelial cells (Park *et al.*, 2013) and heart (accepted) are also included. This specialized pig glycome database will offer a new insight into pig-to-human xenotransplantation as well as into many different areas of biological and biomedical research.

2 METHOD

XDB is based on open-source technology, developed in jruby 1.7.0 and trinidad 1.4.4 Tomcat 7.0.37 and tested on Debian. The database was used in SQLite version 3.0 as a data storage program. The HTML-based web interface has been tested using Safari 6 and Chrome 22. The updated version of web browsers is recommended for the usage of the XDB.

3 RESULTS AND DISCUSSION

Two major query options are implemented in XDB, namely, 'pig tissue/cell specific glycans' and 'mass spectrometry-based glycan searching' (Fig. 1, Supplementary Data). The 'pig tissue/cell specific glycans' option allows the retrieval of glycans derived from a specific pig tissue/cell. Users can obtain the full lists of glycans by clicking a specific pig organ or cell icon. It is also possible to set parameters including glycan epitope (Kawasaki *et al.*, 2006). On the other hand, the 'mass spectrometry-based glycan searching' is a way to look for a specific glycan information from

*To whom correspondence should be addressed.




Fig. 1. The query and result pages of (A) 'pig tissue/cell' specific glycans and (B) 'mass spectrometry-based glycan searching' in XDB

a mass-to-charge ratio (m/z). If the users input the experimental mass value (m/z), the program enables the users to gather the possible glycan structures and organs. In addition, the XDB allows users to set a mass tolerance that can be specified in Daltons \pm of the measured value to avert no search result found because of instrumental variations. The users can also narrow down the list of the glycans by specifying several factors. It includes the followings: mass spectrometry instrument, glycan derivatization method, glycan type, ion mode, ion adducts and origin, but it will be updated continuously.

The search result from XDB contains mass value (m/z), derivatization, glycan type, structure, antigen, quantity and origin. The mass value is an experimental value of the corresponding glycan mass (mono-isotopic) analyzed using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry or electrospray ionization mass spectrometry. The derivatization indicates the chemical derivatization for glycan analysis using mass spectrometry. All the glycans in the XDB are classified into three types: N-linked glycans (at the side chain of Asn), O-linked glycans (at the Ser/Thr residues) in glycoproteins and glycosphingolipid-derived glycans. Especially, the α -galactosylated glycans are labeled with ' α -Gal antigen' in the column of 'Antigen type'. The chemical structures were drawn with Oxford glycan nomenclature (Harvey *et al.*, 2009). The relative quantity of glycans, which was measured by NP-HPLC or quantitative MS analysis, was incorporated in the result table as well. Additionally, pie graphs represent the relative quantities of each type of glycans (e.g. high-mannose glycans, α -Gal glycans, sialylated glycans and so on).

Although many of the well-established glycan databases offer the detailed glycan structures, it is difficult to solely collect the pig glycans according to the specific organ or cell, which is a promising candidate for xenotransplantation. Therefore, we

have built a unique and specialized database to provide reliable information on glycans, which are derived from tissues and cells of pigs. Moreover, it contains both qualitative and quantitative information on xenoantigenic glycans, which play an important role in immune-rejection response. Taken together, this database is potentially promising and facilitates xenotransplantation research as well as basic carbohydrate research. The XDB will allow users to use hyperlink the human glycan database with CFG, which contains glycan profiles in human cells and tissues for pig-to-human xenotransplantation research. In the near future, this XDB will be updated with a list of glycans from α 1,3- galactosyltransferase gene-knock out pig samples as well.

Funding: This work was supported by Converging Research Center Program funded by the Ministry of Education, Science and Technology through the National Research Foundation of Korea (NRF)(2012K001387). This work was also supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) (2013R1A1A1004998) and the KIST Institutional Program (2E23720-13-053).

Conflict of Interest: none declared.

REFERENCES

- Campbell, M.P. *et al.* (2008) GlycoBase and autoGU: tools for HPLC-based glycan analysis. *Bioinformatics*, **24**, 1214–1216.
- Campbell, M.P. *et al.* (2011) UniCarbKB: putting the pieces together for glycomics research. *Proteomics*, **11**, 4117–4121.
- Cooper, C.A. *et al.* (2003) GlycoSuiteDB: a curated relational database of glycoprotein glycan structures and their biological sources. 2003 update. *Nucleic Acids Res.*, **31**, 511–513.
- Ezzelarab, M. *et al.* (2005) Carbohydrates in xenotransplantation. *Immunol. Cell Biol.*, **83**, 396–404.
- Harvey, D.J. *et al.* (2009) Proposal for a standard system for drawing structural diagrams of N- and O-linked carbohydrates and related compounds. *Proteomics*, **9**, 3796–3801.
- Kameyama, A. *et al.* (2005) A strategy for identification of oligosaccharide structures using observational multistage mass spectral library. *Anal. Chem.*, **77**, 4719–4725.
- Kawasaki, T. *et al.* (2006) GlycoEpitope: the integrated database of carbohydrate antigens and antibodies. *Trends Glycosci. Glyc.*, **18**, 267–272.
- Kim, Y.G. *et al.* (2006) The identification and characterization of xenoantigenic nonhuman carbohydrate sequences in membrane proteins from porcine kidney. *Proteomics*, **6**, 1133–1142.
- Kim, Y.G. *et al.* (2008) Structural analysis of α -Gal and new non-Gal carbohydrate epitopes from specific pathogen-free miniature pig kidney. *Proteomics*, **8**, 2596–2610.
- Kim, Y.G. *et al.* (2009a) Qualitative and quantitative comparison of N-glycans between pig endothelial and islet cells by high-performance liquid chromatography and mass spectrometry-based strategy. *J. Mass Spectrom.*, **44**, 1087–1104.
- Kim, Y.G. *et al.* (2009b) Mass spectrometric analysis of the glycosphingolipid-derived glycans from miniature pig endothelial cells and islets: identification of NeuGc epitope in pig islets. *J. Mass Spectrom.*, **44**, 1489–1499.
- Kim, Y.G. *et al.* (2009c) Identification of α -Gal and non-Gal epitopes in pig corneal endothelial cells and keratocytes by using mass spectrometry. *Curr. Eye Res.*, **34**, 877–895.
- Ohtsubo, K. and Marth, J.D. (2006) Glycosylation in cellular mechanisms of health and disease. *Cell*, **126**, 855–867.
- Oriol, R. *et al.* (1993) Carbohydrate antigens of pig tissues reacting with human natural antibodies as potential targets for hyperacute vascular rejection in pig-to-man organ xenotransplantation. *Transplantation*, **56**, 1433–1442.
- Park, H.M. *et al.* (2013) Structural characterization of α -galactosylated O-glycans from miniature pig kidney and endothelial cells. *Carbohydr. Res.*, **369**, 48–53.

- Raman,R. *et al.* (2006) Advancing glycomics: implementation strategies at the consortium for functional glycomics. *Glycobiology*, **16**, 82R–90R.
- Rudd,P.M. *et al.* (2001) Glycosylation and the immune system. *Science*, **291**, 2370–2376.
- Setcavage,T.M. and Kim,Y.B. (1976) Variability of the immunological state of germ-free colostrum-deprived Minnesota miniature piglets. *Infect. Immun.*, **13**, 600–607.
- von der Lieth,C.W. *et al.* (2011) EUROCarbDB: an open-access platform for glycoinformatics. *Glycobiology*, **21**, 493–502.