Systems biology

Advance Access publication January 30, 2014

CoNtRol: an open source framework for the analysis of chemical reaction networks

Pete Donnell^{1,*}, Murad Banaji¹, Anca Marginean² and Casian Pantea³

¹Department of Mathematics, University of Portsmouth, Portsmouth PO1 3HF, UK, ²Computer Science Department, Technical University of Cluj-Napoca, 400027 Cluj-Napoca, Romania and ³Department of Mathematics, West Virginia University, Morgantown, WV 26506-6310, USA

Associate Editor: Alfonso Valencia

ABSTRACT

Summary: We introduce CoNtRol, a web-based framework for analysis of chemical reaction networks (CRNs). It is designed to be both extensible and simple to use, complementing existing CRN-related tools. CoNtRol currently implements a number of necessary and/or sufficient structural tests for multiple equilibria, stable periodic orbits, convergence to equilibria and persistence, with the potential for incorporation of further tests.

Availability and implementation: Reference implementation: reaction-networks.net/control/. Source code and binaries, released under the GPLv3: reaction-networks.net/control/download/. Documentation: reaction-networks.net/wiki/CoNtRol.

Contact: pete.donnell@port.ac.uk

Received on November 18, 2013; revised on January 21, 2014; accepted on January 24, 2014

1 INTRODUCTION

Differential equation models of chemical reaction networks (CRNs), encompassing many ecological, epidemiological and related non-negative systems, are almost universally complicated by parameter uncertainty. However, dynamical properties of large classes of CRN models are remarkably robust to changes in parameter values, leading to a range of results relating network structure to dynamical behavior. Such parameter-free approaches to the analysis of CRNs fall into the scope of *chem*ical reaction network theory (Horn and Jackson, 1972; Feinberg, 1972, 1979). Fuelled in part by its implications to systems biology (Bailey, 2001; Shinar and Feinberg, 2010), chemical reaction network theory has seen a surge of interest in recent years, attacking questions about multistationarity (Craciun et al., 2006; Conradi et al., 2007; Banaji et al., 2007; Banaji and Craciun, 2009; Shinar and Feinberg, 2012), global stability (Craciun et al., 2009; Angeli et al., 2010; Anderson, 2011; Donnell and Banaji, 2013), oscillatory behavior (Angeli et al., 2014) and persistence (Angeli et al., 2007; Pantea, 2012).

Many of the results in chemical reaction network theory lend themselves to algorithmic implementation, which is useful for large networks. In this article, we introduce CoNtRol (CRN tool), a new, fully open source platform, currently coded in C, Java, Octave and PHP, to perform computations on CRNs without the need for any proprietary software. CoNtRol has a

web-based front-end interfacing with a suite of modular tests, to which users may add new tests in any language. With its array of features, CoNtRol complements existing software tools. The Chemical Reaction Network Toolbox (Ellison et al., 2011) is a proprietary Windows program that implements the results of deficiency theory (Feinberg, 1979), as well as checks for concordance (Shinar and Feinberg, 2012), equivalent to network injectivity. It applies to various types of kinetics, including mass action, and generates detailed information about network properties, multiple and degenerate steady states and their stability. Reaction Network Equilibria Study Toolbox (ERNEST) (Soranzo and Altafini, 2009) and CRNreals (Szederkényi et al., 2012) are MATLAB toolboxes aimed at testing multistationarity, and distinguishability and identifiability of CRNs, respectively. Tests for multistationarity of CRNs are also implemented in Maple (Feliu and Wiuf, 2013). While not all criteria included in these tools are currently tested in CoNtRol, its extensible nature allows for their future inclusion.

2 FEATURES AND FUNCTIONALITY

CoNtRol has an intuitive user interface where a list of reactions can be input manually or uploaded in a variety of formats. While the focus so far has been on developing the framework for extensible functionality, a number of tools for the analysis of CRNs have also been implemented and tested.

2.1 Network analysis

The current functionality of CoNtRol includes a number of necessary and/or sufficient structural tests for multiple equilibria, stable oscillation, convergence to equilibria and persistence, assuming mass action or more general kinetics. In particular, the following are checked: sufficient conditions for convergence to equilibria based on the theory of monotone dynamical systems (Angeli et al., 2010; Donnell and Banaji, 2013); conditions of the deficiency zero and deficiency one theorems (Feinberg, 1987), resulting in claims about local stability of equilibria, absence of multiple positive equilibria and absence of periodic orbits, primarily for mass action kinetics; structural conditions for persistence (Angeli et al., 2007; Donnell and Banaji, 2013) based on examining the siphons of the system; and a large number of sufficient/necessary conditions for injectivity and absence of multistationarity gathered from the literature and developed in (Banaji and Pantea, 2013). The outputs are cross-referenced

^{*}To whom correspondence should be addressed.

to the documentation of CoNtRol, where each conclusion and its implications are described in detail. Some of the multistationarity results of CoNtRol are similar to those of the Chemical Reaction Network Toolbox (Ellison *et al.*, 2011).

2.2 DSR graph generation

A useful tool in the parameter-free study of CRNs is the directed species-reaction (DSR) graph: a bipartite graph with signed, labeled and sometimes directed edges. CoNtRol includes a Java Web Start application for drawing DSR graphs. The graph layout is highly customizable, and the 'export to LATEX' feature generates TikZ code for the resulting DSR diagram. An example is illustrated in Figure 1. Easily checked structural conditions on the DSR graph can imply, or contribute toward, powerful conclusions regarding multistationarity, asymptotic stability or oscillatory behavior in a CRN (Craciun et al., 2006; Banaji and Craciun, 2009; Angeli et al., 2010, 2014; Shinar and Feinberg, 2013). Several of these are implemented or lend themselves naturally to future implementation.

2.3 Batch processing

This feature allows users to analyze a number of networks in a single run, thus generating statistics about the occurrence of certain behaviors. An archive containing individual files describing each network is uploaded, and the user is emailed a link to the output once it is complete. For example, Table 1 summarizes the results of some injectivity tests implemented in CoNtRol, applied to the database of all 4082 non-isomorphic bimolecular CRNs with three species and three reactions (Deckard *et al.*, 2009).

2.4 Miscellaneous

A number of features that further improve usability have been implemented. These include multiple input formats, the option to receive the output via email and automatic generation of LATEX code (e.g. list of reactions, stoichiometric matrix).




Fig. 1. Futile cycle DSR graph $\begin{cases} E+S & \rightleftharpoons & ES \rightarrow & E+P \\ F+P & \rightleftharpoons & FP \rightarrow & F+S \end{cases}$

Table 1. Injectivity of 3 species-3 reaction bimolecular reactions

$GK\alpha$	$GK\beta$	$MA\alpha$	$MA\beta$
44%	45%	45%	47%

GK, general kinetics; MA, mass action kinetics; α , the system is injective on the relative interior of each non-trivial stoichiometry class; β , the fully open system is injective on the non-negative orthant. For example, with mass action kinetics, 45% (MA α) of the networks was injective on the relative interior on each non-trivial stoichiometric class, only slightly more than with general kinetics (GK α). See (Banaji and Pantea, 2013) for more details.

3 CONCLUSION

We have written CoNtRol aiming for a user-friendly tool with maximal functionality, implementing state-of-the-art theory on CRNs. We expect future inclusion of further results from the rich theory of CRNs, with contributions from the research community.

Funding: C.P.'s work was funded by Leverhulme Trust grant F/07 058/BU. M.B and P.D.'s work was supported by EPSRC grant EP/J008826/1.

Conflict of Interest: none declared.

REFERENCES

Anderson, D. (2011) A proof of the global attractor conjecture in the single linkage class case. SIAM J. Appl. Math., 71, 1487–1508.

Angeli, D. et al. (2007) A Petri net approach to the study of persistence in chemical reaction networks. Math. Biosci., 210, 598–618.

Angeli, D. et al. (2010) Graph-theoretic characterizations of monotonicity of chemical networks in reaction coordinates. J. Math. Biol., 61, 581–616.

Angeli, D. et al. (2014) Combinatorial approaches to Hopf bifurcations in systems of interacting elements. Commun. Math. Sci., (in press).

Bailey, J.E. (2001) Complex biology with no parameters. Nat. Biotechnol., 19, 503–504.

Banaji, M. et al. (2007) P matrix properties, injectivity and stability in chemical reaction systems. SIAM J. Appl. Math., 67, 1523–1547.

Banaji, M. and Craciun, G. (2009) Graph-theoretic approaches to injectivity and multiple equilibria in systems of interacting elements. *Commun. Math. Sci.*, 7, 867–900.

Banaji, M. and Pantea, C. (2013) Some Results on Injectivity and Multistationarity in Chemical Reaction Networks. http://arXiv.org/abs/1309.6771.

Conradi, C. et al. (2007) Subnetwork analysis reveals dynamic features of complex (bio) chemical networks. PNAS, 104, 19175–19180.

Craciun, G. et al. (2006) Understanding bistability in complex enzyme-driven reaction networks. PNAS, 103, 8697–8702.

Craciun, G. et al. (2009) Toric dynamical systems. J. Symb. Comput., 44, 1551–1565.Deckard, A. et al. (2009) Enumeration and Online Library of Mass-Action Reaction Networks, http://arXiv.org/abs/0901.3067.

Donnell, P. and Banaji, M. (2013) Local and global stability of equilibria for a class of chemical reaction networks. SIAM J. Appl. Dyn. Syst., 12, 899–920.

Ellison, P. et al. (2011) The Chemical Reaction Network Toolbox ver. 2.2a. http://www.crnt.osu.edu/CRNTWin

Feinberg,M. (1972) Complex balancing in general kinetic systems. Arch. Ration. Mech. Anal., 49, 187–194.

Feinberg,M. (1979) Lectures on Chemical Reaction Networks (Lectures delivered at the Mathematics Research Center, University of Wisconsin-Madison). http://www.chbmeng.ohio-state.edu/~FEINBERG/LecturesOnReactionNetworks/.

Feinberg, M. (1987) Chemical reaction network structure and the stability of complex isothermal reactors—I: the deficiency zero and deficiency one theorems. Chem. Eng. Sci., 42, 2229–2268.

Feliu, E. and Wiuf, C. (2013) A computational method to preclude multistationarity in networks of interacting species. *Bioinformatics*, **29**, 2327–2334.

Horn, F. and Jackson, R. (1972) General mass action kinetics. Arch. Ration. Mech. Anal., 47, 81–116.

Pantea, C. (2012) On the persistence and global stability of mass-action systems. SIAM J. Math. Anal., 44, 1636–1673.

Shinar, G. and Feinberg, M. (2010) Structural sources of robustness in biochemical reaction networks. Science. 327, 1389.

Shinar, G. and Feinberg, M. (2012) Concordant chemical reaction networks. Math. Biosci., 240, 92–113.

Shinar, G. and Feinberg, M. (2013) Concordant chemical reaction networks and the species-reaction graph. *Math. Biosci.*, 241, 1–23.

Soranzo, N. and Altafini, C. (2009) ERNEST: a toolbox for chemical reaction network theory. *Bioinformatics*, 25, 2853–2854.

Szederkényi,G. et al. (2012) CRNreals: a toolbox for distinguishability and identifiability analysis of biochemical reaction networks. Bioinformatics, 28, 1549–1550.