

Name: _____

Student number: _____

Chemistry 1A03

Test 2

Nov 13, 2015

McMaster University

VERSION 1

Instructors: D. Brock, G. Goward, A. Hitchcock, L. Davis

Duration: 120 minutes

This test contains 20 numbered pages printed on both sides. There are **30** multiple-choice questions appearing on pages numbered 3 to 17. Page 18 is extra space for rough work. Page 19 includes some useful data and equations, and there is a periodic table on page 20. You may tear off the last page to view the periodic table and the data provided.

You must enter your name and student number on this question sheet, as well as on the answer sheet. Your invigilator will be checking your student card for identification.

You are responsible for ensuring that your copy of the question paper is complete. Bring any discrepancy to the attention of your invigilator.

All questions are worth 2 marks - the total marks available are 60. There is **no** penalty for incorrect answers.

BE SURE TO ENTER THE CORRECT VERSION OF YOUR TEST (shown near the top of page 1), IN THE SPACE PROVIDED ON THE ANSWER SHEET.

ANSWER ALL QUESTIONS ON THE ANSWER SHEET, IN PENCIL.

Instructions for entering multiple-choice answers are given on page 2.

SELECT ONE AND ONLY ONE ANSWER FOR EACH QUESTION from the answers (A) through (E). **No work written on the question sheets will be marked.** The question sheets may be collected and reviewed in cases of suspected academic dishonesty.

Academic dishonesty may include, among other actions, communication of any kind (verbal, visual, *etc.*) between students, sharing of materials between students, copying or looking at other students' work. If you have a problem please ask the invigilator to deal with it for you. Do not make contact with other students directly. Try to keep your eyes on your own paper – looking around the room may be interpreted as an attempt to copy.

Only Casio FX 991 electronic calculators may be used; but they must NOT be transferred between students. Use of any aids other than those provided, is not allowed.

1. What is the **pH** of a 0.0342 M solution of HCl?

- A) 1.466
- B) 4.456
- C) 7.568
- D) 3.492
- E) 1.081

2. Which of the following statements about periodic trends are **TRUE**?

- i) The correct sequence for decreasing ionic radius is: $\text{Br}^- > \text{Rb}^+ > \text{Sr}^{2+}$.
- ii) The ground-state electron configuration of Si has no unpaired electrons.
- iii) The oxide of calcium is a basic oxide.
- iv) Rb loses electrons more easily than Na.
- v) The electronegativity of chlorine is smaller than that of phosphorus.

- A) ii, v
- B) i, iv
- C) iii, iv, v
- D) i, iii, iv
- E) i, ii, v

3. Which of the following statement(s) are **FALSE** ?

- i) Sodium has a larger first ionization energy than potassium.
- ii) Sodium has a larger atomic size than chlorine.
- iii) Oxygen has a larger first ionization energy than nitrogen.
- iv) Sulfur has a larger electronegativity than chlorine.

- A) ii, iv
- B) ii
- C) i, ii
- D) iii, iv
- E) i, iii

4. When preparing a 0.100 mol L^{-1} solution of NaOH, instead of weighing out 1.00 kg of NaOH and diluting it with the appropriate amount of water, the individual unknowingly used KOH by accident. How would this affect the **concentration** of the solution and the **volume** of base needed to titrate an acid?

<u>Concentration of Base Solution</u> <u>(relative to 0.100 mol L^{-1})</u>	<u>Volume of base needed to reach</u> <u>equivalence point</u>
A) more concentrated	more
B) less concentrated	less
C) more concentrated	less
D) same concentration	same volume
E) less concentrated	more

Name: _____

Student number: _____

5. What is the **concentration (in mol L⁻¹)** of an 8.00 mass % aqueous solution of NH₃, which has a density of 0.9651 g mL⁻¹?

- A) 12.9
- B) 4.53
- C) 0.264
- D) 7.55
- E) 18.2

6. The threshold wavelength of photons that are able to eject electrons from Cesium metal is 500. nm. Which of the following statements are **FALSE**?

- i) The energy of each photon of 500. nm light is 3.97×10^{-19} J.
- ii) Shorter wavelengths of light will also eject electrons from Cesium.
- iii) If ionization occurs, the number of electrons ejected is directly proportional to the brightness of the incident light.
- iv) If the energy of the incident photons is doubled, the electron speed increases by a factor of four.
- v) If the incident wavelength is tuned to 400. nm, the speed of the electrons is 8.67×10^5 ms⁻¹.

- A) ii, v
- B) i, iii
- C) i, iv
- D) iv, v
- E) ii, iii

Name: _____

Student number: _____

7. Rank the following species in **increasing terminal Se-O bond order** (from lowest to highest). (H atoms are attached to O)

SeO_2F^- , SeO_3 , HSeO_4^-

- A) $\text{HSeO}_4^- < \text{SeO}_3 < \text{SeO}_2\text{F}^-$
B) $\text{HSeO}_4^- < \text{SeO}_2\text{F}^- < \text{SeO}_3$
C) $\text{SeO}_3 < \text{SeO}_2\text{F}^- < \text{HSeO}_4^-$
D) $\text{SeO}_2\text{F}^- < \text{SeO}_3 < \text{HSeO}_4^-$
E) $\text{SeO}_2\text{F}^- < \text{HSeO}_4^- < \text{SeO}_3$

8. How many **resonance structures** does HSeO_4^- have? (H atom is attached to O)

- A) 6
B) 1
C) 3
D) 2
E) 4

Name: _____

Student number: _____

9. Which VSEPR class(es) would have **non-ideal** bond angles?

- A) AX_4E_2
- B) AX_3
- C) AX_2E_3
- D) AX_4E
- E) More than one of the above.

10. Which of the following has an atom with **less than an octet**?

- i) NO
 - ii) BCl_3
 - iii) CO_2
- A) i, iii
 - B) i, ii
 - C) ii, iii
 - D) all of the above
 - E) none of the above

Name: _____

Student number: _____

11. In class we discussed that methane clathrate ice is **unusual**. If water was frozen after exposing it to the following compounds, which compound would **MOST LIKELY** be found in the water lattice?

- A) F_2
- B) CH_2Cl_2
- C) CO_2
- D) SiO_2
- E) O_2


12. How many **non-bonding** electrons are there in SeO_3 ?

- A) 14
- B) 16
- C) 12
- D) 10
- E) 18

13. What is the **average formal charge** on the atoms in SeO_2F^- ?

	<u>Se</u>	<u>O</u>	<u>F</u>
A)	0	-1	0
B)	+1	-1/2	0
C)	+1	-1	-1
D)	0	-1/2	0
E)	0	-1/3	-1/3

14. Which of the following perturbations will shift the given equilibrium **toward products**?



- i) The temperature is increased.
- ii) Some NO_2 is removed from the reaction mixture.
- iii) The reaction mixture is transferred to a vessel with twice the volume.
- iv) The O_3 partial pressure is increased.
- v) The total pressure is increased.

- A) iii, v
- B) i, iv
- C) ii, iv
- D) i, v
- E) ii, iii

Name: _____

Student number: _____

15. AgCl(s) has a K_{sp} of 1.8×10^{-10} . Consider a system in which AgCl(s) is added to a solution of 1.0×10^{-3} M KCl. When equilibrium is established, what is the **concentration of $\text{Ag}^+(\text{aq})$** in moles per litre?






- A) 5.5×10^{-7}
- B) 3.9×10^{-9}
- C) 1.8×10^{-7}
- D) 2.5×10^{-3}
- E) 7.2×10^{-6}

16. **Persistent organic pollutants** are characterized by which of the following:


- i) high vapour pressure
- ii) resistance to chemical change
- iii) large octanol-water partition coefficient
- iv) high electron affinity
- v) electron pair geometry

- A) iv, v
- B) ii, iii
- C) i, iii
- D) i, iv
- E) ii, v

17. Which one of the following is **NOT** the correct form of the **equilibrium constant** for the given equilibrium?



18. Find the **equilibrium partial pressure of IBr**, for the equilibrium between:



The initial pressure of IBr is 0.017 bar, and the value of K is 8.5×10^{-3} .

- A) 0.081
- B) 0.0055
- C) 0.0094
- D) 0.014
- E) 0.0021

Name: _____

Student number: _____

19. Determine the **molar solubility** of CaF_2 . $K_{sp} = 3.45 \times 10^{-11}$

- A) 2.38×10^{-2}
- B) 1.69×10^{-3}
- C) 2.05×10^{-4}
- D) 4.25×10^{-8}
- E) 8.43×10^{-4}

20. Which of the following set(s) of observations is/are **FALSE** for the reaction products of experiment #2, the cycles of copper.


- i) $\text{Cu}(\text{NO}_3)_2$, aqueous
- ii) CuSO_4 , precipitate
- iii) CuO , precipitate
- iv) $\text{Cu}(\text{OH})_2$, precipitate
- v) NO_2 , gas

- A) iii, iv
- B) ii
- C) iv, v
- D) iv
- E) i, v

Name: _____

Student number: _____

21. Consider the reaction between water vapour and carbon monoxide, where each are initially present at a 1.0 bar. The value of K for the reaction at a certain temperature, T , is 0.63. What is the **equilibrium partial pressure of carbon dioxide** in bar?



- A) 0.59
- B) 0.44
- C) 0.25
- D) 0.32
- E) 0.67

22. Consider a weak acid, HA with $K_a = 1.0 \times 10^{-4}$. Which of the following is **FALSE** with respect to a 1.0 M solution of HA.

- A) $\text{p}K_b$ of the conjugate base = $-\log(10^{-10})$
- B) $[\text{OH}^-]$ in solution is 10^{-10} M
- C) $K_a < [\text{H}_3\text{O}^+]$ in solution.
- D) The small x approximation is valid when solving for the pH.
- E) The conjugate base is a weak base.

Name: _____

Student number: _____

23. Which of the following is **FALSE** with regard to acid strength?

- A) $\text{HF} > \text{HI}$
- B) $\text{HBr} > \text{H}_2\text{S}$
- C) $\text{HClO}_3 > \text{HClO}_2$
- D) $\text{H}_2\text{SO}_3 > \text{H}_3\text{PO}_3$
- E) $\text{H}_2\text{O} > \text{NH}_3$

24. What is the **pH** of a solution that is comprised of 2.0 L of water and 2.0 g of NaOCl?
 $K_a [\text{HOCl}] = 2.9 \times 10^{-8}$

- A) 5.26
- B) 8.11
- C) 8.59
- D) 10.24
- E) 9.83

Name: _____

Student number: _____

25. Determine the **FALSE** statement:

- A) A Brønsted base is a proton acceptor.
- B) H_2F^+ would be a strong acid.
- C) 1 mole of $\text{Mg}(\text{OH})_2$ would react with 2 moles of HCl to produce a neutral solution.
- D) NH_4^+ is the conjugate acid of NH_3 .
- E) F^- is a Lewis acid

26. If the pH of a solution is 12.100, what is the **concentration of OH^-** (mol L^{-1}) in the solution?

- A) 0.106
- B) 0.259
- C) 0.0984
- D) 0.0548
- E) 0.0126

Name: _____

Student number: _____

27. Determine the **FALSE** statement below.

- A) NaH is a strong base.
- B) When dissolved in water, CH_3NH_2 will produce OH^- .
- C) When dissolved in water, ClOH would produce a solution with $\text{pH} > 7$.
- D) Both HI and HCl are strong acids.
- E) H_2CO_3 is not a strong acid

28. A solution of ethylamine ($\text{CH}_3\text{CH}_2\text{NH}_2$; $K_b = 4.30 \times 10^{-4}$) produces a $\text{pH} = 12.67$. What is the **% ionization** of the base?

- A) 2.2 %
- B) 0.054 %
- C) 0.92 %
- D) 0.0015%
- E) 1.5 %

Name: _____

Student number: _____

29. When a particular gas expands against a constant external pressure of 2.50 atm, the volume increases by 9.75 L. During this transformation the gas also absorbs 1000. J of heat. What is the energy change, ΔU (in kJ), for the gas?

- A) +3.47
- B) +4.39
- C) -3.47
- D) +1.47
- E) -1.47

30. Which one of the following reactions has **no work done ON or BY** the system?

- A) $2 \text{NO}_2(\text{g}) + 7 \text{H}_2(\text{g}) \rightarrow 2 \text{NH}_3(\text{g}) + 4 \text{H}_2\text{O}(\text{l})$
- B) $\text{PCl}_5(\text{g}) \rightarrow \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$
- C) $\text{Br}_2(\text{g}) \rightarrow 2 \text{Br}(\text{g})$
- D) $2 \text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2 \text{SO}_3(\text{g})$
- E) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2 \text{NO}(\text{g})$

Name: _____

Student number: _____

Extra space for rough work:

Name: _____

Student number: _____

- Some general data are provided on this page.
- A Periodic Table with atomic weights is provided on the next page.

STP = 273.15 K, 1 atm

 $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ $h = 6.6256 \times 10^{-34} \text{ Js}$ density(H_2O , l) = 1.00g/mL

Specific heat of water = 4.184 J / g·°C

 $R = 8.3145 \text{ J K}^{-1} \text{ mol}^{-1} = 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1} = 0.083145 \text{ L bar K}^{-1} \text{ mol}^{-1}$ $F = 96485 \text{ C/mol}$ $c = 2.9979 \times 10^8 \text{ m/s}$ $m_e = 9.109 \times 10^{-31} \text{ kg}$ $\Delta H^\circ_{\text{vap}}[\text{H}_2\text{O}] = 44.0 \text{ kJ mol}^{-1}$

1 bar = 100.00 kPa = 750.06 mm Hg = 0.98692 atm

0°C = 273.15 K

1 J = 1 kg m² s⁻² = 1 kPa L = 1 Pa m³1 m = 10⁶ μm = 10⁹ nm = 10¹⁰ Å1 cm³ = 1 mL1 g = 10³ mg

1 Hz = 1 cycle/s

De Broglie wavelength:

$$\lambda = h / mu = h / p$$

Hydrogen atom energy levels:

$$E_n = -R_H / n^2 = -2.179 \times 10^{-18} \text{ J} / n^2$$

$$KE = \frac{1}{2}mu^2$$

Nernst Equation:

$$E = E^\circ - \frac{RT}{zF} \ln Q = E^\circ - \frac{0.0257 \text{ V}}{z} \ln Q = E^\circ - \frac{0.0592 \text{ V}}{z} \log_{10} Q$$

Entropy change: $\Delta S = \frac{q_{\text{rev}}}{T}$

Solubility Guidelines for Common Ionic Solids

Follow the lower-numbered guideline when two guidelines are in conflict. This leads to the correct prediction in most cases.

1. Salts of group 1 cations and the NH_4^+ cation are soluble . Except LiF and Li_2CO_3 which are insoluble.
2. Nitrates, acetates, bicarbonates, and perchlorates are soluble.
3. Salts of silver, lead and mercury (I) are insoluble. Except AgF which is soluble.
4. Fluorides, chlorides, bromides, and iodides are soluble. Except Group 2 fluorides which are insoluble
5. Carbonates, phosphates, chromates, sulfides, oxides, and hydroxides are insoluble. Except Group 2 sulfides and hydroxides of Ca^{2+} , Sr^{2+} , and Ba^{2+} which are soluble.).
6. Sulfates are soluble except for those of calcium, strontium, and barium.

