




6.3 Graphing Using Intercepts Worksheet #2


Math 9

1. Identify the x- and y-intercepts of each graph.




2. For each of the following, plot the intercepts and graph the line.

	x-intercept	y-intercept
a)	4	7
b)	-3	1
c)	1	-4
d)	-5.5	-2




3. Determine the x- and y-intercepts and use them to graph each line.


a) $2x + 5y = 10$




b) $-2x - 3y = 12$




c) $3x + 6y = -9$




d) $4x - y = 6$



e) $3y - 7x = 21$




f) $y - x = 11$




4. Draw a graph and determine the slope of each line whose x- and y-intercepts are given.


a) x-intercept = 2; y-intercept = 4




b) x-intercept = -3; y-intercept = 5




c) x-intercept = 6; y-intercept = -3



d) no x-intercept; y-intercept = 3



5. The distance-time graph shows Kyla's motion in front of a motion sensor.



a) What is the d-intercept?

b) What is the t-intercept?


c) What information does each intercept give about Kyla's motion?

d) Use the graph or the slope formula to determine Kyla's speed. Remember the speed is the slope on a distance-time graph.

6. A candle burns at a constant rate of 2 cm/h. The candle is 12 cm tall when it is first lit.

a) State the dependent and the independent variables.

b) Set up a graph of length, l , in centimeters versus time, t , in hours.



c) Plot points to represent the length of candle remaining after 0, 1, 2 hours.

d) Join the points and extend the line so that it crosses the t-axis.

e) What is the l – *intercept*? What information does it give?

f) What is the t – *intercept*? What information does it give?

g) Why does this linear model have no meaning below the t -axis?

h) Why does this linear model have no meaning to the left of the l -axis?