

BEE 4750/5750 Homework 1

Jason Shao (jls647)

2022-09-15

Problem 1

Problem 1.1

```
julia> using GraphRecipes, Plots

julia> A = [0 1 1 1;
            0 0 0 1;
            0 0 0 1;
            0 0 0 0];

julia> names = ["Plant", "Land Treatment", "Chem Treatment", "Pristine Brook"];

julia> shapes=[:hexagon, :rect, :rect, :hexagon];

julia> xpos = [0, -2, 2, 0];

julia> ypos = [1, 0, 0, -1];

julia> edgelabels=Array{String}(undef,4,4);

julia> for i=1:4
    for j=1:4
        edgelabels[i,j]=string("edgelabel");
    end
end

julia> edgelabels[1,2]="X1*1kg/m^3";


julia> edgelabels[1,3]="X2*1kg/m^3";

julia> edgelabels[1,4]="100-(X1*1kg/m^3)-(X2*1kg/m^3)";

julia> edgelabels[2,4]=".2*(X1*1kg/m^3)";

julia> edgelabels[3,4]="(.005*X2)*(X2*1kg/m^3)";

julia> graphplot(A, names=names, markersize=0.15, edgelabel=edgelabels,
smarkersshapes=shapes, markercolor=:white, x=xpos, y=ypos)
```



Model of mass [kg/day] moving between each location. X_1 and X_2 in m^3/day

Problem 1.2

The plant originally starts off discharging 100 kg/day, so the rate of mass discharge starts off with this value

$$d(X_1, X_2) = 100 - \dots \left[\frac{kg}{day} \right]$$

next you can subtract the amounts X_1 and X_2 since the original concentration is $1 \frac{kg}{m^3}$

$$d(X_1, X_2) = 100 - X_1 * 1 \frac{kg}{m^3} - X_2 * 1 \frac{kg}{m^3} \dots \left[\frac{kg}{day} \right]$$

then you have to take into account the efficiency of the treatments and add back mass

$$d(X_1, X_2) = 100 - X_1 - X_2 + .2X_1 + X_2(.005X_2) \left[\frac{kg}{day} \right]$$

This rearranges to

$$d(X_1, X_2) = -.8X_1 + .005X_2^2 - X_2 + 100 \quad \left[\frac{\text{kg}}{\text{day}}\right]$$

The cost is equal to the sum of both treatments

$$C(X_1, X_2) = \frac{X_1^2}{20} + 1.5X_2 \quad [\text{USD}]$$

Problem 1.3

```
julia> function ConcCost(X1,X2)
    d=(-.8*X1)+(.005*X2^2)-(X2)*(100)
    C=X1^2/20+1.5*X2
    return d,C
end
ConcCost (generic function with 1 method)
```

Problem 1.4

```
julia> using Plots, Distributions

julia> d=zeros(100,100);

julia> C=zeros(100,100);


julia> a,b = ConcCost(0,0);

julia> for i=1:100
    for j=1:100
        d[i,j]=a;
        C[i,j]=b;
    end
end

julia> for i=1:100
    for j=1:101-i
        a,b = ConcCost(i-1,j);
        d[i,j]=a;
        C[i,j]=b;
    end
end

julia> scatter(C, d, xguide="Cost (USD)",yguide="Daily Effluent YUK Mass
Discharge (kg/day)", legend=false, markercolor=:blue);

julia> hline!([20])
```



Problem 1.5

Using the plot from 1.4, different treatment plans can be made by envisioning the priorities of different groups of interest. The factory owners might use the cheapest treatment plan that is under 20 kg/day. They could look at the plot and identify the leftmost point that is under the horizontal line at 20 kg/day and find the corresponding X_1 and X_2 combination. If it were up to the public or regulatory agencies, they might opt for a plan which costs more but will be safer. I iterated through all integer values of X_1 and X_2 that add up to 100 in order to get these points. This means that the cheapest point in the plot that meets the standard might not be the optimal minimizing cost plan. If I had included decimal values then there may be a cheaper option that still meets the standard that I would have chosen.

Problem 1.6

One thing that can be looked further into is what the actual optimal treatment plan is that minimizes cost. As mentioned earlier, the treatment plans I plotted have X_1 and X_2 take integer values only. Knowing this assumption, the results cannot be interpreted as the true optimal treatment plan that minimizes cost, as the actual optimal treatment plan might not have whole numbers. This optimal treatment plan could be found by minimizing cost with effluent mass discharge as a constraint.

Problem 1.7

References