

BEE 4750/5750 Homework 2

Jason Shao (jls647)

2022-09-28

Problem 1

Problem 1.1

julia> #A function which takes distance x in km, and initial DO, BOD, and NOD concentrations and outputs the DO concentration at that distance

```
function C(x, Co, CBODo, NBODo)
    #given values
    ka=.55
    kc=.35
    kn=.25
    Cs=10
    U=6

    #alpha value calculations
    a1=exp(-ka*x/U)
    a2=(kc/(ka-kc))*(exp(-kc*x/U)-exp(-ka*x/U))
    a3=(kn/(ka-kn))*(exp(-kn*x/U)-exp(-ka*x/U))
    DO=Cs*(1-a1)+Co*a1-CBODo*a2-NBODo*a3
    return DO
end
```

C (generic function with 1 method)

julia> #Decay of CBOD function

```
function CBOD(x, CBODo)
    kc=.35
    U=6
    CBOD=CBODo*exp(-kc*x/U)
    return CBOD
end
```

CBOD (generic function with 1 method)

julia> #Decay of NBOD function

```
function NBOD(x, NBODo)
    kn=.35
    U=6
    NBOD=NBODo*exp(-kn*x/U)
    return NBOD
end
```

NBOD (generic function with 1 method)

To find initial concentration of DO:

$$C_{o1} = \frac{C_{River} * Q_{River} + C_{Waste1} * Q_{Waste1}}{Q_{River} + Q_{Waste1}}$$

$$C_{o1} = \frac{7.5 \frac{mg}{L} * 10^8 \frac{L}{d} + 5 \frac{mg}{L} * 10^7 \frac{L}{d}}{10^8 \frac{L}{d} + 10^7 \frac{L}{d}} = 7.27 \frac{mg}{L}$$

The same process is done for CBOD and NBOD at the start of the first inflow and again at the second waste flow

$$CBOD_{o1} = \frac{5 \frac{mg}{L} * 10^8 \frac{L}{d} + 50 \frac{mg}{L} * 10^7 \frac{L}{d}}{10^8 \frac{L}{d} + 10^7 \frac{L}{d}} = 9.09 \frac{mg}{L}$$

$$NBOD_{o1} = \frac{5 \frac{mg}{L} * 10^8 \frac{L}{d} + 35 \frac{mg}{L} * 10^7 \frac{L}{d}}{10^8 \frac{L}{d} + 10^7 \frac{L}{d}} = 7.72 \frac{mg}{L}$$

```
julia> using Plots, Distributions

julia> DO=zeros(51); #initialize DO vector

julia> DO[1]=7.27 ; #set initial DO concentration with both river inflow and
waste source 1

julia> for i=1:14
    DO[i+1]=C(i,7.27,9.09,7.72); #calculate DO concentration for 1 to 14 km
downstream
end

julia> #calculate initial DO, BOD, and NOD concentrations of inflow diverging
with waste source 2
    CBODo2=(1.1*10^8*CBOD(15,9.09)+45*1.5*10^7)/(1.25*10^8);

julia> NBODo2=(1.1*10^8*NBOD(15,7.72)+35*1.5*10^7)/(1.25*10^8);


julia> DO[16]=(C(15,7.27,9.09,7.72)*1.1*10^8+1.5*10^7*5)/(1.25*10^8);

julia> #using these initial values, calculate the DO concentration after waste
source 2
    for i=16:50
        DO[i+1]=C(i-15,DO[16],CBODo2,NBODo2);
```

end

```
julia> x=0:1:50; #vector of distance values
```

```
julia> plot(x,DO, xguide="Distance downstream from first waste source, x  
(km)", yguide= "DO concentration (mg/L)", legend=false)
```




Problem 1.2

Using the conditions at the input of the second waste stream as the initial conditions to find DO concentration, you can solve for the distance which achieves 6 mg/L. Since the equation seems difficult to solve algebraically, you can find the distance finding the root of $C(x)-6$. This can be one using the Roots package in Julia. A plot with a horizontal line at $DO=6\text{mg/L}$ is shown to provide an initial guess of x .

```
julia> using Roots
```

```
julia> plot(x,DO, xguide="Distance downstream from first waste source, x  
(km)", yguide= "DO concentration (mg/L)", legend=false);
```

```
julia> hline!([6])
```



```
julia> ka=.55;
julia> kc=.35;
julia> kn=.25;
julia> Cs=10;
julia> U=6;
julia> Co=D0[16];
julia> CBODo=(1.1*10^8*CBOD(15,9.09)+45*1.5*10^7)/(1.25*10^8);
julia> NBODo=(1.1*10^8*NBOD(15,7.72)+35*1.5*10^7)/(1.25*10^8);
julia>
f(x)=(Cs*(1-(exp(-ka*x/U)))+Co*(exp(-ka*x/U))-CBODo*((kc/(ka-kc))*(exp(-kc*x/U)-exp(-ka*x/U)
f (generic function with 1 method)

julia> #redefine function with only one input, x
      find_zero(f, 30)
25.01130063650589

julia> minimum(D0)
3.9066566044718316
```

Finding the root of the equation shows that at 25.0 km downstream from waste stream 2, the dissolved oxygen concentration rebounds back to 6 mg/L.

Problem 1.3

In order to find the treatment needed to prevent the dissolved oxygen concentration from dropping below 4 mg/L, you can test a range of treatment percentages and finding the treatment which has a minimum DO level just at or above 4 mg/L.

```
julia> function C(x, Co, CBODo, NBODo)
```

```
    #given values
```

```
    ka=.55
```

```
    kc=.35
```

```
    kn=.25
```

```
    Cs=10
```

```
    U=6
```

```
    #alpha value calculations
```

```
    a1=exp(-ka*x/U)
```

```
    a2=(kc/(ka-kc))*(exp(-kc*x/U)-exp(-ka*x/U))
```

```
    a3=(kn/(ka-kn))*(exp(-kn*x/U)-exp(-ka*x/U))
```

```
    DO=Cs*(1-a1)+Co*a1-CBODo*a2-NBODo*a3
```

```
    return DO
```

```
end
```

```
C (generic function with 1 method)
```

```
julia> #Decay of CBOD function
```

```
function CBOD(x, CBODo)
```

```
    kc=.35
```

```
    U=6
```

```
    CBOD=CBODo*exp(-kc*x/U)
```

```
    return CBOD
```

```
end
```

```
CBOD (generic function with 1 method)
```

```
julia> #Decay of NBOD function
```

```
function NBOD(x, NBODo)
```

```
    kn=.35
```

```
    U=6
```

```
    NBOD=NBODo*exp(-kn*x/U)
```

```
    return NBOD
```

```
end
```

```
NBOD (generic function with 1 method)
```

```
julia> Co=DO[16]
```

```
4.897609716593195
```

```
julia> minConc=zeros(20);
```

```
julia> treatment=zeros(20);
```

```
julia> for x=1:20
```

```
    D0treat=zeros(45);
```

```
    CBODo=(1.1*10^8*CBOD(15,9.09)+45*(1-x*.01)*1.5*10^7)/(1.25*10^8);
```

```
    NBODo=(1.1*10^8*NBOD(15,7.72)+35*(1-x*.01)*1.5*10^7)/(1.25*10^8);
```

```
    for i=1:45
```

```
        D0treat[i]=C(i, Co, CBODo, NBODo)
```

```
    end
```

```
    minConc[x]=minimum(D0treat);
```

```
    treatment[x]=x*.01
```

end

```
julia> minConc
20-element Vector{Float64}:
 3.927451121268696
 3.9482456380655586
 3.9690401548624235
 3.989834671659286
 4.01062918845615
 4.0314237052530135
 4.052218222049878
 4.0730127388467405
 4.0930141861362666
 4.112150237577911
 4.131286289019555
 4.1504223404611995
 4.169558391902845
 4.188694443344488
 4.207830494786133
 4.226966546227778
 4.246102597669422
 4.265238649111067
 4.2843747005527115
 4.303510751994356
```

```
julia> treatment
20-element Vector{Float64}:
 0.01
 0.02
 0.03
 0.04
 0.05
 0.06
 0.07
 0.08
 0.09
 0.1
 0.11
 0.12
 0.13
 0.14
 0.15
 0.16
 0.17
 0.18
 0.19
 0.2
```

A treatment of 5% removal will achieve a minimum DO concentration of 4.01 mg/L, just over 4 mg/L.

Problem 1.4

Now the same can be done with treatment done at both waste streams. Treatment done at the first waste stream will affect the initial DO, CBOD, and NBOD concentrations at

the second waste stream.

```
julia> function C(x, Co, CBODo, NBODo)
```

```
    #given values
```

```
    ka=.55
```

```
    kc=.35
```

```
    kn=.25
```

```
    Cs=10
```

```
    U=6
```

```
    #alpha value calculations
```

```
    a1=exp(-ka*x/U)
```

```
    a2=(kc/(ka-kc))*(exp(-kc*x/U)-exp(-ka*x/U))
```

```
    a3=(kn/(ka-kn))*(exp(-kn*x/U)-exp(-ka*x/U))
```

```
    DO=Cs*(1-a1)+Co*a1-CBODo*a2-NBODo*a3
```

```
    return DO
```

```
end
```

```
C (generic function with 1 method)
```

```
julia> #Decay of CBOD function
```

```
function CBOD(x, CBODo)
```

```
    kc=.35
```

```
    U=6
```

```
    CBOD=CBODo*exp(-kc*x/U)
```

```
    return CBOD
```

```
end
```

```
CBOD (generic function with 1 method)
```

```
julia> #Decay of NBOD function
```

```
function NBOD(x, NBODo)
```

```
    kn=.35
```

```
    U=6
```

```
    NBOD=NBODo*exp(-kn*x/U)
```

```
    return NBOD
```

```
end
```

```
NBOD (generic function with 1 method)
```

```
julia> minConc=zeros(11);
```

```
julia> treatment=zeros(11);
```

```
julia> for x=0:10
```

```
    DOTreat=zeros(45);
```

```
    CBODo1=(5*10^8+50*(1-x*.01)*10^7)/(1.1*10^8);
```

```
    NBODo1=(5*10^8+35*(1-x*.01)*10^7)/(1.1*10^8);
```

```
    Co2=(C(15,7.27,CBODo1,NBODo1)*1.1*10^8+1.5*10^7*5)/(1.25*10^8);
```

```
    CBODo2=(1.1*10^8*CBOD(15,CBODo1)+45*(1-x*.01)*1.5*10^7)/(1.25*10^8);
```

```
    NBODo2=(1.1*10^8*NBOD(15,NBODo1)+35*(1-x*.01)*1.5*10^7)/(1.25*10^8);
```

```
    for i=1:45
```

```
        DOTreat[i]=C(i,Co2,CBODo2,NBODo2)
```

```
    end
```

```
    minConc[x+1]=minimum(DOTreat);
```

```
    treatment[x+1]=x*.01
```

```
end
```

```
julia> minConc
```

```
11-element Vector{Float64}:
```

```
 3.905171614769302
```

```

3.9416618343430034
3.9781520539167032
4.014642273490404
4.051132493064105
4.087622712637804
4.1241129322115055
4.160603151785207
4.197093371358905
4.233583590932606
4.2700738105063065

```

```

julia> treatment
11-element Vector{Float64}:
 0.0
 0.01
 0.02
 0.03
 0.04
 0.05
 0.06
 0.07
 0.08
 0.09
 0.1

```

A 3% treatment of both streams achieves a minimum DO concentration of 4.01 mg/L.

Problem 1.5

I would treat each waste stream equally. This is because both waste streams contribute to the depletion of dissolved oxygen. While waste stream 2 brings the DO concentration to its minimum point, this minimum is still affected by the DO depletion from waste stream 1. Additional information may change this strategy however. For example, if the cost for treatment is in USD/mg/d, treating only the second waste stream would be less costly.

Problem 1.6

To estimate the probability that

Problem 1.7

Problem 1.8

'''

References