

BEE 4750 Homework 1: Introduction to Using Julia

Name:

ID:

Due Date

Thursday, 9/11/25, 9:00pm

Overview

Instructions

- Problem 1 consist of a series of code snippets for you to interpret and debug. You will be asked to identify relevant error(s) and fix the code.
- Problem 2 asks you to write code to implement several simple functions using more general programming and some which are unique to Julia (which might mean that you need to look at the documentation or use Julia's help function, which is accessed with `?`, e.g. `?sum` for help with the `sum()` function) or to analyze Julia syntax.
- Problem 3 asks you to convert a verbal description of a wastewater treatment system into a Julia function, and then to use that function to explore the impact of different wastewater allocation strategies.

Load Environment

The following code loads the environment and makes sure all needed packages are installed. This should be at the start of most Julia scripts.

```
import Pkg
Pkg.activate(@__DIR__)
Pkg.instantiate()
```

Activating project at `~/BEE 4750/bee4750-hw1-pdc76`

Standard Julia practice is to load all needed packages at the top of a file. If you need to load any additional packages in any assignments beyond those which are loaded by default, feel free to add a `using` statement, though [you may need to install the package](#).

```
using Random
using Plots
using GraphRecipes
using LaTeXStrings
using Distributions
```

```
# this sets a random seed, which ensures reproducibility of random number generation. You sh
Random.seed!(1)
```

TaskLocalRNG()

Problems (Total: 30 Points)

Problem 1 (12 points)

The following subproblems all involve code snippets that require debugging. You are encouraged to use online resources (*e.g.* Julia documentation and forums, Stack Overflow, Reddit, etc) to help find diagnose error messages and find solutions, but make sure that you clearly document which resources you used and how you used them.

Problem 1.1

You've been tasked with writing code to identify the minimum value in an array. You cannot use a predefined function. Your colleague suggested the function below, but it does not return the minimum value.

```
function minimum(array)
    # initialize the minimum value counter
    min_value = 0
    # update minimum values
    for i in 1:length(array)
        if array[i] < min_value
            min_value = array[i]
        end
    end
end
```

```

    end
    # return found minimum
    return min_value
end

array_values = [89, 90, 95, 100, 100, 78, 99, 98, 100, 95]
@show minimum(array_values);

```

What is the logical flaw in the code? Fix it and show that your fixed code solves the problem.

Solution to Problem 1.1

The issue with the original code is that the value, 0, held by the variable, “min_value,” was less than the minimum value in the array, “array_values.” To address this, I initialized the min_value variable to be 1/0, which Julia interprets to be infinity. This guarantees that any value in the list will be less than or equal to the originally initialized minimum value. Therefore, the function “minimum” will always return the correct value minimum value of an array. The corrected code is below.

```

function minimum(array)
    # initialize the minimum value counter
    min_value = 1/0
    # update minimum values
    for i in 1:length(array)
        if array[i] < min_value
            min_value = array[i]
        end
    end
    # return found minimum
    return min_value
end

array_values = [89, 90, 95, 100, 100, 78, 99, 98, 100, 95]
@show minimum(array_values);

```

Problem 1.2

Your team is trying to compute the average grade for your class, but the following code produces an `UndefVarError`.

```
# enter student grade vector
student_grades = [89, 90, 95, 100, 100, 78, 99, 98, 100, 95]
# compute class average
function class_average(grades)
    average_grade = mean(student_grades)
    return average_grade
end

@show average_grade;
```

What does this `UndefVarError` mean? Why does this code produce one? Fix the error and solve the problem.

Problem 1.2 Solution

`UndefVarError` indicates an issue with global and local variables within a function. Since “average_grade” is only defined within the function, “class_average,” you cannot show “average_grade” without defining it in the global space. To correct this, you must define “average_grade” by calling “class_average” with the array “student_grades.” The corrected code is below:

```
# enter student grade vector
student_grades = [89, 90, 95, 100, 100, 78, 99, 98, 100, 95]
# compute class average
function class_average(grades)
    average_grade = mean(student_grades)
    return average_grade
end

average_grade = class_average(student_grades)
@show average_grade;
```

Problem 1.3

Your team wants to know the expected payout of an old Italian dice game called *passadieci* (which was analyzed by Galileo as one of the first examples of a rigorous study of probability). The goal of *passadieci* is to get at least an 11 from rolling three fair, six-sided dice. Your strategy is to compute the average wins from 1,000 trials, but the code you’ve written below produces a `MethodError`.

```

# function to simulate a passadieci roll: 3 6-sided dice
function passadieci()
    # this rand() call samples 3 values from the vector [1, 6]
    roll = rand(1:6, 3)
    return roll
end
# set number of trials and initialize outcome vector
n_trials = 1_000
outcomes = zero(n_trials)
# simulate number of passadieci rolls and count wins
for i = 1:n_trials
    outcomes[i] = (sum(passadieci()) > 11)
end
win_prob = sum(outcomes) / n_trials # compute average number of wins
@show win_prob;

```

```

MethodError: MethodError: no method matching setindex!{::Int64, ::Bool, ::Int64}
The function `setindex!` exists, but no method is defined for this combination of argument types.
MethodError: no method matching setindex!{::Int64, ::Bool, ::Int64}

```

```

The function `setindex!` exists, but no method is defined for this combination of argument types.

```

Stacktrace:

```
[1] top-level scope
```

```
@ ~ /BEE 4750/bee4750-hw1-pdc76/jl_notebook_cell_df34fa98e69747e1a8f8a730347b8e2f_X22sZmls
```

What does this `MethodError` mean? Why does this code produce one? Fix the error and solve the problem.

Problem 1.3 Solution

A `MethodError` is an error where a function is passed an argument that it cannot handle or recognize. This occurs in 1.3 when the variable `outcomes` is created by calling the function `zero()`. It is passed an argument that is an integer, which is correct, but the output of this is a single integer, 0. `0[i]` is not valid syntax, so `outcomes[i]` throws a method error. To fix this, we use the function `zeros()` which creates an array of zeros with length of the integer argument. See corrected code below:

```

# function to simulate a passadieci roll: 3 6-sided dice
function passadieci()
    # this rand() call samples 3 values from the vector [1, 6]
    roll = rand(1:6, 3)
    return roll
end
# set number of trials and initialize outcome vector
n_trials = 1_000
outcomes = zeros(n_trials)
# simulate number of passadieci rolls and count wins
for i = 1:n_trials
    outcomes[i] = (sum(passadieci()) > 11)
end
win_prob = sum(outcomes) / n_trials # compute average number of wins
@show win_prob;

```

```
win_prob = 0.369
```

Problem 1.4

You're interested in writing some code to remove the mean of a vector from all of its components. You've written the following code and tried to test it on a random vector, but your code returns a `MethodError`.

```

# function to remove mean from a vector
function remove_mean(vect)
    # fuction to compute the mean
    function compute_mean(vect)
        element_sum = 0 # initialize sum
        # compute mean and return
        for v in vect
            element_sum += v
        end
        return element_sum / length(vect)
    end

    m = compute_mean(vect) # compute mean
    # return demeaned vector
    return vect - m
end

```

```
random_vect = rand(1_000)
@show remove_mean(random_vect)
```

MethodError: MethodError: no method matching `- (::Vector{Float64}, ::Float64)`
For element-wise subtraction, use broadcasting with dot syntax: `array .- scalar`
The function ``-`` exists, but no method is defined for this combination of argument types.

Closest candidates are:

```
-(!Matched::Complex{Bool}, ::Real)
  @ Base complex.jl:329
-(!Matched::Missing, ::Number)
  @ Base missing.jl:123
-(!Matched::BigFloat, ::Union{Float16, Float32, Float64})
  @ Base mpfr.jl:565
...
```

MethodError: no method matching `- (::Vector{Float64}, ::Float64)`

For element-wise subtraction, use broadcasting with dot syntax: `array .- scalar`

The function ``-`` exists, but no method is defined for this combination of argument types.

Closest candidates are:

```
-(!Matched::Complex{Bool}, ::Real)

  @ Base complex.jl:329
-(!Matched::Missing, ::Number)

  @ Base missing.jl:123
-(!Matched::BigFloat, ::Union{Float16, Float32, Float64})

  @ Base mpfr.jl:565
...
```

Stacktrace:

```
[1] remove_mean(vect::Vector{Float64})

 @ Main ~/BEE 4750/bee4750-hw1-pdc76/jl_notebook_cell_df34fa98e69747e1a8f8a730347b8e2f_X30sZmls

[2] macro expansion

 @ show.jl:1232 [inlined]

[3] top-level scope

 @ ~/BEE 4750/bee4750-hw1-pdc76/jl_notebook_cell_df34fa98e69747e1a8f8a730347b8e2f_X30sZmls
```

What does this `MethodError` mean? Why does this code produce one? Fix the error and solve the problem.

Problem 1.4 Solution

This `MethodError` indicates that we are trying to do element-wise subtraction, but what we are subtracting is not an array. This occurs in the `remove_mean([])` function when we try to subtract `vect - m`. Julia doesn't know how to deal with subtraction of an integer from an array, so we have to use the operator `.-` in order to handle this. See corrected code below:

```
# function to remove mean from a vector
function remove_mean(vect)
    # function to compute the mean
    function compute_mean(vect)
        element_sum = 0 # initialize sum
        # compute mean and return
        for v in vect
            element_sum += v
        end
        return element_sum / length(vect)
    end

    m = compute_mean(vect) # compute mean
    # return demeaned vector
    return vect .- m
end
```

```
random_vect = rand(1_000)
@show remove_mean(random_vect)
```

```
remove_mean(random_vect) = [-0.3742068799795504, -0.22137438001670173, 0.13983318238138076, ...]
```

```
1000-element Vector{Float64}:
```

```
-0.3742068799795504
-0.22137438001670173
 0.13983318238138076
 0.17615354307930053
 0.1770106929061267
-0.4024537712628796
-0.00543377164007619
-0.45454765208074355
-0.19388551551759303
-0.2831164582780967
```

```
-0.1542603211615855
-0.2502250217142279
 0.22997610173145833
-0.002288954204928473
-0.014550916429255989
 0.12551199798981827
 0.3772353827400948
 0.05815824435508066
-0.08918753570927829
```

Problem 2 (18 points)

Cheap Plastic Products, Inc. is operating a plant that produces $100\text{m}^3/\text{day}$ of wastewater that is discharged into Pristine Brook. The wastewater contains $1\text{kg}/\text{m}^3$ of YUK, a toxic substance. The US Environmental Protection Agency has imposed an effluent standard on the plant prohibiting discharge of more than $20\text{kg}/\text{day}$ of YUK into Pristine Brook.

Cheap Plastic Products has analyzed two methods for reducing its discharges of YUK. Method 1 is land disposal, which costs $X_1^2/20$ dollars per day, where X_1 is the amount of wastewater disposed of on the land (m^3/day). With this method, 20% of the YUK applied to the land will eventually drain into the stream (*i.e.*, 80% of the YUK is removed by the soil).

Method 2 is a chemical treatment procedure which costs $\$1.50$ per m^3 of wastewater treated. The chemical treatment has an efficiency of $e = 1 - 0.005X_2$, where X_2 is the quantity of

wastewater (m^3/day) treated. For example, if $X_2 = 50\text{m}^3/\text{day}$, then $e = 1 - 0.005(50) = 0.75$, so that 75% of the YUK is removed.


Cheap Plastic Products is wondering how to allocate their wastewater between these three disposal and treatment methods (land disposal, chemical treatment, and direct disposal) to meet the effluent standard while keeping costs manageable.

The flow of wastewater through this treatment system is shown in Figure 1. Modify the edge labels (by editing the `edge_labels` dictionary in the code producing Figure 1) to show how the wastewater allocations result in the final YUK discharge into Pristine Brook. For the `edge_label` dictionary, the tuple (i, j) corresponds to the arrow going from node i to node j . The syntax for any entry is $(i, j) \Rightarrow \text{"label text"}$, and the label text can include mathematical notation if the string is prefaced with an L, as in `L"x_1"` will produce x_1 .

```
A = [0 1 1 1;
      0 0 0 1;
      0 0 0 1;
      0 0 0 0]

names = ["Plant", "Land Treatment", "Chem Treatment", "Pristine Brook"]
# modify this dictionary to add labels
edge_labels = Dict((1, 2) => L"x_1", (1,3) => L"x_2", (1, 4) => L"x_3", (2, 4) => L"0.2*x_1",
shaped=[:hexagon, :rect, :rect, :hexagon]
xpos = [0, -1.5, -0.25, 1]
ypos = [1, 0, 0, -1]

p = graphplot(A, names=names,edgelabel=edge_labels, markersize=0.15, markershapes=shapes, ma
display(p)
```



Problem 2 Graph Explanation

The graph shows that the initial plant output, X , is subdivided into 3 parts: X_1 , X_2 , X_3 . X_1 refers to the effluent that receives land treatment, X_2 is the chemically treated outflow, and X_3 is the effluent that runs directly into the stream. Land treatment removes 80% of the YUK so the effluent from the land to the stream is $0.2 X_1$. The chemical treatment removes $(1 - e) X_2$ YUK, where e is the efficiency and $e = (1 - 0.005 * X_2)$. This means that $(1 - e) X_2$ YUK runs into the stream from the chemical treatment plant.

Problem 2.1

Formulate a mathematical model for the treatment cost and the amount of YUK that will be discharged into Pristine Brook based on the wastewater allocations. This is best done with some equations and supporting text explaining the derivation. Make sure you include, as additional equations in the model, any needed constraints on relevant values. You can find some basics on writing mathematical equations using the LaTeX typesetting syntax [here](#), and a cheatsheet with LaTeX commands can be found on the course website's [Resources page](#).

Problem 2.1 Solution

These are the given and maximum values for the effluent, YUK concentration, and mass flow rate. While $C = 1 \text{ kg/day}$, I think it is important to include this value in my equations, as it the most complete equation and would make for more simple changes to have a singular global variable, C .

$$X = 100 \text{ m}^3/\text{day}$$

$$C = 1 \text{ kg/m}^3 \text{ of YUK}$$

$$\dot{m} = 100 \text{ kg/day}$$

$$\dot{m}_{\max} = 20 \text{ kg/day}$$

This is the given collective volumetric flow rate for all 3 effluent discharges, with X_3 being solved for as the difference between the total and the two reduction methods.

$$X = X_1 + X_2 + X_3 \quad \Rightarrow \quad X_3 = X - X_1 - X_2$$

$\dot{m}_1$ and $\dot{m}_2$ represent the mass flow rate of each treated effluent stream, and $\dot{m}_3$ is the untreated effluent.

$$\dot{m}_1 = 0.2X_1C$$

$$\dot{m}_2 = (1 - e)X_2C$$

$$= (1 - (1 - 0.005X_2))X_2C$$

$$\dot{m}_3 = X_3C$$

These equations of total mass flow rate and total mass flow rate effluent. The maximum total mass flow is the total effluent allowed in the stream, $\dot{m}_{\max} = 20$.

$$\dot{m}_{\text{tot}} = \dot{m}_1 + \dot{m}_2 + \dot{m}_3$$

$$\dot{m}_{\text{tot}} \leq \dot{m}_{\max}$$

This is the cost equation. The goal is to minimize this value while staying under the $\dot{m}_{\max} = 20$ maximum.

$$\text{Cost} = \frac{X_1^2}{20} + 1.50X_2$$

Problem 2.2

Implement your systems model as a Julia function which computes the resulting YUK discharge and cost for a particular treatment plan. You can return multiple values from a function with a [tuple](#), as in:

```
function multiple_return_values(x, y)
    return (x+y, x*y)
end
```

```
a, b = multiple_return_values(2, 5)
@show a;
@show b;
```

```
a = 7
b = 10
```

To evaluate the function over vectors of inputs, you can *broadcast* the function by adding a decimal `.` before the function arguments and accessing the resulting values by writing a *comprehension* to loop over the individual outputs in the vector:

```
x = [1, 2, 3, 4, 5]
y = [6, 7, 8, 9, 10]

output = multiple_return_values.(x, y)
a = [out[1] for out in output]
b = [out[2] for out in output]
@show a;
@show b;
```

```
a = [7, 9, 11, 13, 15]
b = [6, 14, 24, 36, 50]
```

Make sure you comment your code appropriately to make it clear what is going on and why.

Problem 2.2 Solution Code

```

function yuk_treatment_method(x1,x2)
    #Initial Givens
    c = 1
    x = 100
    mdot_max = 20

    #Calculate the effluent volume directly into stream
    x3 = x - x1 - x2

    #Solve for mass flow rate of each effluent and find sum of effluent into the stream
    mdot1 = x1 * c * 0.2
    mdot2 = (1-(1-0.005*x2)) * (x2 * c)
    mdot3 = x3 * c
    mdot_tot = mdot1 + mdot2 + mdot3

    #Calculate the cost of the chosen treatment plan
    cost = ((x1^2)/20) + (1.5*x2)

    #return the total YUK in the stream and the cost of the treatment
    return (mdot_tot,cost)
end

```

```

eff = 54.5
cost = 80.0

```

The function “yuk_treatment_method(x1,x2)” takes 2 parameters, x1 and x2, which are used to calculate the total effluent entering stream, as well as the proportion that receives land and chemical treatment. It also calculates the cost associated with each set of the chosen proportions entered as function arguments.

Problem 2.3

Use your function to experiment with 1,000 different combinations of wastewater discharge and treatment. You can do this with either a grid search or by sampling from a [Dirichlet distribution](#) (a $\text{Dirichlet}(1, n)$ distribution will generate uniformly-weighted n -dimensional vectors whose components add up to 1; see the [Distributions.jl documentation](#) for how to sample from probability distributions in Julia). Plot the results of these experiments. Do any satisfy the YUK effluent standard (plot this as well as a dashed red line). What was the cost of solutions satisfying the standard? What can you say about the tradeoff between treatment cost and YUK discharge? You don’t have to find an “optimal” solution to this problem, but what do you think would be needed to find a better solution?

```

#Define the number of samples and total volumetric flow rate
N = 1000
X = 100 #m3/day

#Take a Dirichlet sample and store it in a 3xN matrix
dir = Dirichlet(3,1)
sample = rand(dir,N)

#Define the vectors for X1 and X2
x1 = X .* vec(sample[1,:])
x2 = X .* vec(sample[2,:])

#Broadcast samples over the function using method shown above
output = yuk_treatment_method.(x1,x2)
eff = [out[1] for out in output]
cost = [out[2] for out in output]

#Define lists for viable and non-viable effluent treatment means
viable = []
nviabile = []

#Iterate through all of the samples check to see if each distribution of x1 and x2 is a viable
#combination by making sure the effluent is below 20 kg/day and assigning it to a list according
for i in eachindex(x1,x2,eff,cost)
    if eff[i] <= 20
        push!(viable,(x1[i],x2[i],eff[i],cost[i]))
    else
        push!(nviabile,(x1[i],x2[i],eff[i],cost[i]))
    end
end

#Define the effluent and cost for valid and invalid samples (X<20 kg/day) and separate for each
eff_viable = [out[3] for out in viable]
cost_viable = [out[4] for out in viable]
eff_nviabile = [out[3] for out in nviabile]
cost_nviabile = [out[4] for out in nviabile]


#Plot each data set, a vertical line showing the cutoff threshold, and min/max of cost and effluent
plt = scatter(eff_viable, cost_viable, color=:green, label = "Valid Samples", xlabel="YUK Effluent",
    ylabel="Total Cost per day (\$/day)", title="Cost vs YUK Effluent")

scatter!(eff_nviabile, cost_nviabile, color=:orange, label = "Invalid Samples", xlabel="YUK Effluent",
    ylabel="Total Cost per day (\$/day)", title="Cost vs YUK Effluent")

```

```
ylabel="Total Cost per day (\$/day)", title="Cost vs YUK Effluent")

vline!(plt, [20]; linestyle=:dash, color=:red, label="Maximum [YUK] in Effluent")
```



Problem 2.4

Find the strategies which minimize cost and YUK discharge (these will be different strategies) analytically and find the values of the objective metrics. Plot these values in the plot that you created for Problem 2.3. How do their values compare to the spread of values that you found in that problem? Would you select either of them (explain why or why not)?

Problem 2.3/2.4 Answers

References

List any external references consulted, including classmates.