

BEE 4750 Homework 2: Systems Modeling and Simulation

Name: Adrián Cardona Young

ID: alc354

Due Date

Thursday, 09/25/25, 9:00pm

Overview

Instructions

- Problem 1 asks you to draw a systems diagram and identify the type of a feedback.
- Problem 2 asks you to model contaminant concentrations in a river and use simulation to compare the concentrations to a regulatory standard.
- Problem 3 asks you to explore the implications of an ice-albedo feedback in the Earth's climate system by understanding the equilibria of the modeled system and their stabilities.

Load Environment

The following code loads the environment and makes sure all needed packages are installed. This should be at the start of most Julia scripts.

```
import Pkg
Pkg.activate(@__DIR__)
Pkg.instantiate()
```

Activating project at `c:\Users\adric\OneDrive - Cornell University\Documents\3. Env Sys A

```
Pkg.add("Measures")
```

```
Updating registry at `C:\Users\adric\.julia\registries\General.toml`
```

```
Resolving package versions...
```

```
No Changes to `C:\Users\adric\OneDrive - Cornell University\Documents\3. Env Sys Analysis\I
```

```
No Changes to `C:\Users\adric\OneDrive - Cornell University\Documents\3. Env Sys Analysis\I
```

```
using Plots
using LaTeXStrings
using CSV
using DataFrames
using Roots
using GraphRecipes
using Measures
```

Problems (Total: 30 Points)

Problem 1 (5 points)

Draw a systems diagram for the relationship between global mean temperature, atmospheric CO₂ concentrations, and ocean CO₂ concentrations. What are the signs of the interactions between these components and why? What does this suggest about the overall feedback between temperature and the ocean carbon cycle?

Tip

Think about Henry's law for CO₂.

```
A = [0 1 1;
      1 0 1;
      1 1 0]
```

```
names = [L"Global \ Mean \ Temperature", L"Atmospheric \ CO_2", L"Oceanic \ CO_2"]
```

```
# modify this dictionary to add labels
```

```
edge_labels = Dict{Tuple{Int,Int}}{String}((1, 2) => "+", (1,3) => "-", (2, 3) => "+")
```

```
edgecurves = Dict{Tuple{Int,Int}}{Float64}((1, 2) => -0.05, (1,3) => -0.05, (2, 3) => 0.05)
```


```
shapes=[:rect, :rect, :rect]
```

```
xpos = [1, 0, 2]
```

```
ypos = [1, 0, 0]
```

```
p = graphplot(A, names=names, edgelabel=edge_labels, markersize=0.07, markershapes=shapes, ma
```

```
size=(700, 400), fontsize = 10, curvature_scalar=edgecurves)
display(p)
```



Problem 2 (15 points)

A river which flows at 10 km/d is receiving discharges of wastewater contaminated with CRUD from two sources which are 15 km apart, as shown in the Figure below. CRUD decays exponentially in the river at a rate of 0.36 d^{-1} and is deposited by the atmosphere along the river at a rate of 54 kg/km/d.




Figure 1: river_diagram.png

Problem 2.1

Draw a systems diagram with the relevant control volume(s) denoted and any relevant in/out-flows between these boxes. How did you decide how many boxes were needed?




Figure 2: p2.2 diagram.png

Problem 2.2

Develop a model for the concentration of CRUD downriver by formulating and solving the appropriate differential equation(s) analytically.

Tip

Formulate your model in terms of distance downriver, rather than leaving it in terms of time from discharge.

Problem 2.3

Determine if the system is in compliance with a regulatory limit of 2.3 kg/(1000m³). You can do this analytically or computationally.

Volumetric Flow Balance

$$Q_3 = Q_0 + Q_1 = 250,000 + 40,000 = 290,000 \text{ m}^3/\text{d}$$

$$Q_f = Q_3 + Q_2 = 290,000 + 60,000 = 350,000 \text{ m}^3/\text{d}$$

Mass Flow Balance

First Mixing:

$$Q_3 C_3 = Q_0 C_0 + Q_1 C_1 = 250,000 (0.5/1000) + 40,000 (9/1000) = 485 \text{ kg}/(1000 \text{ m}^3)$$

$$C_3 = 485/290,000 * 1000 = 1.67 \text{ kg}/(1000 \text{ m}^3)$$

Between First and Second Mixing:

$$d(CV)/dt = Q_3 C_3 - k Q_3 C \quad d/v + d \cdot 54, \text{ where } d = \text{distance}, v = \text{velocity}$$

$$\text{Mass change between first and second mixing } \rightarrow d(CV)/dt = Q_3 C_3 - k Q_3 C \quad d/v + (54 - v)/t, \text{ where } d = \text{distance}, v = \text{velocity}$$

$$Q_3 C_3^* = (485 - 0.36 \cdot 485 \cdot 15/10 + (54 - 10)/1.5) \cdot 1.5 = 389 \text{ kg}/\text{d}$$

Where 1.5 = time in days, C_3^* = concentration flowing into the second mixing (not the concentration flowing out the first mixing)

Should it be:

$$C(t) = C_f e^{-k \cdot d/v} + \text{depo rate}$$

Second Mixing

$$Q_f C_f = Q_3 C_3^* + Q_2 C_2 = 389 + 60,000 \cdot 7/1000 = 809 \text{ kg}/\text{d}$$

Concentration Downstream of Second Mixing

$$Q \quad C(d) = C_f e^{-\{k/d/v\}}$$

Problem 3 (10 points)

In class, we discussed the ice-albedo feedback and its possible influence on melting the hypothesized “Snowball Earth”. In this problem, we’ll introduce a simple model of the Earth’s energy balance with includes this feedback and examine the stability of the climate.

This simple model of the energy balance (averaged over the entire planet) is:

$$\underbrace{C \frac{dT}{dt}}_{\text{change in heat}} = \underbrace{\frac{(1-\alpha)S}{4}}_{\text{incoming radiation}} - \underbrace{(A - BT)}_{\text{outgoing radiation}} + \underbrace{a \ln \left(\frac{[CO_2]}{[CO_2]_{PI}} \right)}_{\text{greenhouse effect}}, \quad (1)$$

where:

- T is the Earth’s global mean temperature (in °C);
- C is the heat capacity of the atmosphere and shallow ocean, taken to be 51 J/m²/°C;
- α is the planetary albedo, or the fraction of incoming radiation reflected by the Earth, which has a present-day value of approximately 0.3;
- S is the solar constant, or the amount of solar radiation received by the Earth averaged over area, which has a present-day value of 1368 W/m² but during the Neoproterozoic Era had a value of 1272 W/m² (this is divided by four because the Earth is a sphere but S is the radiation captured by a disc with radius equal to that of the Earth);
- A and B are coefficients from the linearization of outgoing radiation physics, and have corresponding values $B = -1.3$ W/m²/°C (estimated from a number of lines of evidence about the sensitivity of outgoing radiation to temperature; this is negative due to a sign convention about the direction of incoming vs. outgoing radiation) and $A = 221.2$ W/m² (estimated by assuming the pre-industrial temperature of 14°C was stable without anthropogenic greenhouse gas emissions).

We will ignore the greenhouse effect term as we are considering the Earth system well before humans were around.

Problem 3.1

Discretize the climate model (Equation 1) using forward Euler integration and a time step of $\Delta t = 1$ yr.

$$T_{t+\Delta t} = T_t + \frac{\Delta t}{C} \left(\frac{(1-\alpha)S}{4} - A + BT \right)$$

Problem 3.2

Rather than assuming a constant value for the albedo α , we will represent the ice-albedo feedback by letting α depend on T :

$$\alpha(T) = \begin{cases} \alpha_i & \text{if } T \leq -10^\circ\text{C} \\ \alpha_i + (\alpha_0 - \alpha_i)((T + 10)/20) & \text{if } -10^\circ\text{C} \leq T \leq 10^\circ\text{C} \\ \alpha_0 & \text{if } T \geq 10^\circ\text{C}. \end{cases}$$

Let $\alpha_i = 0.5$ and $\alpha_0 = 0.3$. Simulate the simple climate model using the Neoproterozoic value of S with temperature-varying albedo for initial values of T spanning $-60^\circ\text{C} \leq T_0 \leq 30^\circ\text{C}$ over a period of 200 years. Plot the temperature trajectories. How many equilibria are there and what are their stabilities?

```
C = 51 #J/m2/deg C
ai = 0.5
a0 = 0.3
S = 1272 #W/m2, Neoproterozoic solar radiation
A = 221.2 #W/m2
B = -1.3 #W/m2
step = 1 #year

init_temps = collect(range(-60, 30, 10)) #create range of initial temperatures
time = collect(range(0, 200, 20)) #time array (year 0 through 200)

function SimPlot(temps, times, ai, a0, S, A, B, step)
    res = zeros(10,201)
    res[:, 1] = temps
    for i in 1:length(res[:,1])
        for y in 1:size(res, 2)-1
            t_year = res[i, y]
            if t_year < -10
                alpha = ai
            elseif -10 < t_year <= 10
                alpha = ai + (a0 - ai)*((t_year + 10)/(20))
            else t_year > 10
                alpha = a0
            end
            res[i, y+1] = t_year + step * (1/C) * (((1-alpha)*S)/(4) - A + B*t_year)
        end
    end
end
```

```

    return res
end

```

SimPlot (generic function with 1 method)

```


res = SimPlot(init_temps, time, ai, a0, S, A, B, step);

```

```

p1 = plot(time, res[1,:], xlabel=L"Time (years)", ylabel=L"Temperatures (\degree C)", legend=
title=L"Temperature \ Profiles \ for \ varying \ T_0")
for i in 1:9
    plot!(time, res[i+1,:])
end
display(p1)

```




Problem 3.3

One might hypothesize that one cause for the transition from Snowball Earth (the stable frozen equilibrium from Problem 3.2) was an increasing amount of incoming solar radiation

(as expressed by an increase in S). Examine this hypothesis using our simple model. Is this factor enough to cause the planet to warm to the pre-industrial temperature of 14°C ?

```
res = SimPlot(init_temps, time, ai, a0, 1368, A, B, step);

p2 = plot(time, res[1,:], xlabel=L"Time (years)", ylabel=L"Temperatures (\degree C)", legend=
title=L"Temperature \ Profiles \ for \ varying \ T_0")
for i in 1:9
    plot!(time, res[i+1,:])
end
display(p2)
```



The above plot shows the difference in the trends when changing the incoming solar radiation to $1368 \text{ W}/\text{m}^2$ as opposed to the $1272 \text{ W}/\text{m}^2$ we used in Problem 3.2. This suggests that an increase in solar radiation could contribute to the warming to preindustrial levels. With temperatures somewhere between 0 and 10°C as starting temperature, it would tend towards 14°C and stabilize there. This, in conjunction with other changes, could result in the pre-industrial temperatures we saw. It would depend, however, on the initial temperature, as temperatures below a certain value all tend to almost -40°C . This means it might have taken

an event that raised the Earth's temperature in combination with increased solar radiation to make it happen.

References

List any external references consulted, including classmates.