

BEE 4750 Homework 2: Systems Modeling and Simulation

Name:

ID:

Due Date

Thursday, 09/25/25, 9:00pm

Overview

Instructions

- Problem 1 asks you to draw a systems diagram and identify the type of a feedback.
- Problem 2 asks you to model contaminant concentrations in a river and use simulation to compare the concentrations to a regulatory standard.
- Problem 3 asks you to explore the implications of an ice-albedo feedback in the Earth's climate system by understanding the equilibria of the modeled system and their stabilities.

Load Environment

The following code loads the environment and makes sure all needed packages are installed. This should be at the start of most Julia scripts.

```
import Pkg
Pkg.activate(@__DIR__)
Pkg.instantiate()
```

```
Activating project at `c:\Users\holly\4750 HW\hw2-holly`
Precompiling project...
 22029.3 ms    Plots → UnitfulExt
 46358.6 ms    DataFrames
  3286.0 ms    Latexify → DataFramesExt
3 dependencies successfully precompiled in 103 seconds. 252 already precompiled.
```

```
Pkg.add("GraphRecipes")
using Plots
using LaTeXStrings
using CSV
using DataFrames
using Roots
using Random
using Plots
using GraphRecipes
using LaTeXStrings
using Distributions
```

```
Resolving package versions...
No Changes to `C:\Users\holly\4750 HW\hw2-holly\Project.toml`
No Changes to `C:\Users\holly\4750 HW\hw2-holly\Manifest.toml`

ArgumentError: Package Distributions not found in current path.
- Run `import Pkg; Pkg.add("Distributions")` to install the Distributions package.
ArgumentError: Package Distributions not found in current path.

- Run `import Pkg; Pkg.add("Distributions")` to install the Distributions package.
```

Stacktrace:

```
[1] macro expansion
      @ .\loading.jl:2296 [inlined]

[2] macro expansion
      @ .\lock.jl:273 [inlined]

[3] __require(into::Module, mod::Symbol)
```

```

@ Base .\loading.jl:2271

[4] #invoke_in_world#3

@ .\essentials.jl:1089 [inlined]

[5] invoke_in_world

@ .\essentials.jl:1086 [inlined]

[6] require(into::Module, mod::Symbol)

@ Base .\loading.jl:2260

[7] eval

@ .\boot.jl:430 [inlined]

[8] include_string(mapexpr::typeof(REPL.softscope), mod::Module, code::String, filename::String)

@ Base .\loading.jl:2734

[9] #invokelatest#2

@ .\essentials.jl:1055 [inlined]

[10] invokelatest

@ .\essentials.jl:1052 [inlined]

[11] (::VSCoServer.var"#217#218"{VSCoServer.NotebookRunCellArguments, String})()

@ VSCoServer c:\Users\holly\.vscode\extensions\julialang.language-julia-1.149.2\scripts\

[12] withpath(f::VSCoServer.var"#217#218"{VSCoServer.NotebookRunCellArguments, String}, path::String)

@ VSCoServer c:\Users\holly\.vscode\extensions\julialang.language-julia-1.149.2\scripts\

[13] notebook_runcell_request(conn::VSCoServer.JSONRPC.JSONRPCEndpoint{Base.PipeEndpoint, Base.Pipe})

@ VSCoServer c:\Users\holly\.vscode\extensions\julialang.language-julia-1.149.2\scripts\

```

```

[14] dispatch_msg(x::VSCodeServer.JSONRPC.JSONRPCEndpoint{Base.PipeEndpoint, Base.PipeEndpoint})
      @ VSCodeServer.JSONRPC c:\Users\holly\.vscode\extensions\julialang.language-julia-1.149.2\scripts\notebook\notebook.jl
[15] serve_notebook(pipename::String, debugger_pipename::String, outputchannel_logger::Base.Logging.Logger)
      @ VSCodeServer c:\Users\holly\.vscode\extensions\julialang.language-julia-1.149.2\scripts\notebook\notebook.jl
[16] top-level scope
      @ c:\Users\holly\.vscode\extensions\julialang.language-julia-1.149.2\scripts\notebook\notebook.jl

```

Problems (Total: 30 Points)

Problem 1 (5 points)

Draw a systems diagram for the relationship between global mean temperature, atmospheric CO2 concentrations, and ocean CO2 concentrations. What are the signs of the interactions between these components and why? What does this suggest about the overall feedback between temperature and the ocean carbon cycle?


```

A = [0 0 1 ;
      1 0 1 ;
      1 0 0 ]

names = ["Global Mean temp", "Atm CO2", "Ocean CO2"]
# modify this dictionary to add labels
edge_labels = Dict{(:, :)}{String}((2, 1) => "+", (1, 3) => "-", (2, 3) => "+", (3, 1) => "-")
shapes = [:rect, :rect, :rect]
xpos = [0, -0.9, 0.9]
ypos = [1, 0, 0]

p = graphplot(A, names=names, edgelabel=edge_labels, markersize=0.15, markershapes=shapes, margin=10)
display(p)

```



From Henry's law, we know that as the partial pressure of CO₂ increases above a body of water, the solubility of CO₂ in the water increases. This led us to choose a "+" for the relationship between Atm CO₂ and Ocean CO₂: As atm CO₂ increases, so will oceanic CO₂. On the other hand, as surface temperature increases, the ability of CO₂ to dissolve in water decreases. So as global mean temperature rises (along with ocean surface temperature), less CO₂ will dissolve, resulting in a "-" relationship. Additionally for this inverse, or dampening, relationship, oceanic carbon is a carbon sink, with more dissolved carbon resulting in slower warming. This solidifies our selection of using "-". We also know from the greenhouse effect that as more CO₂ resides in the atmosphere, more radiation will be trapped in the atmosphere contributing to a rise in temperature. Given this, we chose a "+" relationship.

Tip

Think about Henry's law for CO₂.

Problem 2 (15 points)

A river which flows at 10 km/d is receiving discharges of wastewater contaminated with CRUD from two sources which are 15 km apart, as shown in the Figure below. CRUD decays expo-

nentially in the river at a rate of 0.36 d^{-1} and is deposited by the atmosphere along the river at a rate of 54 kg/km/d .

Schematic of the river system in Problem 2

Problem 2.1

Draw a systems diagram with the relevant control volume(s) denoted and any relevant in/out-flows between these boxes. How did you decide how many boxes were needed?

Problem 2.2

Develop a model for the concentration of CRUD downriver by formulating and solving the appropriate differential equation(s) analytically.

Tip

Formulate your model in terms of distance downriver, rather than leaving it in terms of time from discharge.

Problem 2.3

Determine if the system is in compliance with a regulatory limit of $2.3 \text{ kg/(1000m}^3\text{)}$. You can do this analytically or computationally.

Problem 3 (10 points)

In class, we discussed the ice-albedo feedback and its possible influence on melting the hypothesized “[Snowball Earth](#)”. In this problem, we’ll introduce a simple model of the Earth’s energy balance with includes this feedback and examine the stability of the climate.

This simple model of the energy balance (averaged over the entire planet) is:

$$\underbrace{C \frac{dT}{dt}}_{\text{change in heat}} = \underbrace{\frac{(1-\alpha)S}{4}}_{\text{incoming radiation}} - \underbrace{(A - BT)}_{\text{outgoing radiation}} + \underbrace{a \ln \left(\frac{[CO_2]}{[CO_2]_{PI}} \right)}_{\text{greenhouse effect}}, \quad (1)$$

where:

- T is the Earth’s global mean temperature (in $^{\circ}\text{C}$);
- C is the heat capacity of the atmosphere and shallow ocean, taken to be $51 \text{ J/m}^2/^{\circ}\text{C}$;
- α is the planetary albedo, or the fraction of incoming radiation reflected by the Earth, which has a present-day value of approximately 0.3;

- S is the solar constant, or the amount of solar radiation received by the Earth averaged over area, which has a present-day value of 1368 W/m^2 but during the Neoproterozoic Era had a value of 1272 W/m^2 (this is divided by four because the Earth is a sphere but S is the radiation captured by a disc with radius equal to that of the Earth);
- A and B are coefficients from the linearization of outgoing radiation physics, and have corresponding values $B = -1.3 \text{ W/m}^2/\text{°C}$ (estimated from a number of lines of evidence about the sensitivity of outgoing radiation to temperature; this is negative due to a sign convention about the direction of incoming vs. outgoing radiation) and $A = 221.2 \text{ W/m}^2$ (estimated by assuming the pre-industrial temperature of 14°C was stable without anthropogenic greenhouse gas emissions).

We will ignore the greenhouse effect term as we are considering the Earth system well before humans were around.

Problem 3.1

Discretize the climate model (Equation 1) using forward Euler integration and a time step of $\Delta t = 1 \text{ yr}$.

Problem 3.2

Rather than assuming a constant value for the albedo α , we will represent the ice-albedo feedback by letting α depend on T :

$$\alpha(T) = \begin{cases} \alpha_i & \text{if } T \leq -10^\circ\text{C} \\ \alpha_i + (\alpha_0 - \alpha_i)((T + 10)/20) & \text{if } -10^\circ\text{C} \leq T \leq 10^\circ\text{C} \\ \alpha_0 & \text{if } T \geq 10^\circ\text{C}. \end{cases}$$

Let $\alpha_i = 0.5$ and $\alpha_0 = 0.3$. Simulate the simple climate model using the Neoproterozoic value of S with temperature-varying albedo for initial values of T spanning $-60^\circ\text{C} \leq T_0 \leq 30^\circ\text{C}$ over a period of 200 years. Plot the temperature trajectories. How many equilibria are there and what are their stabilities?

Problem 3.3

One might hypothesize that one cause for the transition from Snowball Earth (the stable frozen equilibrium from Problem 3.2) was an increasing amount of incoming solar radiation (as expressed by an increase in S). Examine this hypothesis using our simple model. Is this factor enough to cause the planet to warm to the pre-industrial temperature of 14°C ?

References


List any external references consulted, including classmates.

```
# 3.2
# initial conditions/defining variables

A = 221.2
B = -1.3
C = 51
S = 1272

# albedo
ai = 0.5
a0 = 0.3

p = plot(xlabel="Step", ylabel="Temperature (T)", lw=2, legend = :outerbottomright)
# loop through T-values
for T0 in -60:1:30
    Ti = T0
    T = []
    I = []
    # 200 years of 1-year timesteps
    for i = 1:200
        push!(I, i)
        push!(T, Ti)
        # albedo based on temperature
        if Ti <= -10
            a = ai
        elseif Ti >= 10
            a = a0
        else
            a = ai + ((a0-ai)*((Ti+10)/20))
        end
        # new temperature
        T_new = Ti + 1/C * (((1-a)*S/4) - A + B*Ti)
        Ti = T_new
    end
    plot!(I, T, label="T initial=$T0")
end
display(p)
```

3.3 After testing with $S=1360$, another equilibrium point arises at around 14-15 degrees, C. This shows that an increase in solar radiation does lead to an increase in steady state temperature, and this may be attributed to Earth's proximity to the sun, closer leading to stronger radiation.