

BEE 4750 Homework 3: Dissolved Oxygen and Monte Carlo

Name: Claire Jacobson

ID: cpj35

Due Date

Thursday, 10/16/25, 9:00pm

Overview

Instructions

- Problem 1 asks you to implement a model for dissolved oxygen in a river with multiple waste releases and use this to develop a strategy to ensure regulatory compliance.

Load Environment

The following code loads the environment and makes sure all needed packages are installed. This should be at the start of most Julia scripts.

```
import Pkg
Pkg.activate(@__DIR__)
Pkg.instantiate()
```

Activating project at `~/hw3-claire`

```
using Random
using Plots
using LaTeXStrings
using Distributions
```

Problems (Total: 30 Points)

Problem 1 (30 points)

A river which flows at 6 km/d is receiving waste discharges from two sources which are 15 km apart. The oxygen reaeration rate is 0.55 day⁻¹, and the decay rates of CBOD and NBOD are 0.35 and 0.25 day⁻¹, respectively. The river's saturated dissolved oxygen concentration is 10 mg/L.

If the characteristics of the river inflow and waste discharges are given in Table 1, write a Julia model to compute the dissolved oxygen concentration from the first wastewater discharge to an arbitrary distance d km downstream. Use your model to compute the minimum dissolved oxygen concentration up to 50 km downstream and how far downriver this maximum occurs.

Parameter	River Inflow	Waste Stream 1	Waste Stream 2
Inflow	100,000 m ³ /d	10,000 m ³ /d	15,000 m ³ /d
DO Concentration	7.5 mg/L	5 mg/L	5 mg/L
CBOD	5 mg/L	50 mg/L	45 mg/L
NBOD	5 mg/L	35 mg/L	35 mg/L

Table 1: River inflow and waste stream characteristics for Problem 1.

Problem 1.1

Implement the Streeter-Phelps (analytic) solution for the dissolved oxygen concentration. Plot the dissolved oxygen concentration from the first waste stream to 50 km downriver. What is the minimum value in mg/L?

```
x2 = 15 # km
xf = 50 # km
dx = 0.1 # km
u = 6 # km/d

C_sat = 10 # mg/L

kr = 0.55 # day-1
kc = 0.35 # day-1
kn = 0.25 # day-1

Qr = 100000 # m3/day
Cdo_r = 7.5 # mg/L
```

```

Cc_r = 5 # mg/L
Cn_r = 5 # mg/L

Q1 = 10000 # m^3/day
Cdo_1 = 5 # mg/L
Cc_1 = 50 # mg/L
Cn_1 = 35 # mg/L

Q2 = 15000 # m^3/day
Cdo_2 = 5 # mg/L
Cc_2 = 45 # mg/L
Cn_2 = 35 # mg/L

C_new(q1, c1, q2, c2) = (q1 * c1 + q2 * c2) / (q1 + q2)

function mix(DO_0, CBOD0, NBOD0, t, k1, k2, k3, sat)
    DO_new = DO_0 * exp(-k1 * t)
    CBOD_new = CBOD0 * (k2 / (k1 - k2)) * (exp(-k2 * t) - exp(-k1 * t))
    NBOD_new = NBOD0 * (k3 / (k1 - k3)) * (exp(-k3 * t) - exp(-k1 * t))
    D = DO_new + CBOD_new + NBOD_new
    return sat - D
end

Q_first = Qr + Q1
DO_mix1 = C_new(Qr, Cdo_r, Q1, Cdo_1)
CBOD_mix1 = C_new(Qr, Cc_r, Q1, Cc_1)
NBOD_mix1 = C_new(Qr, Cn_r, Q1, Cn_1)
DO_first = C_sat - DO_mix1

decay(C_0, k, t) = C_0 * exp(-k * t)

xs = 0.0:dx:xf

for (i, x) in enumerate(xs)
    if x <= x2
        t = x / u
        DOs[i] = mix(DO_first, CBOD_mix1, NBOD_mix1, t, kr, kc, kn, C_sat)
    else
        t_x2 = x2 / u
        DO_x2 = mix(DO_first, CBOD_mix1, NBOD_mix1, t_x2, kr, kc, kn, C_sat)
        CBOD_x2 = decay(CBOD_mix1, kc, t_x2)
        NBOD_x2 = decay(NBOD_mix1, kn, t_x2)
    end
end

```

```

        DO_mix2 = C_new(Q_first, DO_x2, Q2, Cdo_2)
        CBOD_mix2 = C_new(Q_first, CBOD_x2, Q2, Cc_2)
        NBOD_mix2 = C_new(Q_first, NBOD_x2, Q2, Cn_2)
        DO_second = C_sat - DO_mix2
        t_after2 = (x - x2) / u
        DOs[i] = mix(DO_second, CBOD_mix2, NBOD_mix2, t_after2, kr, kc, kn, C_sat)
    end
end

min_conc=minimum(DOs)

p = plot(xs, DOs,
        xlim = (0, 50),
        ylim = (0,10),
        legend = false,
        xlabel = "Distance [km]",
        ylabel = "DO Conc [mg/L]",
        title = "DO Concentration Downstream of Waste Outlets"
)

vline!(p, [15], color=:red, linestyle=:dash)

display(p)
println("Minimum dissolved oxygen concentration = ", min_conc)

```

```

LoadError: UndefVarError: `DOs` not defined in `Main`
Suggestion: check for spelling errors or missing imports.
UndefVarError: `DOs` not defined in `Main`
Suggestion: check for spelling errors or missing imports.
Stacktrace:
 [1] top-level scope
      @ ./In[33]:50

```

Problem 1.2

Implement a numerically-integrated (discretized) solution for the dissolved oxygen concentration. Using a resolution of 0.5km, conduct a simulation and plot the results on the same axis as the plot from Problem 1.1. How has the minimum value changed? What do you attribute this difference to (be specific about the source of the difference(s) in terms of the simulation dynamics, not just that one is a numerical approximation).

The new minimum is ~0.9 mg/L higher than in the analytic solution. This can likely be attributed to some truncation error from the larger time step as compared to the analytical solution and the fact that the values are only incrementally evaluated. Also, the numerical integration method assumes constant values and only has values compared to those 0.5 km before, ignoring what goes on in between those time steps meaning a minimum will be mostly ignored if it lies between two time steps. This means a change could be ignored even further as the effects of evaluating based on that point will also be ignored.

```
dx_int = 0.5 # km
xs_int = 0.0:dx_int:xf
dt_int = dx_int / u

DO_int=zeros(length(xs_int))
CBOD_int = zeros(length(xs_int))
NBOD_int = zeros(length(xs_int))

DO_int[1] = DO_mix1
D=C_sat-DO_mix1
CBOD_int[1] = CBOD_mix1
NBOD_int[1] = NBOD_mix1

for i in 2:length(xs_int)
    CBOD_int[i] = CBOD_int[i-1] * exp(-kc*dt_int)
    NBOD_int[i] = NBOD_int[i-1] * exp(-kn*dt_int)

    D = D * exp(-kr*dt_int) + (CBOD_int[i-1] - CBOD_int[i])+(NBOD_int[i-1]-NBOD_int[i])

    DO_int[i] = C_sat - D

    if xs_int[i-1] < x2 < xs_int[i]
        DO_int[i] = C_new(Q_first, DO_int[i], Q2, Cdo_2)
        CBOD_int[i] = C_new(Q_first, CBOD_int[i], Q2, Cc_2)
        NBOD_int[i] = C_new(Q_first, NBOD_int[i], Q2, Cn_2)
        DO= C_sat - DO_int[i]
        Q_first += Q2
    end
end

min_conc_new=minimum(DO_int)


p2 = plot(xs_int, DO_int,
    xlim = (0, 50),
    ylim = (0,10),
```

```

    legend = false,
    xlabel = "Distance [km]",
    ylabel = "DO Conc [mg/L]",
    title = "DO Concentration Downstream: Numerical Integration"
)

display(p2)
println("Minimum dissolved oxygen concentration = ", min_conc_new)

```



Minimum dissolved oxygen concentration = 4.612126754365276

Problem 1.3

Using the analytic model, what is the minimum level of treatment (% removal of organic waste; assume this is equivalent to the same level of reduction of the CBOD and NBOD) for waste stream 1 that will ensure that the dissolved oxygen concentration is in compliance with the 4 mg/L standard along with a 5% margin of safety, assuming that waste stream 2 remains untreated? How about if only waste stream 2 is treated?

```

dx=0.1
x2=15
xf=50
u=6

function find_treat(tr1, tr2)
    Cc_1_t = Cc_1*(1-tr1) # mg/L
    Cn_1_t = Cn_1*(1-tr1) # mg/L

    Cc_2_t = Cc_2*(1-tr2)
    Cn_2_t = Cn_2*(1-tr2)

    Q_first = Qr + Q1
    DO_mix1 = C_new(Qr, Cdo_r, Q1, Cdo_1)
    CBOD_mix1_t = C_new(Qr, Cc_r, Q1, Cc_1_t)
    NBOD_mix1_t = C_new(Qr, Cn_r, Q1, Cn_1_t)
    DO_first = C_sat - DO_mix1

    decay(C_0, k, t) = C_0 * exp(-k * t)

    xt = 0.0:dx:xf
    DOs_t = zeros(length(xt))

    for (i, x) in enumerate(xt)
        if x <= x2
            t = x / u
            DOs_t[i] = mix(DO_first, CBOD_mix1_t, NBOD_mix1_t, t, kr, kc, kn, C_sat)
        else
            t_x2 = x2 / u
            DO_x2_t = mix(DO_first, CBOD_mix1_t, NBOD_mix1_t, t_x2, kr, kc, kn, C_sat)
            CBOD_x2_t = decay(CBOD_mix1_t, kc, t_x2)
            NBOD_x2_t = decay(NBOD_mix1_t, kn, t_x2)
            DO_mix2_t = C_new(Q_first, DO_x2_t, Q2, Cdo_2)
            CBOD_mix2_t = C_new(Q_first, CBOD_x2_t, Q2, Cc_2_t)
            NBOD_mix2_t = C_new(Q_first, NBOD_x2_t, Q2, Cn_2_t)
            DO_second_t = C_sat - DO_mix2_t
            t_after2_new = (x - x2) / u
            DOs_t[i] = mix(DO_second_t, CBOD_mix2_t, NBOD_mix2_t, t_after2_new, kr, kc, kn, C_sat)
        end
    end
end

# Compute minimum concentration (lowest DO)

```

```

    min_conc = minimum(DOs_t)
    return xt, DOs_t, min_conc
end

```

find_treat (generic function with 1 method)

```

C_std = 4*(1.05)

# Case 1: treat stream 1 only
for r1 in 0:0.01:1
    _, _, min_conc = find_treat(r1, 0)
    if min_conc >= C_std
        println("Stream 1 treatment required = ", round(r1*100, digits=1), "%")
        break
    end
end

# Treating stream 2 only
for r2 in 0:0.01:1
    _, _, min_conc = find_treat(0, r2)
    if min_conc >= C_std
        println("Stream 2 treatment required = ", round(r2*100, digits=1), "%")
        break
    end
end

```

Stream 1 treatment required = 28.0%

Stream 2 treatment required = 22.0%

Problem 1.4

Suppose you are responsible for designing a waste treatment plan for discharges into the river, with a regulatory mandate to keep the dissolved oxygen concentration above 4 mg/L. Discuss whether you'd opt to treat waste stream 2 alone or both waste streams equally. What other information might you need to make a conclusion, if any?

I would prefer to treat stream both equally. Although stream 2 requires less percentage of treatment, treating both streams would limit the effects throughout the entire stream. When only the second effluent is treated it is more of a reactive approach than proactive, allowing the 15 km stretch of the river to receive all of the waste from the first effluent. In order to make a more accurate conclusion I would like to know the difference in cost of the two strategies

(will that much money really be saved by treating only outlet 2). I would also like to know how much the BOD concentrations fluctuate. Having both outlets treated allows to prepare for increases in input from either one.

Problem 1.5

Suppose that it is known that the DO concentrations at the river inflow can vary according to a LogNormal(2.0, 0.15) distribution. Conduct a Monte Carlo simulation of the DO concentration in the river. If you treat only waste stream 1 (based on your analysis from Problem 1.3), what is the expected probability and 95% confidence interval that the river fails to comply with the regulatory standard of 4 mg/L? How did you decide that your Monte Carlo sample size was sufficiently large?

References

List any external references consulted, including classmates.