

BEE 4750 Lab 2: Monte Carlo

Name: Eliana Olson

ID: ero23

Due Date

Thursday, 10/02/25, 9:00pm

Setup

The following code should go at the top of most Julia scripts; it will load the local package environment and install any needed packages. You will see this often and shouldn't need to touch it.

```
import Pkg
Pkg.activate(".")
Pkg.instantiate()
```

Activating project at `~/lab-2-ergocat`

```
using Random # random number generation
using Distributions # probability distributions and interface
using Statistics # basic statistical functions, including mean
using Plots # plotting
```

Overview

In this lab, we will conduct a Monte Carlo experiment and explore how sensitive Monte Carlo estimates are to the underlying probability distribution(s).

You should always start any computing with random numbers by setting a “seed,” which controls the sequence of numbers which are generated (since these are not *really* random, just “pseudorandom”). In Julia, we do this with the `Random.seed!()` function.

```
Random.seed!(1)
```

`TaskLocalRNG()`

It doesn't matter what seed you set, though different seeds might result in slightly different values. But setting a seed means every time your notebook is run, the answer will be the same.

Seeds and Reproducing Solutions

If you don't re-run your code in the same order or if you re-run the same cell repeatedly, you will not get the same solution. If you're working on a specific problem, you might want to re-use `Random.seed()` near any block of code you want to re-evaluate repeatedly.

Probability Distributions and Julia

Julia provides a common interface for probability distributions with the [Distributions.jl package](#). The basic workflow for sampling from a distribution is:

1. Set up the distribution. The specific syntax depends on the distribution and what parameters are required, but the general call is the similar. For a normal distribution or a uniform distribution, the syntax is

```
# you don't have to name this "normal_distribution"
#  is the mean and  is the standard deviation
normal_distribution = Normal( , )
# a is the upper bound and b is the lower bound; these can be set to +Inf or -Inf for an
uniform_distribution = Uniform(a, b)
```

There are lots of both [univariate](#) and [multivariate](#) distributions, as well as the ability to create your own, but we won't do anything too exotic here.

2. Draw samples. This uses the `rand()` command (which, when used without a distribution, just samples uniformly from the interval $[0, 1]$.) For example, to sample from our normal distribution above:

```
# draw n samples
rand(normal_distribution, n)
```

Putting this together, let's say that we wanted to simulate 100 six-sided dice rolls. We could use a [Discrete Uniform distribution](#).

```
dice_dist = DiscreteUniform(1, 6) # can generate any integer between 1 and 6
dice_rolls = rand(dice_dist, 100) # simulate rolls
```


100-element Vector{Int64}:

```
1
3
5
4
6
2
5
5
5
2

3
6
5
5
6
3
6
6
6
```

And then we can plot a histogram of these rolls:

```
histogram(dice_rolls, legend=:false, bins=6)
ylabel!("Count")
xlabel!("Dice Value")
```



Instructions

Remember to:

- Evaluate all of your code cells, in order (using a `Run All` command). This will make sure all output is visible and that the code cells were evaluated in the correct order.
- Tag each of the problems when you submit to Gradescope; a 10% penalty will be deducted if this is not done.

Problem

A common engineering problem is to quantify flood risk, which is typically computed by propagating a flood hazard distribution through a *depth-damage function* relating flood depths to economic damages. A reasonable depth-damage function for a house without a basement is a bounded logistic function,

$$d(h) = \mathbb{1}_{h>0} \frac{L}{1 + \exp(-k(h - h_0))},$$

where d is the damage as a percent of total structure value, h is the water depth (relative to the house's ground floor elevation) in m, $\mathbb{1}_{x>0}$ is the indicator function, L is the maximum loss in USD, k is the slope of the depth-damage relationship, and h_0 is the inflection point. We'll assume $L = \$200,000$, $k = 0.8$, and $h_0 = 3$.

For this problem, suppose that we have two different probability distributions characterizing annual maxima flood depths:

1. $h \sim \text{LogNormal}(1.2, 0.3)$;
2. $h \sim \text{GeneralizedExtremeValue}(3, 0.9, -0.15)$.

Plot histograms of samples from these distributions and conduct a Monte Carlo experiment for annual maximum flood damages using each. Answer the following questions:

- What are the Monte Carlo estimates of the expected value and the 99% quantile for the annual maximum damage the structure would suffer for each of these flood hazard distributions?
- How did you decide on your sample size?
- Why do you think the estimates differed or did not differ (you can also plot the depth-damage function to compare with the samples)?
- How might you decide (based on getting additional information, if needed) which distribution is “better”?

```
L = 200000 #maximum loss in USD
k = 0.8 #slope of the dept-damage relationship
h0 = 3 #inflection point
h1_dis = LogNormal(1.2, 0.3)
h2_dis = GeneralizedExtremeValue(3, 0.9, -0.15)

#Simulate the annual maxima flood depth arrays
h1 = rand(h1_dis, 500000)
h2 = rand(h2_dis, 500000)

damage1 = zeros(500000)
damage2 = zeros(500000)

for i in 1:length(h1)
    if h1[i] > 0
        damage1[i] = L / (1 + exp(-k * (h1[i]-h0)))
    elseif h1[i] < 0
        damage1[i] = 0
    end
end
end
```

```

for i in 1:length(h2)
    if h2[i] > 0
        damage2[i] = L / (1 + exp(-k * (h2[i]-h0)))
    else
        damage2[i] = 0
    end
end

end


histogram(damage1, legend=:false, bins=100)
ylabel!("frequency for h1")
display(xlabel!("Flood Damage"))


histogram(damage2, legend=:false, bins=100)
ylabel!("frequency for h2")
display(xlabel!("Flood Damage"))

print("The estimate of the expected value for the LogNormal distribution mean is:", mean(damage1))
print("The estimate of the expected value for the GEV distribution mean is:", mean(damage2))

print("The estimate of the 99% quantile for the LogNormal distribution mean is:", quantile(damage1, 0.99))
print("The estimate of the 99% quantile for the GEV distribution mean is:", quantile(damage2, 0.99))

```





The estimate of the expected value for the LogNormal distribution mean is:115163.12828427345
The estimate of the expected value for the GEV distribution mean is:113550.08803271444

The estimate of the 99% quantile for the LogNormal distribution mean is:189877.4274980219
The estimate of the 99% quantile for the GEV distribution mean is:183229.56025684162

1.) What are the Monte Carlo estimates of the expected value and the 99% quantile for the annual maximum damage the structure would suffer for each of these flood hazard distributions?

The Monte Carlo estimate for the expected value is **115,163.128** for the LogNormal distribution and **113,550.088** for the GEV distribution.

The 99% quantile is **189,877.427** for the LogNormal distribution and **183,229.560** for the GEV distribution.

2.) How did you decide on your sample size?

I started at 50 and multiplied by 10 until the curve looked relatively smooth without significant outliers/noise. This ended up being 500,000.

3.) Why do you think the estimates differed or did not differ (you can also plot the depth-damage function to compare with the samples)?

The Lognormal results had a mean that was lower and a longer tail, meaning a higher 99% percentile value. The estimates differed due to the difference in distributions, where GEV is specifically for extreme events, and the lognormal is worse at capturing the extreme values (tail values)

4.) How might you decide (based on getting additional information, if needed) which distribution is “better”?

The means are slightly shifted, and the mean that more closely aligns with the real data may show a more accurate results? It also depends on what data we are looking for, the lognormal may show a more accurate value for the mean, but given how a GEV distribution works, it may give better results if we are looking for the extreme tail values.

References

Put any consulted sources here, including classmates you worked with/who helped you.