
NUMERICAL COMPUTATION OF HADAMARD FINITE PART OF SOME HYPERSINGULAR INTEGRALS IN 1, 2 AND 3 DIMENSIONS

Ziyue Qi

School of Computing and Information
University of Pittsburgh
Pittsburgh, PA 15213
ziq2@pitt.edu

Zixuan Lu

School of Computing and Information
University of Pittsburgh
Pittsburgh, PA 15213
ZIL50@pitt.edu

Yuchen Lu

School of Computing and Information
University of Pittsburgh
Pittsburgh, PA 15213
yul217@pitt.edu

July 15, 2026

ABSTRACT

Data mining is a good way to find the relationship between raw data and predict the target we want which is also widely used in different field nowadays. In this project, we implement a lots of technology and method in data mining to predict the sale of an item based on its previous sale. We create a strong model to predict the sales. After evaluating this model, we conclude that this model can be used in normal life for future sale's prediction.

1 Introduction

Our project is a competition on Kaggle (Predict Future Sales). We are provided with daily historical sales data (including each products' sale date, block ,shop price and amount). And we will use it to forecast the total amount of each product sold next month. Because of the list of shops and products slightly changes every month. We need to create a robust model that can handle such situations.

2 Notations

For every physical quantity, the corresponding reference space quantity is denoted by a hat, so that, for example, if $\mathbf{x}$ is a point on a d -dimensional variety, then $\hat{\mathbf{x}}$ is the corresponding point in the reference space. If u is a function defined on the variety, then $\hat{u}$ is the corresponding function in the reference space. Furthermore, we denote by $\oint$ the Hadamard finite part of an integral.

3 Model presentation

Let's consider an finite part integral of the form

$$I = \oint_{\hat{\tau}} f(\hat{\mathbf{y}}) d\hat{\mathbf{y}}, \quad f(\hat{\mathbf{y}}) \underset{\hat{\mathbf{y}} \rightarrow \hat{\mathbf{x}}}{\sim} \frac{1}{\|\hat{\mathbf{y}} - \hat{\mathbf{x}}\|^{d+1}}. \quad (1)$$

4 Injecting Richardson extrapolation into Guiggiani's final formula

4.1 Richardson extrapolation - general idea

For any angle θ , let's consider a finite sequence of n radial points $\rho_i(\theta)$ such that

$$0 < \rho_1(\theta) < \rho_2(\theta) < \dots < \rho_n(\theta) < \hat{\rho}(\theta), \quad (2)$$

Recall: we can express the two first terms of the Laurent expansion of the complete polar kernel $F(\rho, \theta) = \rho K(\chi(\hat{\mathbf{x}}), \chi(\hat{\mathbf{y}}(\rho, \theta)))J(\hat{\mathbf{y}}(\rho, \theta))\hat{u}(\hat{\mathbf{y}}(\rho, \theta))$ as the following limits

$$F_{-2}(\theta) = \lim_{\rho \rightarrow 0} \rho^2 F(\rho, \theta), \quad F_{-1}(\theta) = \lim_{\rho \rightarrow 0} \rho \left(F(\rho, \theta) - \frac{F_{-2}(\theta)}{\rho^2} \right). \quad (3)$$

such that we can write the complete polar kernel as

$$F(\rho, \theta) = \frac{F_{-2}(\theta)}{\rho^2} + \frac{F_{-1}(\theta)}{\rho} + O_{\rho \rightarrow 1}(1) \quad (4)$$

Let $c_{1,i}$ and $c_{2,i}$ be the coefficients used in the Richardson extrapolation of order n to compute the previous limits, so that we can write

$$\boxed{F_{-2}(\theta) \approx \tilde{F}_{-2}(\theta) = \sum_{i=1}^n c_{2,i} \rho_i^2(\theta) F(\rho_i(\theta), \theta)}, \quad F_{-1}(\theta) \approx \tilde{F}_{-1}(\theta) = \sum_{i=1}^n c_{1,i} \rho_i(\theta) \left(F(\rho_i(\theta), \theta) - \frac{\tilde{F}_{-2}(\theta)}{\rho_i^2(\theta)} \right). \quad (5)$$

By injecting the formula of $\tilde{F}_{-2}(\theta)$ into the formula of $\tilde{F}_{-1}(\theta)$, so that we can write $\tilde{F}_{-1}(\theta)$ only as a linear combination of the complete polar kernel evaluated at the radial points $\rho_i(\theta)$, we obtain

$$\tilde{F}_{-1}(\theta) = \sum_{i=1}^n c_{1,i} \rho_i(\theta) F(\rho_i(\theta), \theta) - \sum_{i=1}^n c_{1,i} \frac{1}{\rho_i(\theta)} \sum_{j=1}^n c_{2,j} \rho_j^2(\theta) F(\rho_j(\theta), \theta).$$

By letting $P_n(\theta) = \sum_{k=1}^n \frac{1}{\rho_k(\theta)}$ be a constant that is a polynomial in $\cos \theta$ and $\sin \theta$, we can write the previous formula as

$$\boxed{\tilde{F}_{-1}(\theta) = \sum_{i=1}^n F(\rho_i(\theta), \theta) \rho_i(\theta) \left(c_{1,i} - c_{2,i} P_n(\theta) \rho_i(\theta) \right)} \quad (6)$$

4.2 Lagrange basis method

For this section and the following, we will no longer use θ in the notation, it will be implicit. Let's denote the regular function $G(\rho) = \rho^2 F(\rho)$, so that $G(0) = F_{-2}$ and $G'(0) = F_{-1}$. Let $\ell_i(\rho)$ be the Lagrange basis of degree $n-1$ associated to the radial points ρ_i , defined as

$$\ell_i(\rho) = \prod_{j \neq i} \frac{\rho - \rho_j}{\rho_i - \rho_j}, \quad i = 1, \dots, n. \quad (7)$$

The lagrange interpolant of G and its derivative are given by

$$\Pi_{n-1} G(\rho) = \sum_{i=1}^n G(\rho_i) \ell_i(\rho), \quad (\Pi_{n-1} G)'(\rho) = \sum_{i=1}^n G(\rho_i) \ell'_i(\rho). \quad (8)$$

So that the numerical Laurent's coefficients, denoted $\tilde{F}_{-2}$ and $\tilde{F}_{-1}$, are simply the evaluation of the interpolant and its derivative at the origin $\rho = 0$:

$$\tilde{F}_{-2} = \Pi_{n-1}G(0) = \sum_{i=1}^n G(\rho_i)\ell_i(0), \quad \tilde{F}_{-1} = (\Pi_{n-1}G)'(0) = \sum_{i=1}^n G(\rho_i)\ell'_i(0). \quad (9)$$

with $\ell_i(0) = \prod_{j \neq i} \frac{\rho_j}{\rho_j - \rho_i}$ and $\ell'_i(0) = \ell_i(0) \left(\frac{1}{\rho_i} - P_n \right)$, so that we can finally write the Laurent's coefficients as

$$\tilde{F}_{-2} = \sum_{i=1}^n \ell_i(0)\rho_i^2 F(\rho_i), \quad \tilde{F}_{-1} = \sum_{i=1}^n \ell_i(0)\rho_i(1 - P_n\rho_i) F(\rho_i). \quad (10)$$

This is a particular case of the Richardson extrapolation, with the two families of coefficients $c_{1,i}$ and $c_{2,i}$ collapsing into a single family of coefficients $\ell_i(0)$:

$$c_{1,i} = c_{2,i} = \ell_i(0) = c_i = \prod_{j \neq i} \frac{\rho_j}{\rho_j - \rho_i}.$$

Finally, the numerical Laurent's coefficients can be written as

$$\tilde{F}_{-2}(\theta) = \sum_{i=1}^n A_i(\theta)F(\rho_i(\theta), \theta), \quad \tilde{F}_{-1}(\theta) = \sum_{i=1}^n B_i(\theta)F(\rho_i(\theta), \theta). \quad (11)$$

With $A_i(\theta) = c_i(\theta)\rho_i^2(\theta)$ and $B_i(\theta) = c_i(\theta)\rho_i(\theta)(1 - P_n(\theta)\rho_i(\theta))$.

Each of these formula simply consist in the numerical evaluation of the polar kernel at the radial points $\rho_i(\theta)$, multiplied by some weights.

4.2.1 Properties of $A_i(\theta)$ and $B_i(\theta)$

$$\sum_{i=1}^n A_i = \begin{cases} \rho_1, & n = 1 \\ -\rho_1\rho_2, & n = 2 \\ 0, & n \geq 3 \end{cases}, \quad \sum_{i=1}^n B_i = \begin{cases} 0, & n = 1 \\ \rho_1 + \rho_2, & n = 2 \\ 0, & n \geq 3 \end{cases} \quad (12)$$

These three cases illustrates the following definition :

$$\sum_{i=1}^n A_i = (\Pi_{n-1}[\rho^2])(0), \quad \sum_{i=1}^n B_i = (\Pi_{n-1}[\rho^2])'(0) \quad (13)$$

4.3 Final quadrature rule

Let's then denote $I(\theta)$ the integral of the complete polar kernel over the radial direction, defined as:

$$I(\theta) = \int_0^{\hat{\rho}(\theta)} \left\{ F(\rho, \theta) - \frac{F_{-1}(\theta)}{\rho} - \frac{F_{-2}(\theta)}{\rho^2} \right\} d\rho + F_{-1}(\theta) \ln \left| \frac{\hat{\rho}(\theta)}{\beta(\theta)} \right| - F_{-2}(\theta) \left\{ \frac{\gamma(\theta)}{\beta^2(\theta)} + \frac{1}{\hat{\rho}(\theta)} \right\} \quad (14)$$

A final quadrature rule for it is the following:

$$I(\theta) \approx \tilde{I}(\theta) = \sum_{i=1}^n \tilde{w}_i(\theta)F(\rho_i(\theta), \theta) \quad (15)$$

With the new weights $\tilde{w}_i(\theta)$ defined as

$$\tilde{w}_i(\theta) = w_i(\theta) - A_i(\theta) \left\{ W_{-2,n}(\theta) + \frac{\gamma(\theta)}{\beta^2(\theta)} + \frac{1}{\hat{\rho}(\theta)} \right\} + B_i(\theta) \left\{ \ln \left| \frac{\hat{\rho}(\theta)}{\beta(\theta)} \right| - W_{-1,n}(\theta) \right\}$$

and the auxiliary constants :

- $w_i(\theta) = \frac{\hat{\rho}(\theta)}{2} w_i^{\text{Ref}}$ are the weights of the reference quadrature over the segment $[-1, 1]$.
- $W_{-1,n}(\theta) = \sum_{i=1}^n \frac{w_i(\theta)}{\rho_i(\theta)} = \sum_{i=1}^n \frac{w_i^{\text{Ref}}}{1+x_i^{\text{Ref}}}$.
- $W_{-2,n}(\theta) = \sum_{i=1}^n \frac{w_i(\theta)}{\rho_i^2(\theta)} = \frac{2}{\hat{\rho}(\theta)} \sum_{i=1}^n \frac{w_i^{\text{Ref}}}{(1+x_i^{\text{Ref}})^2}$.

Remark: in the case of Gauss-Legendre quadrature, the constants $W_{-1,n}$ and $W_{-2,n}$ can be computed analytically as:

$$W_{-1,n} = 2H_n, \quad W_{-2,n}(\theta) = \frac{2}{\hat{\rho}(\theta)} n(n+1), \quad P_n(\theta) = \frac{n(n+1)}{\hat{\rho}(\theta)} \quad (16)$$

With $H_n = \sum_{i=1}^n \frac{1}{i}$ the n -th harmonic number. These last two formulas are, for the moment, a conjecture. (Elaborate to prove it?)

4.4 Properties of the new weights $\tilde{w}_i(\theta)$

$$\text{For } n \geq 3, \quad \sum_{i=1}^n \tilde{w}_i(\theta) = \hat{\rho}(\theta) \quad (17)$$

This is easy to understand: either look at the definition of $\tilde{w}_i(\theta)$ and use the properties of $A_i(\theta)$ and $B_i(\theta)$, or integrate the constant function $F(\rho, \theta) = 1$ between 0 and $\hat{\rho}(\theta)$, which is equivalent. In a more general way, if the integrand F is not singular (i.e. $F \sim \mathcal{O}(1)$), then the quadrature rule is the standard quadrature as if reference quadrature was used to integrate the function F between 0 and $\hat{\rho}(\theta)$. If the integrand is singular as $1/\rho$, the customized quadrature just performs a Cauchy principal value. The true ingredients are the values of the Laurent coefficients F_{-1} and F_{-2} , which are computed numerically; they can be zero (up to the algorithm precision) or not, depending on the considered singularity.

4.5 Convergence and error study

4.5.1 General idea

To begin our reasoning, assume that we have computed the Laurent's coefficients up to a relative error ϵ_{-k} , $k \in \{1, 2\}$:

$$\tilde{F}_{-k} = F_{-k}(1 + \epsilon_{-k}), \quad k \in \{1, 2\} \quad (18)$$

Now let's denote

$$R(\rho, \theta) = F(\rho, \theta) - \frac{F_{-1}(\theta)}{\rho} - \frac{F_{-2}(\theta)}{\rho^2} \quad (19)$$

$$\tilde{R}(\rho, \theta) = F(\rho, \theta) - \frac{\tilde{F}_{-1}(\theta)}{\rho} - \frac{\tilde{F}_{-2}(\theta)}{\rho^2} \quad (20)$$

$$(21)$$

$$I = \int_0^{2\pi} \int_0^{\hat{\rho}(\theta)} R(\rho, \theta) d\rho d\theta, \quad \tilde{I} = \int_0^{2\pi} \int_0^{\hat{\rho}(\theta)} \tilde{R}(\rho, \theta) d\rho d\theta \quad (22)$$

On one hand, after the theory of the Bernstein's ellipses, if we assume that R and $\tilde{R}$ are analytic, we can make a error majoration regarding the Gauss-Legendre quadrature of I and $\tilde{I}$:

$$|I - Q_n[R]| \leq \frac{64}{15} \frac{M}{(1 - \rho_B^{-2})\rho_B^{2n}} \quad (23)$$

$$|\tilde{I} - Q_n[\tilde{R}]| \leq \frac{64}{15} \frac{\tilde{M}}{(1 - \tilde{\rho}_B^{-2})\tilde{\rho}_B^{2n}} \quad (24)$$

On the other hand, we can subtract $\tilde{I}$ to I to get:

$$|I - \tilde{I}| \leq C(\hat{x}) \sum_{k=1}^2 \frac{\epsilon_{-k}}{\rho_{\min}^k} \|F_{-k}\|_{\infty} \quad (25)$$

where $C(\hat{x}) = \int_0^{2\pi} \hat{\rho}(\theta) d\theta$.

Thus,

$$|I - Q_n[\tilde{R}]| = |I - \tilde{I} + \tilde{I} - Q_n[\tilde{R}]| \leq |I - \tilde{I}| + |\tilde{I} - Q_n[\tilde{R}]| \leq C(\hat{x}) \sum_{k=1}^2 \frac{\epsilon_{-k}}{\rho_{\min}^k} \|F_{-k}\|_{\infty} + \frac{64}{15} \frac{\tilde{M}}{(1 - \tilde{\rho}_B^{-2})\tilde{\rho}_B^{2n}} \quad (26)$$

4.5.2 Richardson extrapolation

We can show that the following majorations hold for any sequence $\rho_i(\theta)$ of n radial points:

$$|\tilde{F}_{-2}(\theta) - F_2(\theta)| \leq \frac{\|R^{(n)}\|_{\infty}(\theta)}{n!} \prod_{i=1}^n \rho_i(\theta), \quad |\tilde{F}_{-1}(\theta) - F_1(\theta)| \leq P_n(\theta) \frac{\|R^{(n+1)}\|_{\infty}(\theta)}{n!} \prod_{i=1}^n \rho_i(\theta) \quad (27)$$

Which shows that the speed of convergence is conditioned by the product of the radial points $\prod_{i=1}^n \rho_i(\theta)$. In the case of a geometric sequence $\rho_i(\theta) = \hat{\rho}(\theta)q^{i-1}$, we have $\prod_{i=1}^n \rho_i(\theta) = \hat{\rho}^n(\theta)q^{\frac{n(n-1)}{2}}$, which is a very fast convergence. In the case of a sequence of Gauss-Legendre quadrature points, we have $\prod_{i=1}^n \rho_i(\theta) = \hat{\rho}^n(\theta) \frac{(n!)^2}{(2n)!}$, which is a slower convergence (see Figure 1).

That's why geometric sequences are still preferred over Gauss-Legendre quadrature points for pure Richardson extrapolation.

4.5.3 Gauss-Legendre quadrature

To delve deeper into these general ideas, let's consider :

$$\tilde{I}_{\epsilon,h,n}(\theta) = Q_n[\tilde{R}_{\epsilon,h}](\theta) = \sum_{i=1}^n w_i(\theta) \left\{ F_h(\rho_i(\theta), \theta) - \frac{\tilde{F}_{-1,\epsilon,h}(\theta)}{\rho_i(\theta)} - \frac{\tilde{F}_{-2,\epsilon,h}(\theta)}{\rho_i^2(\theta)} \right\} \quad (28)$$

$$= \sum_{i=1}^n \left\{ F_h(\rho_i(\theta), \theta) - \frac{F_{-1,h}(\theta)}{\rho_i(\theta)} - \frac{F_{-2,h}(\theta)}{\rho_i^2(\theta)} \right\} w_i(\theta) - \sum_{k=1}^2 F_{-k,h}(\theta) \epsilon_{-k} \sum_{i=1}^n \frac{w_i(\theta)}{\rho_i^k(\theta)} \quad (29)$$

$$= Q_n[R_h](\theta) - \sum_{k=1}^2 F_{-k,h}(\theta) \epsilon_{-k} W_{-k,n}(\theta) \quad (30)$$

Where h is the mesh size. Thanks to standard results on the Gauss-Legendre quadrature for any function in $C^{2n}([0, \hat{\rho}(\theta)])$, we have the following error majoration (deleting the θ dependency for clarity):

$$Q_n[R_h] = \int_0^{\hat{\rho}} R_h(\rho) d\rho + \frac{2^{2n+1}(n!)^4}{(2n+1)[(2n)!]^3} R_h^{(2n)}(\rho_0) \quad (31)$$




Figure 1: Convergence comparison between geometric and Gauss-Legendre sequences for the function $\rho \rightarrow 1/(1 + \rho)$

for some $\rho_0 \in [0, \hat{\rho}]$. Our next goal is to show that the error term $R_h^{(2n)}(\rho_0)$ converges with the mesh size h and the number of quadrature points n . By using $G_h(\rho) = \rho^2 F_h(\rho) = \mathcal{O}_{\rho \rightarrow 0}(1)$, we can straightforwardly express R_h in terms of G , and bound its derivatives, assuming, as Guiggiani, that F_h is analytic in a neighborhood of the origin:

$$R_h(\rho) = \sum_{k=0}^{+\infty} F_k \rho^k, \quad \Rightarrow \quad R_h^{(2n)}(\rho) = \sum_{k=2n}^{+\infty} F_k \frac{k!}{(k-2n)!} \rho^{k-2n} \quad (32)$$

$$(33)$$

with $F_k = \frac{G_h^{(k+2)}(0)}{(k+2)!} \leq \frac{1}{(k+2)!} \|G_h^{(k+2)}\|_{\infty, [0, \hat{\rho}]}$ and $\rho^{k-2n} \leq \hat{\rho}^{k-2n}$, so that we can write

$$|R_h^{(2n)}(\rho)| \leq \sum_{k=2n}^{+\infty} k! \frac{\|G_h^{(k+2)}\|_{\infty, [0, \hat{\rho}]}}{(k+2)! (k-2n)!} \hat{\rho}^{k-2n} \quad (34)$$

BOUNDING OF G AND ITS DERIVATIVES: let us defined the following functions:

$$\begin{aligned} \phi_h : \hat{\tau} \subset \mathbb{R}^{d-1} &\rightarrow \mathbb{R} & \psi_{\hat{x}} : \mathbb{R} \times [0, 2\pi[&\rightarrow \mathbb{R}^{d-1} \\ \hat{x} &\mapsto \det M_h(\hat{x}) & (\rho, \theta) &\mapsto \hat{x} + \rho \mathbf{u}(\theta) \end{aligned}$$

$$\begin{aligned} \chi_h : \hat{\tau} \subset \mathbb{R}^{d-1} &\rightarrow \tau \subset \mathbb{R}^d \\ \hat{x} &\mapsto x \end{aligned}$$

where χ is the mapping from the reference space to the physical element, $\psi_{\hat{x}}$ is the mapping from polar coordinates to the reference space, and M_h is the metric tensor of the mapping χ_h , defined as:

$$M_h(\hat{x}) = D\chi_h(\hat{x})^T D\chi_h(\hat{x}), \quad (M_h)_{ab} = \sum_{k=1}^{d-1} \frac{\partial \chi_{h,k}}{\partial \hat{x}_a} \frac{\partial \chi_{h,k}}{\partial \hat{x}_b}, \quad (a, b) \in \{1, \dots, d-1\}^2 = \{1, 2\}^2 \quad (35)$$

Lemma 1: using the Faà di Bruno formula for functions of the form $h(x) = f[g_1(x), \dots, g_m(x)]$ (see Constantine, 1996), one can write, for any function $f : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$,

$$\frac{\partial^k (f \circ \psi_{\hat{x}})}{\partial \rho^k}(\rho, \theta) = \sum_{i=0}^k \binom{k}{i} \cos^i(\theta) \sin^{k-i}(\theta) \frac{\partial^k f}{\partial \hat{x}_1^i \partial \hat{x}_2^{k-i}}(\psi_{\hat{x}}(\rho, \theta)) \quad (36)$$

Finally, the vector $\mathbf{u}(\theta)$ is the unit trigonometric vector defined as $\mathbf{u}(\theta) = (\cos \theta, \sin \theta)$ for our case $d = 3$. We can then expand G as

$$G(\rho, \theta) = H_h(\rho, \theta) J_h \circ \psi_{\hat{x}}(\rho, \theta) \quad (37)$$

where

$$H_h(\rho, \theta) = \rho^3 K(x = \chi_h(\hat{x}), y = \chi_h \circ \psi_{\hat{x}}(\rho, \theta)) \hat{u} \circ \psi_{\hat{x}}(\rho, \theta), \quad J_h(\hat{x}) = \sqrt{\phi_h(\hat{x})} = \sqrt{\det M_h(\hat{x})} \quad (38)$$

As the kernel K is regularized by the term ρ^3 in H_h , we can ensure that $H_h = \mathcal{O}_{\rho \rightarrow 0}(1)$. Applying the Leibniz formula, we have

$$\frac{\partial^m G(\rho, \theta)}{\partial \rho^m} = \sum_{k=0}^m \binom{m}{k} \frac{\partial^k H_h(\rho, \theta)}{\partial \rho^k} \frac{\partial^{m-k} (J_h \circ \psi_{\hat{x}})}{\partial \rho^{m-k}}(\rho, \theta) \quad (39)$$

We then let

$$\begin{cases} \gamma_{h,m} = \sup_{k \leq m} \sup_{\theta \in [0, 2\pi[} \sup_{\rho \in [0, \hat{\rho}]} \left\| \frac{\partial^k H_h(\rho, \theta)}{\partial \rho^k} \right\| = O_{h \rightarrow 0}(1) \\ M_m = \max_{k \leq m} \binom{m}{k} = \binom{m}{\lfloor m/2 \rfloor} \end{cases} \quad (40)$$

So that we can write

$$\left| \frac{\partial^m G(\rho, \theta)}{\partial \rho^m} \right| \leq (m+1) M_m \gamma_{h,m} \sum_{k=0}^m \left| \frac{\partial^k (J_h \circ \psi_{\hat{x}})}{\partial \rho^k}(\rho, \theta) \right| \quad (41)$$

BOUNDING OF THE k -TH RADIAL DERIVATIVE OF $J_h \circ \psi_{\hat{x}}$: using the lemma 36, we can write

$$\frac{\partial^k (J_h \circ \psi_{\hat{x}})}{\partial \rho^k}(\rho, \theta) = \sum_{i=0}^k \binom{k}{i} \cos^i(\theta) \sin^{k-i}(\theta) \frac{\partial^k J_h}{\partial \hat{x}_1^i \partial \hat{x}_2^{k-i}}(\psi_{\hat{x}}(\rho, \theta)) \quad (42)$$

$$\leq (k+1) M_k \sum_{i=0}^k \left| \frac{\partial^k J_h}{\partial \hat{x}_1^i \partial \hat{x}_2^{k-i}}(\psi_{\hat{x}}(\rho, \theta)) \right| \quad (43)$$

BOUNDING OF THE CARTESIAN DERIVATIVES OF J_h : there's no need to expand an explicit formula for the derivatives of J_h in terms parametrization χ_h , because we can use a useful hypothesis on the mesh quality. Indeed, to ensure non-degeneracy of the elements, we can assume that

$$\exists C_Q > 0, \quad \forall h > 0, \quad \forall m > 0, \quad \forall \hat{x} \in \hat{\tau}, \quad \|D^m \chi_h(\hat{x})\| \leq C_Q h^m \quad (44)$$

Where $D^m \chi_h(\hat{x})$ is the tensor of order m defined containing all the m -th order derivatives of χ_h at $\hat{x}$, defined as

$$(D^m \chi_h(\hat{x}))_{i, \alpha_1, \dots, \alpha_m} = \frac{\partial^m \chi_{h,i}}{\partial \hat{x}_{\alpha_1} \dots \partial \hat{x}_{\alpha_m}}(\hat{x}), \quad i \in \{1, 2, 3\}, \quad (\alpha_1, \dots, \alpha_m) \in \{1, 2\}^m \quad (45)$$

In particular, for $\alpha = 1$, we have

$$\|M_h(\hat{x})\| = \|D\chi_h(\hat{x})^T D\chi_h(\hat{x})\| \leq C_Q^2 h^2. \quad (46)$$

Furthermore, the Hadamard's inequality (HI) states that

$$|\phi_h(\hat{x})| = |\det(M_h(\hat{x}))| \stackrel{\text{(HI)}}{\leq} \prod_{i=1}^{d-1} \|M_h(\hat{x})_{i,:}\| \leq \|M_h(\hat{x})\|^{d-1} \leq C_Q^{2(d-1)} h^{2(d-1)} \quad (47)$$

Remarks : theses inequalities holds perfectly if we consider the ∞ -norm, but all the norms are equivalent in finite dimension, so we can use any norm; that leads to different majoration constants but the idea remains the same. This ensure that we can define the following function:

$$\begin{aligned} \mu &: \hat{\tau} \subset \mathbb{R}^{d-1} \rightarrow \mathbb{R} \\ \hat{x} &\mapsto \sqrt{h^{-2(d-1)} \det M_h(\hat{x})} = h^{-(d-1)} J_h(\hat{x}) \leq C_Q^{d-1} \end{aligned}$$

By the non-degeneracy hypothesis, we have $\mu = \mathcal{O}_{h \rightarrow 0}(1)$ and all its derivatives are also bounded. So that we can write $J_h(\hat{x}) = h^{d-1} \mu(\hat{x})$, and

$$\left| \frac{\partial^m G(\rho, \theta)}{\partial \rho^m} \right| \leq h^{d-1} (m+1) M_m \gamma_{h,m} \sum_{k=0}^m (k+1) M_k \sum_{i=0}^k \left\| \frac{\partial^i \mu}{\partial \hat{x}^i \partial \hat{x}^{k-i}} (\psi_{\hat{x}}(\rho, \theta)) \right\| \quad (48)$$

Naturally, let now $\mu_k = \sup_{i \leq k} \left\| \frac{\partial^k \mu}{\partial \hat{x}_1^i \partial \hat{x}_2^{k-i}} \right\|_{\infty, \hat{\tau}}$ and $\lambda_m = \max_{k \leq m} \mu_k$, we have the simplified following majoration:

$$\left| \frac{\partial^m G(\rho, \theta)}{\partial \rho^m} \right| \leq h^{d-1} (m+1)^4 M_m^2 \gamma_{h,m} \lambda_m \quad (49)$$

Thus, the remainder term of Laurent's serie expansion can be bounded as

$$|R_h^{(2n)}(\rho)| \leq h^{d-1} \sum_{k=2n}^{+\infty} k! \frac{(k+3)^4 M_{k+2}^2 \gamma_{h,k+2} \lambda_{k+2}}{(k+2)! (k-2n)!} \hat{\rho}^{k-2n} \quad (50)$$

EXPLICIT MAJORATION OF $\gamma_{h,k}$: let's recall the definition of $\gamma_{h,k}$:

$$\gamma_{h,k} = \sup_{m \leq k} \sup_{\theta \in [0, 2\pi[} \sup_{\rho \in [0, \hat{\rho}]} \left\| \frac{\partial^m H_h(\rho, \theta)}{\partial \rho^m} \right\| \quad (51)$$

where

$$\frac{\partial^m H_h(\rho, \theta)}{\partial \rho^m} = \frac{\partial^m}{\partial \rho^m} \{ \rho^s K(\chi_h(\hat{x}), \chi_h \circ \psi_{\hat{x}}(\rho, \theta)) \hat{u} \circ \psi_{\hat{x}}(\rho, \theta) \} \quad (52)$$

$$= \sum_{k=0}^m \binom{m}{k} \frac{\partial^k}{\partial \rho^k} \left(\rho^s K(\chi_h(\hat{x}), \chi_h \circ \psi_{\hat{x}}(\rho, \theta)) \right) \frac{\partial^{m-k}}{\partial \rho^{m-k}} \left(\hat{u} \circ \psi_{\hat{x}}(\rho, \theta) \right) \quad (53)$$

$$\stackrel{36}{\leq} (m+1)^2 \delta_m M_m^2 \sum_{k=0}^m \frac{\partial^k}{\partial \rho^k} \left(\rho^s K(\chi_h(\hat{x}), \chi_h \circ \psi_{\hat{x}}(\rho, \theta)) \right) \quad (54)$$

where $\delta_m = \sup_{i \leq m} \left\| \frac{\partial^m \hat{u}(\hat{x})}{\partial \hat{x}_1^i \partial \hat{x}_2^{m-i}} \right\|_{\infty, \hat{\tau}}$ is a constant that depends only on the geometry of the element, and s is the order of singularity of the function $\rho \rightarrow K(\chi_h(\hat{x}), \chi_h \circ \psi_{\hat{x}}(\rho, \theta))$ at the origin, such that $K(\chi_h(\hat{x}), \chi_h \circ \psi_{\hat{x}}(\rho, \theta)) = \mathcal{O}_{\rho \rightarrow 0}(\rho^{-s})$.

The idea was originally used for hypersingular 3D kernels, where $s = 3$ (see Guiggiani, 1992), but we write the computations with general order s because the idea remains the same.

Lemma 2: if $f : \mathbb{R} \rightarrow E$ is a function of class $\mathcal{C}^m(E)$, then we have the following useful calculation:

$$\frac{\partial^n}{\partial \rho^n} \left(\rho^s f(\rho) \right) = \sum_{j=\max\{s-n, 0\}}^n \binom{n}{j} \frac{s!}{(s-n+j)!} \rho^{s-n+j} f^{(j)}(\rho) \quad (55)$$

Thus,

$$\frac{\partial^k}{\partial \rho^k} \left(\rho^s K(\chi_h(\hat{x}), \chi_h \circ \psi_{\hat{x}}(\rho, \theta)) \right) = \sum_{j=\max\{k-s, 0\}}^k \binom{k}{j} \frac{s!}{(s-k+j)!} \rho^{s-k+j} \frac{\partial^j}{\partial \rho^j} K(\chi_h(\hat{x}), \chi_h \circ \psi_{\hat{x}}(\rho, \theta)) \quad (56)$$

(57)

Then, we notice, as well as another article (T.G. Anderson, M. Bonnet, L.M. Faria and C. Pérez-Arancibia, 2024, formula 6.27), that the partial derivatives of the kernel K are bounded:

$$\|D_y^\alpha K(x, y)\| \leq C_K(\alpha) \|x - y\|^{-(|\alpha|+s)}, \quad \alpha \in \mathbb{N}^d, \quad |\alpha| = \sum_{i=1}^d \alpha_i \quad (58)$$