

3(a)(i)	Table 3.1 completed correctly ;	1										
	<table><tr><td>type of fruit</td><td>mean of samples</td></tr><tr><td>apple</td><td>17 / 16.7</td></tr><tr><td>orange</td><td>33 / 33.3</td></tr><tr><td>pineapple</td><td>43 / 43.3</td></tr><tr><td>grapes</td><td>14 / 14.3</td></tr></table>	type of fruit	mean of samples	apple	17 / 16.7	orange	33 / 33.3	pineapple	43 / 43.3	grapes	14 / 14.3	
type of fruit	mean of samples											
apple	17 / 16.7											
orange	33 / 33.3											
pineapple	43 / 43.3											
grapes	14 / 14.3											
3(a)(ii)	Table 3.2 completed correctly ;	2										
	<table><tr><td></td><td>type of variable</td><td>type of data</td></tr><tr><td>type of fruit</td><td>categoric / qualitative</td><td>nominal</td></tr><tr><td>volume of juice extracted</td><td>quantitative</td><td>continuous</td></tr></table>		type of variable	type of data	type of fruit	categoric / qualitative	nominal	volume of juice extracted	quantitative	continuous		
	type of variable	type of data										
type of fruit	categoric / qualitative	nominal										
volume of juice extracted	quantitative	continuous										

Question	Answer	Marks
3(a)(iii)	any two from: 1 standardise, (total) mass / volume, of fruit used or standardise number of pieces of fruit used or standardise, dimensions / mass / volume / size / surface area / AW, of (each fruit) piece ; 2 use same, age / ripeness of each fruit ; 3 use a more accurate method to measuring the volume (of pectinase) or use a more accurate method to measure the volume (of juice) ; 4 replace pectinase with, (distilled) water / denatured pectinase / boiled pectinase / AW ;	2
3(b)	any two from: 1 <i>idea that</i> the volume of juice may not be correlated with the quantity of pectin ; 2 age / ripeness, of fruit affects quantity of juice ; 3 <i>idea of</i> cell walls of different fruits have different compositions ;	2