

Question	Marking Guidance	Mark	Comments
03.1	1. Tip produces IAA; 2. IAA diffuses (into shoot); 3. (More) <u>elongation</u> of <u>cells</u> on one side (than other);	3	1. Accept source/release for produces but ignore contains/stores IAA. 1 and 2. Accept auxin for IAA. 2. Accept IAA diffuses down. 3. Accept (more) <u>elongation</u> of <u>cells</u> on left side. 3. Reject any reference to shaded/dark side or away from light.

Question	Marking Guidance	Mark	Comments
03.2	1. Size of shoot/tip; 2. Number of shoot tips; 3. Size/type of agar (block); 4. (Shoots) at same stage of growth/development; 5. Time (period) tips kept on agar OR Time (period) agar/block kept on (cut shoot) OR Time (period shoots) kept in dark; 6. Temperature; 7. (Repeat several times and) calculate a mean; 8. Compare/read degree of curvature (on calibration curve) to determine (IAA) concentration OR Higher the degree of curvature the higher the IAA concentration;	5 max	Mark points 1 to 6 = max 3. 1 to 6. Ignore pH, species, carbon dioxide, humidity, nutrients, water and light. 3. Accept 'amount of agar'. 4. Accept (Shoots/plants) are same age.

Question	Marking Guidance	Mark	Comments
03.3	1. (IAA) is not broken down by light OR (IAA) is produced in the dark OR Light/dark does not affect (IAA) production; 2. (IAA) moves away from light OR (IAA) moves to shaded side;	2	2. IAA accumulates on shaded side is not enough on its own, idea of movement is required.