

Question number	Answer	Additional guidance	Mark
2(a)	using a {thermometer / (temperature) probe} to take the temperature of the {liver / rectum}	ACCEPT into the core / deep into the body / up the anus IGNORE other parts of body	(1)

Question number	Answer	Additional guidance	Mark
2(b)(i)	- drop in body temperature in first 12 hours calculated and subtracted from 11.5°C (1) - this value divided by 0.4, added to 12 hours and answer rounded to nearest hour (1)	$11.5 - (0.78 \times 12) / 11.5 - 9.36 / 2.14$ 17 (hours) 17.35 = 1 mark Correct answer with no working gains 2 marks	(2)

Question number	Answer	Additional guidance	Mark
2(b)(ii)	An explanation that includes the following points: (this) estimate would be {shorter / an under-estimate} (1) - because a body loses <u>heat</u> faster (in cooler conditions) (1)	ACCEPT converse	(2)

Question number	Answer	Additional guidance	Mark
2(c)	An explanation that includes three of the following points: • because temperature affects {rigor / body stiffness} (1) • because deciding when a body is stiff or not stiff is subjective (1) • because if the body is stiff, the time since death can only be estimated as being between 3 and 36 hours (1) • because if the body is not stiff, there is no way of telling if it has been dead for less than 3 hours or more than 36 hours (1)	ACCEPT exercise / body shape / body fat / ATP levels ACCEPT gives a wide range of (time) values ACCEPT if not stiff cannot tell how many hours after 36 hours time of death was	(3)

