

.....[1]

The graph displays two data series over time. The solid line represents deaths from TB, which starts at approximately 60 in 1925, peaks at 80 in 1930, and then declines steadily to near zero by 1970. The dashed line represents new cases of TB, which starts at approximately 80 in 1925, peaks at 100 in 1950, and then declines steadily to near zero by 1970. Both series show a significant downward trend over the 85-year period.

Year	Deaths from TB (per 100,000)	New Cases of TB (per 100,000)
1925	60	80
1930	80	80
1935	60	80
1940	50	90
1945	45	100
1950	25	100
1955	10	90
1960	5	55
1965	2	30
1970	1	20
1975	0.5	15
1980	0.2	10
1985	0.1	8
1990	0.1	7
1995	0.1	6
2000	0.1	5
2005	0.1	4
2010	0.1	3

The vaccine for TB was introduced in Canada for widespread use from 1948.

Use the information in Fig. 3.1 to comment on the effect of the introduction of the vaccine and antibiotics on the number of new cases and deaths from TB.

.....[5]

- (c) In 2010, Swaziland had one of the highest death rates from TB in the world.

Swaziland also had a high number of new cases of HIV infection in its population in 2010.

People who are infected with HIV are described as HIV+ and people who are **not** infected with HIV are described as HIV–.

Table 3.1 shows the number of deaths from TB in Swaziland in 2010.

Table 3.1

HIV status	number of deaths from TB per 100 000 population
HIV+	400
HIV–	91

Using the information in Table 3.1, suggest why the number of deaths per 100 000 population of people who are HIV+ is much higher than that in people who are HIV–.

.....

.....

.....

.....

.....

.....

.....

.....

.....[3]

[Total: 9]