

Question Number	Answer	Additional guidance	Mark
3(a)(i)	- calculation of {percentage / proportion} of rod cells (1) - correct number of rod cells in section (1)	5/6 or 0.83 or 83.3% or 83.33% or 83% 2dp's MAX 4167 Accept 4150 or 4166 or 4170 Accept in standard form 4.166×10^3 or 4.17×10^3 or 4.2×10^3 ignore 4167:833 4167.8 ignore 4166.7 or 4166.6 ignore 4.16×10^3	(2)

Question Number	Answer	Additional guidance	Mark
3(a)(ii)	An explanation that includes three of the following points: - rod cells can detect low light intensities / cone cells require higher light intensities (for colour vision) (1) - squirrel monkey has more cones as active {during the day / when there is more light} (1) - the {grey bellied / night} monkey has {fewer cone cells / more rod cells} as it {is active at night / needs to see in dark} (1) - both monkeys need to be able to see colour (to find food) therefore both have similar order of magnitude of cone cells (1)	ACCEPT squirrel monkey has more cones as spends more time in the light ACCEPT the night monkey needs more peripheral vision ACCEPT converse for squirrel monkey	(3)

Question Number	Answer	Additional guidance	Mark
3(b)	An answer that includes three of the following points: • (seedling 2) auxin caused {growth towards the direction of light / (positive) phototropism / cell elongation on the shaded side} (1) • (seedling 2) auxin inhibited lateral bud growth (1) • (seedling 3) gibberellins caused increased {growth / length} of seedling (1) • (seedling 3) gibberellins caused growth of lateral buds (1)	ignore side shoots ignore 'bending' ACCEPT auxins cause apical dominance ACCEPT increase in internode length / stem elongation / increased height / got taller ACCEPT gibberellins cause {new lateral bud production / more leaves}	(3)