

10(a)	<i>any six from:</i> 1 <u>nucleus</u> in, cell body / soma ; 2 (long) dendron ; 3 (short) axon ; 4 many mitochondria (in cell body) ; 5 many, RER / ribosomes or presence of Nissl's granules (in cell body) ; 6 synaptic, knobs / terminals / boutons ; 7 Schwann cells / myelin sheath ; 8 nodes of Ranvier ;	6
10(b)	<i>any nine from:</i> 1 Na^+ moves out of, cell and K^+ moves into cell ; 2 <u>3</u> Na^+ for every <u>2</u> K^+ ; 3 by, active transport / use of ATP ; 4 sodium-potassium pump / $\text{Na}^+ \text{K}^+$ pump ; 5 against concentration gradient ; 6 K^+ <u>diffuses</u> out of cell and Na^+ <u>diffuses</u> into cell ; 7 by facilitated <u>diffusion</u> / <u>diffusion</u> through (ion) channels ; 8 membrane more permeable to K^+ / more K^+ goes out than Na^+ in ; 9 inside of, cell / membrane, more negative than outside ; ora 10 membrane / cell, polarised / repolarised ; 11 (resting potential is), -60 mV / -65 mV / -70 mV ; 12 AVP ; e.g. ion movement only at nodes of Ranvier	9