

2(a)(i)	<i>max two if, number of (reported) cases / AW, not stated any three from:</i> 1 overall decrease in number of reported cases from 1980 to 2000 / over the time period / AW ; 2 from, 27 600 / 27 700 / 27 800, to, 16 300 / 16 400, cases ; A manipulated data e.g. decline of, 11 200 / 11 300 / 11 400 / 11 500 cases 3, 4 additional trends ;; e.g. decrease from 1980 to 1984 increase between 1988 and 1992 or (overall) increase between 1984 to 1992 decrease after 1992 (to 2000) little change / slight increase and decrease / AW, between 1984 and 1988 5 use of comparative values from Fig. 2.1 to support mps, 3 / 4 ;	3
2(a)(ii)	HIV-infected people may develop a weak(ened) immune system or people with, HIV/AIDS have a weak(ened) immune system ; A poor immune response A <i>idea of</i> more susceptible to disease caused by, <i>Mycobacterium, tuberculosis / bovis</i> A TB is an opportunistic disease (of people with HIV/AIDS) increased risk of latent TB becoming active in (many) people, living with HIV / with HIV/AIDS ; AW detail of weak immune system ; e.g. decrease in number of ,T-lymphocytes / T helper cells poor, humoral / B-lymphocyte, response AVP ; <i>idea of</i> more cases of TB owing to more people infected with HIV increases risk of spreading to uninfected people	2

Question	Answer	Marks
2(b)	any two from: 1 prescribe / take, antibiotics, only when (absolutely) necessary ; A examples e.g. do not use for viral infections do not use as preventative medicine 2 make sure, correct / effective, antibiotic(s), prescribed / used AW A <i>ref. to</i>, broad spectrum / narrow spectrum, antibiotics, qualified or make sure people take the correct dose or complete course / follow instructions for use, of antibiotics A <i>ref. to</i> DOTS ; 3 develop new, antibiotics / antibacterials ; 4 use other antibacterials ; <i>if mp3 and 4 not gained, allow one mark for vary antibiotic treatment</i> or use a number of different antibiotics at the same time 5 (named) measures to, prevent infection / break transmission cycle (to reduce the need for antibiotic use) ; e.g. vaccines good hygiene in hospitals quarantine people who have multi-drug resistant bacterial infections 6 report, cases / patterns, of antibiotic resistance ; 7 reduce / control, antibiotics in, agriculture / animals used for food ; 8 <i>ref. to</i> education for, healthcare professionals / public, to reduce impact of antibiotic resistance ; 9 AVP ; e.g. <i>ref. to</i> monitoring situation to check if antibiotic is effective WHO Global Plan to End TB	2
2(c)(i)	(bronchi) may, collapse / lose (structural) support ;	1

Question	Answer	Marks																				
2(c)(ii)	one mark each correct <u>column</u> = 3 marks ;;; if 0 or 1 mark gained, check for correct row to max 2 <i>In the whole table:</i> • <i>if only ticks are used, assume blanks are crosses</i> • <i>if only crosses are used, assume blanks are ticks</i> <i>If there are ticks <u>and</u> crosses and blanks in the whole table, R the columns or rows with the blank(s)</i> <table><tr><th>cell type</th><th>alveolus</th><th>bronchus</th><th>trachea</th></tr><tr><td>ciliated epithelial cell</td><td>×</td><td>✓</td><td>✓</td></tr><tr><td>goblet cell</td><td>×</td><td>✓</td><td>✓</td></tr><tr><td>smooth muscle cell</td><td>×</td><td>✓</td><td>✓</td></tr><tr><td>squamous epithelial cell</td><td>✓</td><td>×</td><td>×</td></tr></table>	cell type	alveolus	bronchus	trachea	ciliated epithelial cell	×	✓	✓	goblet cell	×	✓	✓	smooth muscle cell	×	✓	✓	squamous epithelial cell	✓	×	×	3
cell type	alveolus	bronchus	trachea																			
ciliated epithelial cell	×	✓	✓																			
goblet cell	×	✓	✓																			
smooth muscle cell	×	✓	✓																			
squamous epithelial cell	✓	×	×																			

Question	Answer	Marks
----------	--------	-------