


Question Number	Answer	Additional guidance	Mark
2(a)	An answer that includes the following points: - capsule correctly drawn and labelled (1) (at least) two pili correctly drawn and labelled (1)	<u>Example of diagram</u> 	(2)

Question Number	Answer	Additional guidance	Mark
2(b)	An answer that makes reference to four of the following points: • increasing sodium chloride concentrations (above 10 g dm³) decreases the growth rate (of both bacteria) (1) • (salt concentration between 0-6 g dm³ caused) an initial {increase / constant} rate (1) • more rapid decrease in <i>L. piscium</i> growth rate (1) • <i>B. thermosphacta</i> had a larger growth rate (than <i>L. piscium</i> at all sodium chloride concentrations) (1) • <i>B. thermosphacta</i> was able to continue growing in sodium chloride concentrations above {22-25} g dm³ whereas <i>L. piscium</i> had no growth / <i>L. piscium</i> stopped growing at a concentration {38-46} g dm³ below the concentration that <i>B. thermosphacta</i> stopped growing (1) • data is more scattered about the line of best fit for <i>B. thermosphacta</i> (than <i>L. piscium</i>)	ACCEPT negative correlation (between sodium chloride concentration and growth rate) ACCEPT converse ACCEPT converse Accept <i>B. thermosphacta</i> is more {tolerant / resistant} (than <i>L. piscium</i>) of higher sodium chloride concentrations Accept <i>B. thermosphacta</i> stopped growing at {62-70} g dm³ whereas <i>L. piscium</i> stopped growing at {22-24} g dm³ ACCEPT converse	(4)