

Question	Marking Guidance	Mark	Comments
09.1	1. Two divisions produce 4 (haploid) cells/gametes/nuclei; 2. Separation of homologous chromosomes (in first division); 3. Separation of chromatids (in second division); 4. Independent segregation/assortment of homologous chromosomes; 5. Crossing over leads to exchange of parts of chromatids/alleles between homologous chromosomes;	5 (5 x AO1)	1. Accept meiosis I and meiosis II 1. Accept description that clearly describes two divisions 3. Accept 'chromosomes' for 'chromatids' but reject homologous chromosomes. Accept annotated diagrams for MP1–MP3 4. Accept description eg maternal and paternal chromosomes are reshuffled in any combination

Question	Marking Guidance	Mark	Comments
09.2	(Formation at arterial end) 1. Water/small molecules/glucose/ions (forced) out of the capillary/blood; 2. (Due to) high hydrostatic/blood pressure (at arterial end); 3. (Plasma) proteins remain in capillary/blood; (Return at venous end) 4. (Plasma) proteins create water potential gradient OR (Plasma) proteins reduce/low(er) water potential (of blood/in capillary) OR (Plasma) proteins cause more negative water potential (of blood/in capillary); 5. Water moves (into blood/capillary) by <u>osmosis</u>; 6. Excess (tissue) fluid enters lymph (system/capillaries/vessels);	5 max (5 x AO1)	2. Accept hydrostatic/blood pressure greater than osmotic effect 3. Accept named plasma proteins 4. and 5. Accept Ψ for water potential but not wp/WP 6. Accept lacteals