

1(a)	<table border="1"> <thead> <tr> <th data-bbox="723 217 1227 272"></th><th data-bbox="1227 217 1482 272">region of kidney</th><th data-bbox="1482 217 1525 272"></th></tr> </thead> <tbody> <tr> <td data-bbox="723 272 1227 328">location of loops of Henle</td><td data-bbox="1227 272 1482 328">C</td><td data-bbox="1482 272 1525 328">;</td></tr> <tr> <td data-bbox="723 328 1227 384">location of Bowman's capsules</td><td data-bbox="1227 328 1482 384">A</td><td data-bbox="1482 328 1525 384">;</td></tr> <tr> <td data-bbox="723 384 1227 440">location of glomeruli</td><td data-bbox="1227 384 1482 440">A</td><td data-bbox="1482 384 1525 440">;</td></tr> <tr> <td data-bbox="723 440 1227 507">contains urine at final concentration</td><td data-bbox="1227 440 1482 507">B + D</td><td data-bbox="1482 440 1525 507">;</td></tr> </tbody> </table>		region of kidney		location of loops of Henle	C	;	location of Bowman's capsules	A	;	location of glomeruli	A	;	contains urine at final concentration	B + D	;	4
	region of kidney																
location of loops of Henle	C	;															
location of Bowman's capsules	A	;															
location of glomeruli	A	;															
contains urine at final concentration	B + D	;															
1(b)	any three from: 1 (aquaporins are) water <u>channel</u> (proteins) ; 2 (more) aquaporins increase (cell surface) membrane permeability (to water) ; ora 3 of collecting duct (cells) ; I distal convoluted tubule cells 4 allow water to, be reabsorbed / move into tissue fluid or blood ;	3															
1(c)	any three from: 1 detected by osmoreceptors ; 2 in hypothalamus ; 3 (osmoreceptors send) fewer impulses to posterior pituitary ; I signals 4 less ADH, released / produced ; I no ADH released	3															