

8(a)	any three from: 1 make cDNA from, messenger RNA / mRNA ; 2 using reverse transcriptase ; 3 synthesise (gene chemically) from <u>nucleotides</u> / join <u>nucleotides</u> ; 4 (using known) amino acid / nucleotide, sequence from database ;				3																
8(b)	<table border="1"><thead><tr><th>role in genetic engineering</th><th>DNA ligase</th><th>DNA polymerase</th><th></th></tr></thead><tbody><tr><td>joins two sections of sugar phosphate backbone in DNA</td><td>✓</td><td>✗</td><td>;</td></tr><tr><td>adds a gene to a plasmid</td><td>✓</td><td>✗</td><td>;</td></tr><tr><td>adds free activated DNA nucleotides to a polynucleotide</td><td>✗</td><td>✓</td><td>;</td></tr></tbody></table>				role in genetic engineering	DNA ligase	DNA polymerase		joins two sections of sugar phosphate backbone in DNA	✓	✗	;	adds a gene to a plasmid	✓	✗	;	adds free activated DNA nucleotides to a polynucleotide	✗	✓	;	3
role in genetic engineering	DNA ligase	DNA polymerase																			
joins two sections of sugar phosphate backbone in DNA	✓	✗	;																		
adds a gene to a plasmid	✓	✗	;																		
adds free activated DNA nucleotides to a polynucleotide	✗	✓	;																		
8(c)	1 90–98°C – DNA, denatures / strands separate ; 2 50–65°C – primers, bind / base pair / anneal (to DNA) ; 3 68–75°C – Taq / DNA, polymerase, makes, DNA / (new) strand ;				3																