

2(a)	any three from: 1 capture a large sample (from one area / more than one area) ; 2 count / record, the number caught ; 3 test <u>and</u>, count / record, the number infected ; 4 number / population, of infected divided by, population / total number, $\times 100$; 5 mark-release-<u>recapture</u> method <u>and</u> Lincoln Index ;	3
2(b)(i)	numbers (of infected badgers / cows) per unit area ; allow (valid) comparison (between infected cows and infected badgers) ;	2
2(b)(ii)	any three from: 1 method of testing (for bTB in badgers and cows) ; 2 year of data collection ; 3 type / breed / of, cattle ; 4 cattle kept, outdoors / indoors ; 5 number of areas counted each year ; 6 season / time of year, (data collected) ;	3
2(c)(i)	no, correlation / relationship, between the number of infected badgers and the number of affected cows. or no, correlation / relationship, in the occurrence of infection between badgers and cows ;	1

Question	Answer	Mark
2(c)(ii)	<i>max 2</i> support 1 combined data / A–J / graph / Fig 2.2, shows a positive, correlation / relationship / correct correlation described ; 2 combined data / A–J / graph / Fig 2.2, shows a significant, correlation / relationship ; 3 (so) culling (infected) badgers lowers infection in cattle ; <i>max 2</i> does not support – graph 4 separated data / A–H and D–J are clustered ; or no data between 0.4 and 0.6 per km² ; 5 data taken from different years ; does not support – table 6 separated data / A–H and D–J, shows a negative, correlation / relationship ; 7 separated data / A–H and D–J, does not show a significant, correlation / relationship ; 8 correlation does not mean causation ; 9 only data sets from 10 areas / only 10 out of 30 study areas sampled ; 10 data only from England / one country ;	3