

6(a)	<table><tr><th colspan="5">Stage of cell cycle</th></tr><tr><td>(G₁ phase)</td><td rowspan="3">}</td><td rowspan="3">interphase ;</td><td rowspan="3">}</td><td rowspan="3"></td></tr><tr><td>S phase</td></tr><tr><td>G₂ phase</td></tr><tr><td>prophase</td><td rowspan="4">}</td><td rowspan="4">(mitosis)</td><td rowspan="4">}</td><td rowspan="4"></td></tr><tr><td>metaphase</td></tr><tr><td>anaphase</td></tr><tr><td>(telophase)</td></tr><tr><td colspan="5">(cytokinesis)</td></tr></table> one mark for interphase one mark for S phase and G₂ phase in correct sequence one mark for prophase, metaphase and anaphase in correct sequence	Stage of cell cycle					(G ₁ phase)	}	interphase ;	}		S phase	G ₂ phase	prophase	}	(mitosis)	}		metaphase	anaphase	(telophase)	(cytokinesis)					3
Stage of cell cycle																											
(G ₁ phase)	}	interphase ;	}																								
S phase																											
G ₂ phase																											
prophase	}	(mitosis)	}																								
metaphase																											
anaphase																											
(telophase)																											
(cytokinesis)																											
6(b)	Vincristine: ✓ ✓ x x ; 5-fluorouracil : x x x x ;	2																									

Question	Answer	Marks
6(c)	<i>any three from</i> <i>blood smear differences</i> large / larger, numbers of lymphocytes ; more lymphocytes than normal blood smear ; <i>as blood smear or lymphocyte difference</i> immature / not fully developed ; <i>ref. to</i> no large nuclei / nuclei not pronounced small nuclei ; A nuclei not visible <i>lymphocyte difference</i> non-functional / AW ; detail of lack of function for either B-lymphocytes or T-lymphocytes ; AVP ; e.g. <i>ref. to</i> difficult to distinguish between lymphocytes and monocytes ; A no monocytes visible	3