

1(a)	maintain / keep / restore, constant / stable / set-point / within narrow limits, internal environment / in body ;	1
1(b)(i)	A – pelvis ; <i>note if labelled medulla as affects ecf in part (ii)</i> B – ureter ;	2
1(b)(ii)	A <i>full labels instead of letters</i> <i>if region A (pelvis) was mislabelled as medulla in (i) can apply:</i> ecf for L placed in pelvis ecf U placed in medulla only if word cortex also written by U/ultrafiltration U – pointing to the cortex ; L – pointing to the medulla ; C – pointing to the renal vein ;	3

Question	Answer	Marks
1(c)	<i>max 5 of:</i> 1 <u>hypothalamus</u> detects (changes in) <u>water potential</u> (of the blood) ; 2 <u>osmoreceptors</u> shrink when, low / less, water in <u>blood</u> ; ora 3 ADH, produced / made, in hypothalamus ; 4 if low, water / Ψ, ADH secreted from <u>posterior pituitary</u> ; ora R ADH <i>produced</i> in posterior pituitary 5 <i>ref. to neurosecretory cells</i> or impulse / ADH transported, from hypothalamus to posterior pituitary ; 6 aquaporins ; 7 ADH increases <u>permeability</u> of, distal convoluted tubule / collecting duct ; ora 8 ADH causes, more water reabsorption / smaller volume of urine / more concentrated urine ; ora A both with and without ADH compared	5