

Question number	Answer	Mark
6(a)	The only correct answer is C (two) <i>A is incorrect because thromboplastin is an active enzyme found in the platelets but is not converted into thrombin</i> <i>B is incorrect because thromboplastin is an active enzyme found in the platelets but is not converted into thrombin</i> <i>D is incorrect because thromboplastin is an active enzyme found in the platelets but is not converted into thrombin</i>	(1)

Question number	Answer	Additional guidance	Mark
6(b)	An answer that includes the following points: • 1.8 to 3.5 (1) • below 1.8 there is a high risk of stroke (1) • above 3.5 there is an increased risk of bleeding (within the skull) (1) OR • 1.8 to 2.5 (1) • below 1.8 there is a high risk of stroke (1) • above 2.5 the risk of bleeding (within the skull) starts to increase (1) OR • 2.0 to 2.5 (1) • both risks are low / they are the lowest (combination) (1) • 3.1-3.5 has a lower risk of stroke but a higher risk of bleeding (1)	ACCEPT other ranges {have either high risk of bleeding or high risk of stroke / both risks are low} = 1 mark if mp 2 and 3 not awarded ACCEPT other ranges {have either higher risk of bleeding or high risk of stroke / both risks are low} = 1 mark if mp 2 and 3 not awarded	(3)

Question number	Answer	Additional guidance	Mark
6(c)	An explanation that includes the following points: - warfarin reduces {a blood clot forming / thrombosis} (1) (if a blood clot does not form) it cannot block the <u>coronary artery</u> (1) - if oxygen is supplied to the heart {muscle / cells} it will continue to {contract / respire} (1)	ACCEPT no blood clots forming less blood clotting reduces blood clot (size)	(3)

Question number	Answer	Additional guidance	Mark
6(d)(i)	1 012 / 1 013 (1)		(1)

Question number	Answer	Additional guidance	Mark
6(d)(ii)	An answer that includes the following points: • select a group of patients who have had a heart attack (1) • of similar {age / sex at birth / background / lifestyle} (as in the first study) (1) • give each person the same quantity of warfarin (used in the first study) (1) • (monitor for) {the same time period / four years} (1) • use of a statistics (1)	ACCEPT other appropriate factors no other health issues ACCEPT in context of one group split into two groups ACCEPT description of double-blind trial / double-blind trial / suitable named statistics test	(4)