

6(a)	G – embryo to aleurone layer ; A – aleurone layer to endosperm ;	2
6(b)	any three from: 1 hydrolyses / breaks down / digests / converts, starch / amylose ; 2 in endosperm ; 3 to, maltose / glucose ; 4 (glucose) used by embryo ; 5 (glucose used) for, respiration / growth ;	3

Question	Answer	Marks
6(c)	<i>any four from:</i> <i>accept distilled water for A, 500mg dm⁻³ for B and 1000mg dm⁻³ for C.</i> 1 seeds of B and C germinated before seeds of A or seeds soaked in gibberellin germinated before seeds soaked in distilled water ; ora 2 rate of (increase in percentage) germination was, highest in C or higher in C than, B / A or higher as the concentration of gibberellin increases ; ora 3 final / day 10 / maximum (percentage) germination was, highest in C or higher in C than, B / A or higher as the concentration of gibberellin increases ; ora 4 data quote between two groups ; mp1 – A by day 6 and B and C by day 2 mp1 – A germinates four days after B and C ora mp2 – A – 2.2 B – 5.2 C – 9.2, percentage germination day⁻¹ mp3 – A – 22% B – 52% C – 92% mp3 – A to C = 70% more, A to B = 30% more, B to C = 40% more 5 percentage germination in C levels off after 9 days ; 6 AVP ; e.g. no seeds germinate by day 1	4