


Question Number	Answer	Additional Guidance	Mark
2ai	An answer that includes the following points • addition of (Benedict's / Fehling's) solution (1) • heat / place in water bath (and observe colour change) (1)	Water bath must be heated or > 50 C	2

Question Number	Answer	Additional Guidance			Mark
2aii		e.g.			2

Question Number	Answer	Additional Guidance	Mark
2bi	An answer that includes the following points: - describe how mean obtained (1) - describe how standard deviation obtained (1)	e.g substitute mean and number of observations into equation.	2

Question Number	Answer	Additional Guidance	Mark
2bii	A graph with the following features • L1 x label mean urine glucose concentration / au (1) • L2 y- mean blood glucose concentration / au (1) • P1 plots correct on a linear scale on both axes (1) • P2 standard deviation plotted correctly (1) • suitable line of best fit drawn (1)	 Mean Blood Glucose Concentration / au Mean Urine Glucose Concentration / au IGNORE extrapolation	5

Question Number	Answer	Additional Guidance	Mark
2biii	An answer that includes the following points: - there is a correlation between the 2 variables / variables may depend on each other / as urine glucose goes up blood glucose goes up (1) - however, the {SDs / error bars} overlap (1) - suggesting that differences not significant)(1)	NOT vice versa to the last alternative NOT range bars	3

Question Number	Answer	Additional Guidance	Mark
2biv	An calculation that includes the following steps : - substitution of values into equation (1) - solving of equation (1) - correct rounding (1)	e.g. $y = 29.3 \times 12 + 154.1$ $= 505.7$ rounded to 506 allow ecf for $12 = 29.3 \times X + 154.1$ $= - 4.849 \text{ (ignore sign)}$ Rounded to 4.85	3