

2(a)	capillary ; <i>two from</i> single celled wall / AW ; A thin wall <i>ref. to</i> wall composed only of one, squamous epithelial / endothelial, cell ; <i>ref. to</i> lumen diameter approx 7 μm ; A 5–8 μm A <i>ref. to</i> similarity to dimension of red blood cell in direct contact with tissue fluid ; <i>if vein named, allow one mark for, thin wall (relative to lumen) or wide / AW, lumen (relative to wall thickness)</i>	3
2(b)(i)	0.24 s / 0.25 s ;	1
2(b)(ii)	0.08 s / 0.09 s ; A range	1
2(b)(iii)	<i>four from</i> <i>similarity</i> <i>ref. to</i> increases and decreases in pressure at same time ; A description for part of the graph e.g.starts to rise at same point as RV, returns to minimum at same point as RV <i>idea that</i> events in cardiac cycle occur are coordinated ; A described e.g. impulses pass up both ventricles at the same time, ventricles both contract at same time <i>difference</i> reaches higher, blood pressure / peak, (than RV) ; systolic pressure higher in LV (than RV) left ventricle pumps blood to (whole) body / RV only to lungs) overcome greater resistance / or for RV walls of left ventricle, thicker / more muscular more force exerted by LV	max 4 <i>any two ;;</i>