

number		
2(a)	The only correct answer is B. <i>A is incorrect because anticoagulants prevent blood clotting.</i> <i>C is incorrect because platelet inhibitors reduce blood clotting.</i> <i>D is incorrect because statins reduce cholesterol levels.</i>	(1)

Question number	Answer	Additional guidance	Mark
2(b)	An explanation that includes the following points: - because there is a link between (total) blood cholesterol levels and heart disease (1) - because {HDL is not thought to be a risk factor / LDL is associated with the development of heart disease / the {ratio / balance} of LDL : HDL determines the level of risk} (1)	ACCEPT CHD / CVD / atherosclerosis for heart disease throughout ACCEPT causes / leads / is a risk factor / correlation ACCEPT HDL reduces risk / HDL needed to get the LDL	(2)

number			
2(c)	An explanation that includes the following points: (10 year CHD risk) increases (1) - because smoking {increases blood pressure / increases heart rate / damages (endothelial) lining of arteries} (1) - damage to lining results in {atheroma / plaque} building up OR {high blood pressure / increase in heart rate} makes the heart work harder (1)	ACCEPT CVD / CHD / heart disease / atherosclerosis / throughout ACCEPT increases the chance of blood clotting / makes platelets sticky ACCEPT blood clots can block arteries	(3)

Question number	Answer		Mark
2(d)(i)	salt intake / fibre intake / alcohol intake / (body) mass / BMI / obesity / hip waist ratio / level of exercise / stress levels / family history / genetic factors / taking statins / diastolic blood pressure	IGNORE LDL / antihypertensives / blood pressure	(1)

Question	Answer	Additional guidance	Mark
----------	--------	---------------------	------

number			
2(d)(ii)	{salt / low fibre / alcohol / high BMI / obesity / little exercise / stress / family history / genetic factors / high diastolic blood pressure} increase the risk value OR {low salt / high fibre / low alcohol / suitable mass / exercise / low stress levels / taking statins / low diastolic blood pressure} decrease the risk value	ACCEPT if answer given in (i)	(1)

Question number	Answer	Additional guidance	Mark
2(d)(iii)	An explanation that includes two of the following points: - because there are other (risk) factors not included (1) - because people will underestimate {their mass / how much they smoke} (1) - because {total cholesterol / HDL / blood pressure} might be an estimate (1)	ACCEPT named factor but not those shown on the risk calculator ACCEPT {total cholesterol / HDL / blood pressure} are variable are not aware that they have diabetes	(2)