

1(a)	3 = (atrial) systole (ventricular) diastole 5 = (atrial) diastole (ventricular) systole 7 = (atrial) diastole (ventricular) diastole	} ; } ; } ;	2
1(b)	<i>allow responses in context of both sides or one side of heart</i> <i>any two from:</i> atria and ventricles, do not contract / are not in systole, at the same time ; AW ventricles, contract / are in systole, after atria ; AW ora <i>idea that allows atria to empty (completely) / allows ventricles to fill ;</i> A idea that (all) blood, pumped / AW, into ventricles (ready for systole)		2

Question	Answer	Marks
1(c)	<i>opening of tricuspid valve</i> pressure in (right) ventricle becomes lower than pressure in (right) atrium or pressure in (right) atrium becomes higher than pressure in (right) ventricle ; <i>opening of pulmonary (semilunar) valve</i> pressure in (right) ventricle becomes higher than pressure in pulmonary artery or pressure in pulmonary artery becomes lower than pressure in (right) ventricle ; <i>if mp1 and 2 have correct ideas but not referred to pressures on both 'sides' of the valve, allow one mark for idea of, increasing / high, pressure opening <u>each</u> of the valves</i> <i>help flow of blood through the heart</i> any two from: one-way flow of blood / prevents backflow of blood / directs blood in correct direction / blood flows in one direction / AW ; A blood flows from atrium to ventricle to, pulmonary artery / lungs blood flows from (area of) higher pressure to (area of) lower pressure ; <i>idea that (blood) pressure changes aid coordination ;</i> e.g. semilunar valve is open (about same time) when tricuspid valve is shut	3