

2(a)(i)	<i>any 2 from:</i> 1 remove all / add, larvae ; 2 measure larvae and select, medium-sized / 9–16 mm, larvae; 3 <i>ref. to</i> way of measuring ; 4 suitable equipment for handling larvae, e.g. paint brush, pooter, (blunt) forceps ; 5 place larvae on plants, randomly / on cotton bolls / on buds / AW ; 6 <i>idea of</i> breeding cotton bollworms to use for treatment 2 ; 7 AVP ;	2
2(a)(ii)	<i>any 2 from:</i> 1 variety / type, of (Bt) cotton ; 2 number of rows of cotton ; 3 (1 m) length of cotton sections per test ; 4 number of plants in each section ; 5 (20 m²) size of the plot ; 6 size of larvae ; 7 density of larvae in each treatment ; 8 time before spraying with insecticide ; 9 cotton bolls picked at same time when all bolls in each plot were open / AW ;	2
2(a)(iii)	(as a control) to compare the, quantity / mass / weight / yield, of cotton (fibres, and /or, seeds) from plots that were not sprayed with insecticide with that from the plots that were sprayed / AW ;	1
2(b)(i)	<i>any 1 from:</i> 1 compares the means of two sets of data ; 2 the data is continuous ; 3 the populations have a normal distribution ; 4 standard deviations are similar ;	1
2(b)(ii)	there is no (significant) difference in the mass of cotton fibre produced when sprayed with insecticide or when not sprayed with insecticide ;	1

Question	Answer	Marks
2(c)	<i>any 2 from:</i> 1 (mean) yield of cotton bolls increases when sprayed with insecticide ; 2 (mean) yield of cotton fibre increases, less / little / unchanged / AW, when sprayed ; 3 spraying makes little difference to the (mean) yields of bolls and fibres between treatments 1 and 2 ; 4 standard deviations appear large (e.g. for bolls sprayed in treatment 1 and 2) ; 5 mean $\pm$ SD overlap so no conclusion can be made ; 6 no results given for, <i>t</i>-tests / statistical tests ; 7 AVP ;	2