

Question number	Answer	Mark
6(a)(i)	The correct answer is A ($\frac{5}{11}$) <i>B is incorrect because it is $(3.3-1.8) \div 3.3 = \frac{5}{11}$</i> <i>C is incorrect because it is $(3.3-1.8) \div 3.3 = \frac{5}{11}$</i> <i>D is incorrect because it is $(3.3-1.8) \div 3.3 = \frac{5}{11}$</i>	(1)

Question number	Answer	Mark
6(a)(ii)	The correct answer is B (logarithmic) <i>A is incorrect because this is already a linear scale</i> <i>C is incorrect because standard deviation is not an axis scale</i> <i>D is incorrect because standard form is not an axis scale</i>	(1)

Question number	Answer	Additional guidance	Mark
6(b)	An answer that includes the following points: - each subtype had {same / similar} antigens (1) - the vaccine contained {two types / more than one type} of antigen (1)	ACCEPT vaccine contained two types of {attenuated / deactivated} virus vaccine stimulates production of more than one type of antibody	(2)

Question number	Answer	Additional guidance	Mark
6(c)(ii)	An answer that includes three of the following points: • pregnant women were used and were given the (RSV) vaccine (1) • one large group of {pregnant vaccinated women / babies of these pregnant vaccinated women / babies with passive immunity to RSV} (1) • babies (from vaccinated pregnant women) monitored for a period of time (1) • number of babies who developed RSV infections requiring medical attention and number not requiring medical attention recorded OR number of babies who developed RSV not requiring medical attention divided by the total number of infected babies ($\times 100$) (1)	ACCEPT stated period of time up to 2 years ACCEPT converse for calculating percentage who did need medical attention	(3)

