

8(a)	<table border="1"> <thead> <tr> <th>function</th><th>part of seed</th><th></th></tr> </thead> <tbody> <tr> <td>gibberellin synthesis</td><td>embryo</td><td>;</td></tr> <tr> <td>starch store</td><td>endosperm</td><td>;</td></tr> <tr> <td>amylase synthesis</td><td>aleurone layer</td><td>;</td></tr> <tr> <td>maltose production</td><td>endosperm</td><td>;</td></tr> </tbody> </table>	function	part of seed		gibberellin synthesis	embryo	;	starch store	endosperm	;	amylase synthesis	aleurone layer	;	maltose production	endosperm	;	4
function	part of seed																
gibberellin synthesis	embryo	;															
starch store	endosperm	;															
amylase synthesis	aleurone layer	;															
maltose production	endosperm	;															
8(b)(i)	any three from: 1 protons / H^+ ions, pumped / actively transported (into cell wall) ; 2 (so) pH (in cell wall) decreases / (cell wall) becomes more acidic ; 3 expansins, activated / AW ; 4 microfibrils (in cellulose) loosen (to allow elongation) ; Ignore weaken	3															
8(b)(ii)	any three from: 1 K^+ / potassium ions, enter, cell / cytoplasm ; R active transport 2 water potential of, cell / cytoplasm, decreases or water potential of the cell wall increases ; 3 water enters (cell through aquaporins) by, osmosis / down the water potential gradient ; 4 cell volume increases / cell expands / increase in turgor pressure (inside cell) ;	3															