

6(a)(i)	1 as the eugenol concentration increases the percentage decrease in sodium ion movement increases ; 2 percentage at two concentrations of eugenol plus units ; <table><tr><td>conc (mmol dm⁻³)</td><td>%</td></tr><tr><td>0.0</td><td>0</td></tr><tr><td>0.2</td><td>30</td></tr><tr><td>0.4</td><td>50</td></tr><tr><td>0.6</td><td>65</td></tr><tr><td>0.8</td><td>72</td></tr><tr><td>1.0</td><td>80</td></tr></table>	conc (mmol dm ⁻³)	%	0.0	0	0.2	30	0.4	50	0.6	65	0.8	72	1.0	80	2
conc (mmol dm ⁻³)	%															
0.0	0															
0.2	30															
0.4	50															
0.6	65															
0.8	72															
1.0	80															
6(a)(ii)	57–58 (%) ;	1														

Question	Answer	Marks
6(b)	<i>five from:</i> 1 reduced entry of Na⁺ into sensory neurone ; 2 eugenol prevents opening of sodium ion channels / described ; 3 no / reduced, depolarisation of sensory neurone <u>membrane</u> ; 4 receptor potential / threshold potential, not reached ; 5 no / fewer, action potentials / impulses ; 6 may affect sodium-potassium pump ; 7 resting potential not restored ; 8 action potentials / impulses, do not reach brain ;	5