

- 9 Green lacewings are a family of insects with more than 1300 species.

The common green lacewing, *Chrysoperla carnea*, is shown in Fig. 9.1.




Fig. 9.1

- (a) Green lacewings have sense organs, known as tympanal organs, that detect sound.

The tympanal organ of green lacewings has evolved to detect the high frequency sounds that bats make when they are hunting. Bats eat green lacewings.

When a green lacewing senses the presence of a bat, it moves away or closes its wings in flight to escape.

- (i) Outline how the tympanal organ of green lacewings could have evolved by natural selection.

.....

.....

.....

.....

.....

.....

.....

.....

..... [3]

- (ii) When high frequency sound is detected, the receptor cells in the tympanal organ stimulate the transmission of impulses in sensory neurones.

Describe the sequence of events that results in an action potential in a sensory neurone.

The first event in the sequence has been given for you.

Calcium ions enter the cytoplasm of the receptor cell

.....

.....

.....

.....

.....

.....

.....

..... [4]

[4]

- (b) Two species of green lacewing, *C. carnea* and *C. downesi*, evolved from a common ancestor.

The two species have populations with overlapping distributions in parts of North America.

Table 9.1 shows a comparison of the characteristics of overlapping populations of the two species.

Table 9.1

characteristic	<i>C. carnea</i>	<i>C. downesi</i>
breeding months	June to September	April to May
courtship song	song with a regular rhythm	song with no regular rhythm
colour	light green	dark green

Suggest how speciation occurred to produce the two different species of green lacewing.

.....

.....

.....

.....

.....

.....

..... [3]

[Total: 10]