

7 The risk of developing cardiovascular disease (CVD) is affected by two groups of factors:

- lifestyle factors that can be changed
- non-lifestyle factors that cannot be changed.

Methods to reduce the risk of developing CVD include drug treatments and lifestyle changes.

(a) (i) Which row of the table identifies one lifestyle factor and one non-lifestyle factor?

(1)

	Lifestyle factor	Non-lifestyle factor
☐ A	body mass index (BMI)	age
☐ B	gender	high alcohol intake
☐ C	genetics	high blood pressure
☐ D	high blood cholesterol	inactivity

(ii) Explain why a person might have to take several types of drugs to reduce the risk of CVD.

(2)

.....

.....

.....

.....

.....

.....



(b) Nutritional studies have shown that dietary antioxidants can reduce the risk of CVD.

(i) Explain why antioxidants in the diet reduce the risk of CVD.

(3)

(ii) Some studies do not assess the nutritional quality of the diet of the participants.

Explain why the results of these studies have to be treated with caution.

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(iii) Explain why changes in diet, other than antioxidants, can reduce the risk of CVD.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 7 = 11 marks)

