

2(a)	EMRO <u>and</u> SEARO / Eastern Mediterranean <u>and</u> South-East Asia (Regions)	1
2(b)	<i>max 1 (tar data mark or nicotine makes platelets sticky) if tar / CO also stated any two from:</i> (AFRO) has more nicotine ; R has more, tar / tar and nicotine / CO and nicotine (see below) nicotine makes platelets sticky / AW ; A damages endothelial lining (so turbulent blood flow) so increases clotting risk R if also state tar makes platelets sticky 0.92 and 0.77–0.79 <u>mg</u> (per cigarette) or 9.2 v. 7.7–7.9 <u>mg</u> (per cigarette) or (AFRO) 1.3–1.5 mg (per cigarette) higher ; <i>correctly extracted numerical data for tar is alternative to gain mp 3 ecf</i> 12.5–12.7 v. 11.3–11.5 mg / (AFRO) 1.0–1.4 mg higher	2
2(c)	<i>allow, Hb / hb, for haemoglobin and allow CO for carbon monoxide any four from:</i> 1 haemoglobin, has a higher affinity for / binds more readily with, carbon monoxide (than oxygen) ; A carbon monoxide displaces oxygen from haemoglobin A carbon monoxide binds to Hb and decreases its affinity for oxygen 2 carboxyhaemoglobin formed ; 3 binding is, (mainly) irreversible / permanent / (more) stable (than oxygen binding) / AW ; 4 decreases saturation of haemoglobin with oxygen (in the lungs) / less haemoglobin available to bind oxygen ; A less oxyhaemoglobin formed / less oxygen binds to haemoglobin <i>cigarette smoke:</i> 5 low concentration of oxygen in inhaled cigarette smoke ; 6 <i>ref. to</i> (in airways), narrowed lumen / inflammation / increased mucus / accumulated, mucus ; A bronchial constriction 7 (so) lower concentration of / less, oxygen, reaching alveolus / in alveolar air or (so) less oxygen, diffuses / AW, (from alveolus) into bloodstream / reaches red blood cells / reaches haemoglobin (in red blood cells) ; R <i>ref. to</i> consequence of emphysema <i>must be in context of mp 5 or 6</i> 8 AVP ; e.g. nicotine causes vasoconstriction and reduces blood supply (to extremities)	4