

2(a)	<i>independent variable:</i> number / amount of (daily) garlic capsules ; <i>dependent variable:</i> (resting systolic) blood pressure ;	2	
2(b)	capsule with no garlic ;	1	ignore 'placebo' unqualified A placebo capsule allow tablet / pill, as AW for capsule A capsule with, sugar / starch / cellulose / AW, (and no garlic)
2(c)(i)	continuous ;	1	ignore quantitative / ordinal / ordered / dependent / interval / normally distributed R qualitative / categoric / discrete / discontinuous / nominal

Question	Answer	Marks	Guidance
2(c)(ii)	any three from: <i>comparing start to end of the investigation</i> (resting systolic) blood pressure decreased in all groups (over the 12 weeks) ; - SE for, start / 0 weeks and end / 12 weeks, do not overlap, therefore (mean resting systolic blood pressure may be) significantly different (in all groups) ; <i>comparing groups</i> - results / decrease in blood pressure, for 1 garlic capsule were not different to the control ; - SE for one garlic capsule and control, overlap (at all times) therefore (mean resting systolic blood pressure) is not significantly different ; - results / decrease in blood pressure, for 2 and 4 garlic capsules were different to the control or (taking) 2 garlic capsules gives, greatest / AW, decrease in blood pressure ; - SE for, 2 garlic capsules and control, do not overlap (at 8 weeks and 12 weeks) therefore (mean resting systolic blood pressure may be) significantly different or SE for, 4 garlic capsules and control, do not overlap (at 8 weeks) therefore (mean resting systolic blood pressure may be) significantly different ;	3	ignore improved blood pressure as AW for blood pressure decreases 2 allow if only one group is stated ignore standard deviation / 95%CI / error bars allow significant, decrease / AW (implies a difference) allow is significantly different ignore 'results are significant' 4 ignore standard deviation / 95%CI / error bars allow is not significantly different ignore no significant decrease ignore insignificant 6 ignore standard deviation / 95%CI / error bars allow significant, decrease / AW (implies a difference) allow is significantly different ignore 'results are significant'
2(c)(iii)	any two from: (data) not normally distributed ; - four / more than two, means to compare ; - more than one mean per group / means change over time ;	2	ignore ref. to standard deviations (because the SDs are approximately the same) R standard deviations are different / data is not continuous ignore fewer than 30 values

Question	Answer	Marks	Guidance
2(d)	any two from: 1 larger sample size / study more than 84 people ; 2 longer investigation / study for more than 12 weeks ; 3 standardise, age / sex / diet / smoking status / alcohol consumption / other medication taken / time of day capsule taken / AW ; 4 measurements of diastolic blood pressure ; 5 <i>ref. to</i> standardise measurement of blood pressure ;	2	ignore repeat unqualified 3 A exclude those already taking meds for blood pressure ignore ref. to take 3 garlic capsules 5 A measure blood pressure, at same time of day / at same location / same body position