

Question	Marking Guidance	Mark	Comments
07.1	1. Species richness – number of species (in a community/habitat/ecosystem/area); 2. Index of diversity – the relationship between the number of species (in a community) and the number of individuals in each species; 3. No SD overlap for species richness so significant difference shown OR No SD overlap for species richness so difference not due to chance; 4. SD overlap so no significant difference in index of diversity OR SD overlap so any difference shown due to chance;	4 (2 x AO1, 2 x AO2)	1. Reject number of species in a population. 2. Accept equation with N and n correctly explained
Question	Marking Guidance	Mark	Comments
07.2	1. DNA/genome sequencing now used; 2. (Now) can analyse every/more prokaryote species (in the community); 3. Rather than just recording measurable/observable characteristics;	2 max (2 x AO2)	1. Accept RNA/amino acid sequencing 2. Accept 'identify' for analyse
Question	Marking Guidance	Mark	Comments
07.3	(For farming method 2 has) 1. Better conservation as higher species richness, higher (prokaryotic) biomass and more carbon stored (than farming method 1); 2. More microbial mass – more organisms for food web; 3. Higher species richness – more organisms for food web; 4. More stored carbon, less CO₂ in the atmosphere; 5. Lower yield so less food/less profit;	2 max (2 x AO3)	Accept converse for farming method 1