

1(a)(i)	A – podocyte ; B – (capillary) endothelial cell ;				2																
1(a)(ii)	high(er) , hydrostatic / blood, pressure ; ignore <i>ref. to</i> water potential <u>basement membrane</u> acts as a filter / only small molecules can pass through <u>basement membrane</u> ;				2																
1(b)	<table><tr><th>component of blood plasma in glomerulus</th><th>component in newly formed glomerular filtrate</th><th>component in urine</th><th></th></tr><tr><td>glucose</td><td>same</td><td>not present</td><td>;</td></tr><tr><td>large proteins</td><td>not present</td><td>not present</td><td>;</td></tr><tr><td>urea</td><td>same</td><td>increased</td><td>;</td></tr></table>				component of blood plasma in glomerulus	component in newly formed glomerular filtrate	component in urine		glucose	same	not present	;	large proteins	not present	not present	;	urea	same	increased	;	3
component of blood plasma in glomerulus	component in newly formed glomerular filtrate	component in urine																			
glucose	same	not present	;																		
large proteins	not present	not present	;																		
urea	same	increased	;																		
1(c)	<i>posterior pituitary gland</i> – releases (more) ADH ; A secretes <i>kidney</i> – reabsorbs more water / urine more concentrated / lower volume of urine ; OR (more) aquaporins added to membrane of collecting duct (cells) / membrane of collecting duct (cells) more permeable to water ;				2																