

6(a)(i)	[A] C E D B F ;	1												
6(a)(ii)	I early and late except where indicated <table><tr><th>event in the cell cycle</th><th>name of the stage in the cell cycle</th></tr><tr><td>DNA replication</td><td>interphase / S phase ;</td></tr><tr><td>division of centromeres</td><td>anaphase ; R late anaphase</td></tr><tr><td>condensation of chromatin</td><td>prophase ;</td></tr><tr><td>contraction of spindle fibres</td><td>anaphase ;</td></tr><tr><td>organisation of chromosomes at the equator</td><td>metaphase</td></tr></table>	event in the cell cycle	name of the stage in the cell cycle	DNA replication	interphase / S phase ;	division of centromeres	anaphase ; R late anaphase	condensation of chromatin	prophase ;	contraction of spindle fibres	anaphase ;	organisation of chromosomes at the equator	metaphase	4
event in the cell cycle	name of the stage in the cell cycle													
DNA replication	interphase / S phase ;													
division of centromeres	anaphase ; R late anaphase													
condensation of chromatin	prophase ;													
contraction of spindle fibres	anaphase ;													
organisation of chromosomes at the equator	metaphase													

Question	Answer	Marks
6(b)	I antibodies have receptors <i>any three from</i> 1 clones / AW, of B/T cells consist of very few cells ; AW = types of / specific / groups of 2 (mitosis occurs) during <u>clonal expansion</u> ; 3 many, B cells / plasma cells, to make antibodies ; 4 many, T-helper / T_h, cells to secrete, cytokines / lymphokines / interleukins ; 5 many, T-killer cells / T-cytotoxic cells, to kill, infected cells / cancer cells ; 6 memory cells are produced for, secondary response / faster response (when antigen encountered again) ; A during a secondary infection (implying of the same pathogen) 7 need to make genetically identical cells ;	3