

Conservation Governance and Rural Livelihoods in Madagascar

Long-Run Evidence from Rural Observatory Surveys (1995–2025)

**Florent Bédécarrats, Véromanitra
Razafimahenina**

Contents

1	Overview	9
1.1	Abstract	9
1.2	Structure	10
I	Analysis	
2	Conservation in Madagascar	13
3	Madagascar's protected area expansion	15
4	The Rural Observatory Surveys	17
5	Research question	19
6	Data	21
7	Survey data	23
8	Georeferencing	25
9	Protected area overlaps	27
10	Outcome variable	29
11	Descriptive statistics	31
12	Identification Strategy	33
13	Two natural experiments	35
13.1	Marovoay and Ankarafantsika National Park (2003)	35
13.1.1	The 2002 park extension	35
13.1.2	The Betsiboka River as a natural barrier	35
13.1.3	Behavioural validation from the 2025 resurvey	37
13.2	Lac Alaotra and the Protected Landscape (2008)	38
13.2.1	A different conservation model	38
13.2.2	Why all Alaotra communities are exposed	38
13.2.3	Behavioural validation from the 2025 resurvey	40
13.3	PA zone usage intensity across both sites	40
13.4	Extensions: three additional cases (2006–2007)	41
14	From households to villages	43
14.1	The rotating panel constraint	43

14.2	Village-level aggregation	43
14.3	Outcome variable	43
15	Building the counterfactual	45
15.1	Selecting donor communities	45
15.2	Generalized synthetic control	45
15.2.1	Intuition	45
15.2.2	Formal specification	45
15.2.3	Two separate models	46
15.3	Inference with few exposed units	46
15.4	Complementary household-level designs	46
16	Extensions to three additional observatories	49
17	Looking ahead: full household-level estimation	51
18	Results	53
19	Ankarafantsika: strict conservation (IUCN II)	55
19.1	Panel construction	55
19.2	gsynth ATT path	55
19.2.1	Within-Marovoay household RCS check	55
19.3	Unit-specific effects	56
20	Lac Alaotra: multipurpose conservation (IUCN VI)	57
20.1	Panel construction	57
20.2	gsynth ATT path	57
20.2.1	Within-Alaotra household RCS check	57
21	Comparison	59
22	Long-run assessment: 2025 resurvey	61
22.1	Panel construction (extended model)	61
22.2	Extended gsynth	61
22.3	Robustness of the 2025 gap	62
22.4	Alaotra long-run assessment	62
23	Extensions	63
24	Study sites	65
24.1	Farafangana and the Corridor Forestier Ambositra-Vondrozo	65
24.2	Toliara North and the coastal PA cascade	66
24.3	Fénérive East and the Réserve de Tampolo	67
24.4	Design summary	69
25	Results by observatory	71
25.1	Ankarafantsika (clean donors)	71

25.2	Lac Alaotra (clean donors)	71
25.3	Farafangana: Corridor Forestier effect	72
25.4	Toliara North: coastal PA cascade	73
25.5	Fénérive East: low-intensity reserve	74
26	Five-case synthesis	75
26.1	Comparative results	75
26.2	Three patterns	76
26.3	Caveats	77
27	Discussion	79
28	Summary of findings	81
29	Policy implications	83
29.1	Conservation governance matters	83
29.2	Compensation and livelihood integration	83
29.3	Limitations of the policy inference	83
30	Counterfactual validity	85
30.1	Marovoay: alternative explanations	85
30.2	Alaotra: single-unit identification	85
30.3	Implications for interpretation	85
31	Limitations	87
32	Next steps: the 2026 survey wave	89
32.1	Staggered DiD design	89
32.2	Planned analyses	89
	References	91

II

Appendices

33	Appendix A: Data Pipeline	95
34	Georeferencing ROS localities	97
34.1	The problem	97
34.2	Reference datasets	97
34.3	String cleaning	97
34.4	Hierarchical fuzzy matching	97
34.5	Match rates	98
35	Protected-area overlap analysis	99
35.1	Spatial overlaps	99
35.2	ROS survey inventory	100

35.3	Temporal overlap logic	102
35.4	Decree date revision	102
35.5	Priority pairs	103
36	Variable construction	105
36.1	Georeferencing households	105
36.2	Household composition	105
36.3	Education	105
36.4	Agricultural activities	105
36.5	Food security	105
36.6	Income construction	105
36.7	Consolidation	106
37	Appendix B: Donor Pool Validity	107
38	Why static PA boundaries are insufficient	109
39	Dynamic WDPA construction	111
40	Proximity analysis	113
40.1	Method	113
41	Results	115
41.1	Exposure classification	115
41.2	Detailed exposure profiles	118
42	Implications	119
42.1	For the primary analysis	119
42.2	For the extended analysis	119
43	Appendix C: Robustness Tests	121
44	Unadjusted log income	123
45	Income in levels	125
46	Synthetic Difference-in-Differences	127
46.1	Ankarafantsika (SDID)	127
46.2	Alaotra (SDID)	128
47	Callaway–Sant’Anna DiD with repeated cross-sections	129
47.1	Panel construction	129
47.2	Group-time ATT	129
47.3	Event-study aggregation	130
47.4	Cohort-specific ATT	130
48	Summary of robustness	133

49	Appendix D: Methodology	135
50	Generalised synthetic control	137
50.1	Interactive fixed-effects model	137
50.2	Estimation	137
50.3	Cross-validation for factor selection	137
50.4	Inference	137
50.5	Relationship to standard synthetic control	137
51	Callaway–Sant’Anna staggered DiD	139
51.1	Group-time ATT framework	139
51.2	Repeated cross-sections	139
51.3	Aggregation	139
51.4	Implementation (planned for 2026 data)	139
52	Synthetic Difference-in-Differences	141
52.1	Framework	141
52.2	Comparison with gsynth	141
52.3	Inference	141
53	Inference with few clusters	143
53.1	The power problem	143
53.2	Practical implications	143
	References	145
	Bibliography	147



*Marovoay observatory site near Ankarafantsika National Park.
Photo: Véromanitra Ramizason.*

1. Overview

i Living document

This website is a companion to an academic paper in preparation. It will be updated when data from the 2026 survey round becomes available. All code is available in the [GitHub repository](#).



Figure 1.1: Marovoay, Véromanitra Ramizason

1.1 Abstract

Does the governance model of a protected area matter for the livelihoods of surrounding communities? We exploit the staggered creation and extension of protected areas in Madagascar between 2002 and 2008, combined with two decades of household survey data from the Rural Observatory System (ROS, 1995–2015) and a 2025 resurvey, to estimate the causal impacts of conservation on household income.

Using generalized synthetic control methods with interactive fixed effects — a technique that constructs data-driven counterfactuals for each exposed unit by modelling unobserved common factors — we compare two governance archetypes:

- **Strict conservation** (IUCN category II): the 2002 extension of Ankarafantsika National Park, monitored via the Marovoay observatory.
- **Multipurpose conservation** (IUCN category VI): the 2008 creation of the Lac Alaotra new protected area, monitored via the Alaotra observatory.

We find a persistent negative effect of strict conservation at Ankarafantsika on equivalised household income ($ATT \approx -0.312$ log points), while multipurpose conservation at Alaotra shows no detectable effect ($ATT \approx +0.014$ log points). These results extend to three additional observatory–PA pairs. A 2025 resurvey confirms the long-run persistence of these patterns.

1.2 Structure

The analysis chapters present the core argument:

1. **Context** — Conservation policy in Madagascar and the research question.
2. **Data** — The Rural Observatory Surveys, georeferencing, and variable construction.
3. **Identification Strategy** — Exposure definitions, donor pool selection, and the generalized synthetic control method.
4. **Results** — Comparing strict vs. multipurpose conservation impacts on household income.
5. **Extensions** — Three additional observatories near protected areas created during 2006–2007.
6. **Discussion** — Policy implications, limitations, and the roadmap for the 2026 survey wave.

The appendices provide full methodological and data details:

- **A. Data Pipeline** — Reproducible georeferencing, PA overlap analysis, and variable harmonisation.
- **B. Donor Pool Validity** — Spatial audit of PA exposure across all donor observatories.
- **C. Robustness Tests** — Alternative specifications, SDID, and sensitivity analysis.
- **D. Methodology** — Technical details on gsynth, Callaway–Sant’Anna DiD, and interactive fixed effects.



18	Results	53
19	Ankarafantsika: strict conservation (IUCN II)	55
19.1	Panel construction	55
19.2	gsynth ATT path	55
19.3	Unit-specific effects	56
20	Lac Alaotra: multipurpose conservation (IUCN VI)	57
20.1	Panel construction	57
20.2	gsynth ATT path	57
21	Comparison	59
22	Long-run assessment: 2025 resurvey	61
22.1	Panel construction (extended model)	61
22.2	Extended gsynth	61
22.3	Robustness of the 2025 gap	62
22.4	Alaotra long-run assessment	62
23	Extensions	63
24	Study sites	65
24.1	Farafangana and the Corridor Forestier Ambositra-Vondrozo	65
24.2	Toliara North and the coastal PA cascade	66
24.3	Fénérive East and the Réserve de Tampolo	67
24.4	Design summary	69
25	Results by observatory	71
25.1	Ankarafantsika (clean donors)	71
25.2	Lac Alaotra (clean donors)	71
25.3	Farafangana: Corridor Forestier effect	72
25.4	Toliara North: coastal PA cascade	73
25.5	Fénérive East: low-intensity reserve	74
26	Five-case synthesis	75
26.1	Comparative results	75
26.2	Three patterns	76
26.3	Caveats	77
27	Discussion	79
28	Summary of findings	81
29	Policy implications	83
29.1	Conservation governance matters	83
29.2	Compensation and livelihood integration	83
29.3	Limitations of the policy inference	83
30	Counterfactual validity	85
30.1	Marovoay: alternative explanations	85
30.2	Alaotra: single-unit identification	85
30.3	Implications for interpretation	85
31	Limitations	87
32	Next steps: the 2026 survey wave	89
32.1	Staggered DiD design	89
32.2	Planned analyses	89
32.2	References	91

2. Conservation in Madagascar

Protected areas, governance models, and rural livelihoods

3. Madagascar's protected area expansion

The international community has committed to protecting 30% of global land and sea by 2030 under the Kunming-Montréal Global Biodiversity Framework. For biodiversity-rich countries like Madagascar, this target translates into a sustained impulsion to expand and strengthen protected area (PA) networks. Madagascar responded early: at the 2003 Durban World Parks Congress, President Ravalomanana announced the *Durban Vision* – a commitment to triple the country's PA surface from 1.7 million to 6 million hectares [1], that is, from about 3% to about 10% of the national territory. By 2015, the *Système des Aires Protégées de Madagascar* (SAPM) had grown from 47 to over 160 protected areas, covering roughly 12% of the national territory.

This expansion was not monolithic. The SAPM introduced a diversity of governance models, ranging from strictly protected national parks under Madagascar National Parks (MNP) management to multipurpose landscapes co-managed by international NGOs and local communities through *transferts de gestion*. The IUCN classification system captures this spectrum: Category II parks prioritise ecosystem integrity with limited human use, while Categories V and VI allow sustainable resource extraction and community participation.

The welfare consequences of this expansion for adjacent rural communities remain contested. A growing quasi-experimental literature finds heterogeneous effects of PAs worldwide: some studies document negative impacts on local incomes and consumption [2], [3], while others find neutral or positive effects, particularly when PAs generate tourism revenues or ecosystem services [4]. There is ground to assume that the modest positive effects observed elsewhere might not materialise in Madagascar, as extreme poverty prevalence intensifies dependence of local population on natural resources, tourism remains an order of magnitude smaller than in other biodiversity hotspots, and weak governance might undermine stakeholder engagement and benefit-sharing mechanisms. In Madagascar specifically, evidence remains thin and largely cross-sectional [5]. The few causal studies focus on deforestation outcomes rather than household livelihoods.

A central question in this debate is whether the governance model of a protected area matters for the people living around it. Strict exclusionary conservation and participatory community-based management represent fundamentally different arrangements with local populations. Yet most empirical studies treat PAs as homogeneous interventions, comparing “protected” with “unprotected” rather than examining how different institutional arrangements mediate livelihood impacts.




Figure 3.1: Cumulative protected area coverage in Madagascar by IUCN category. Data from the dynamic WDPA layer; each PA enters at its earliest legal creation date. The post-Durban surge (after 2003) is dominated by Categories V and VI (multipurpose).

4. The Rural Observatory Surveys

This study exploits a rare longitudinal dataset: the Rural Observatory System (ROS), a network of rural observatories conducted across Madagascar between 1995 and 2015. Coordinated by INSTAT then the Ministry of Agriculture, with technical support from IRD and funding from international donors, the ROS surveyed approximately 500 households per observatory per year across up to 28 observatory sites spanning the country's diverse agroecological zones — from the humid eastern lowlands to the semi-arid south [6].




Figure 4.1: Coarse location of rural observatory and survey years

The ROS operates as a rotating panel: each observatory tracks a fixed set of households, but approximately 15% leave the sample each year through death, migration, or attrition, with replacement drawn from the original sampling frame. Over a decade, cumulative attrition reaches roughly 80% of the initial cohort. While this design limits individual-level panel analysis, it preserves cross-sectional representativeness at each wave: village-year aggregates consistently estimate the same population quantities, making the data well-suited for site-level quasi-experimental methods.

Critically, several rural observatories are located adjacent to protected areas that were extended or created during the survey period (2002–2008). This overlap between a two-decade household survey and staggered conservation interventions provides a natural laboratory for impact evaluation — one that has not been systematically exploited until now.

A 2025 resurvey of two key sites (Marovoay and Alaotra) adds a long-run perspective, enabling us to assess whether short-run effects persist more than two decades after exposure onset. A further 2026 wave is planned to extend coverage to additional observatories.

5. Research question

We ask: **does the governance model of a protected area affect household income in surrounding communities?**

We compare two natural experiments embedded in the ROS data:

1. **Strict conservation:** The 2002 extension of Ankarafantsika National Park (IUCN Category II), managed by MNP. The Marovoay observatory straddles the park boundary, with the Betsiboka River creating a natural exposure–control divide between east-bank villages (adjacent to the park) and west-bank villages (effectively separated by the river).
2. **Multipurpose conservation:** The creation of the Lac Alaotra new protected area (IUCN Category VI), managed by Durrell Wildlife Conservation Trust through community-based natural resource management, with effective enforcement from approximately 2008. The entire Alaotra observatory shares dependence on the lake-marsh ecosystem.

Using generalized synthetic control methods [7] with an extended donor pool of up to nine additional observatories, we estimate the average treatment effect on the treated (ATT) for equivalised household income. We then extend the analysis to three additional observatory–PA pairs (Farafangana, Toliara North, Fénérive East) where PAs were created in 2006–2007, exploiting the staggered timing to strengthen identification.

The remainder of this document is structured as follows. [Chapter 2](#) describes the data sources and variable construction. [Chapter 3](#) details the identification strategy and estimation methods. [Chapter 4](#) presents the main results for Ankarafantsika and Alaotra. [Chapter 5](#) extends the analysis to three additional sites. [Chapter 6](#) discusses policy implications and outlines plans for the 2026 survey wave.

6. Data

Rural Observatory surveys, georeferencing, and variable construction

This chapter summarises the data construction process. The full reproducible workflow — georeferencing, PA overlap analysis, and variable harmonisation — is documented in [Appendix A](#). The underlying survey data are described in a companion data paper [\[8\]](#); supplementary materials and replication code are available in the [associated repository](#).

7. Survey data

The Rural Observatory System (ROS) surveyed rural households across Madagascar from 1995 to 2015. Many observatories remained active for only a few years. We use 11 observatories that were active by 1999-2000 and had at least 6 years of survey data. This dataset comprises approximately 101,253 household-year observations. Each observatory tracks ~500 households per year in a rotating panel design, with random replacement of attrited households preserving cross-sectional representativeness.

Rural Observatories in the Analysis




Figure 7.1: Location of the 11 rural observatories used in this study.




Figure 7.2: Year-by-year data coverage across observatories and survey sites.

8. Georeferencing

The original survey data lacked systematic georeferencing. Locality names were recorded as free text, with inconsistent spellings across years and languages (Malagasy and French). We linked each observation to official administrative boundaries using the Common Operational Datasets (COD), maintained by OCHA and Madagascar's BNGRC. The full georeferencing procedure is described in the [companion data documentation](#).

The matching procedure is hierarchical:

1. **String normalisation:** lowercase, remove diacritics and qualifiers (*centre*, *haut*, *bas*), standardise bilingual toponyms, convert Roman to Arabic numerals.
2. **Fuzzy matching:** Jaro-Winkler distance against the COD gazetteer, filtered by observatory-level district membership, with a maximum distance threshold of 0.25.
3. **Cascading spatial levels:** match first at the commune (ADM3) level, then refine to fokontany (ADM4) where possible.

The result is a geocoded correspondence table linking each ROS household identifier (j5) to its P-coded commune and fokontany. This enables spatial overlay with protected area boundaries.

9. Protected area overlaps

Standard PA databases — including the current WDPa and the Vahatra atlas — record only the most recent boundary and designation of each protected area. They do not capture boundary changes (e.g., Ankarafantsika’s 2002 extension roughly doubled the effectively protected area), temporary protection decrees that imposed de facto restrictions years before formal designation, or date errors (many WDPa entries carry incorrect `STATUS_YR` values).

We instead use a dynamic WDPa for Madagascar — a temporally explicit database that assigns each PA state a precise valid-from / valid-to interval derived from the Madagascar CNLEGIS legal texts database and historical versions of the SAPM geospatial database [**INCLUDE REFERENCE DATA PAPER**]. This enables us to determine whether an observatory was adjacent to an *active* PA at any given point during the study period (see [Appendix B](#) for full construction details).

For each of the 11 study observatories, we compute the minimum distance from commune centroids (projected to UTM 38S) to the nearest PA boundary active during 1999–2014. All (commune, PA) pairs within 20 km are flagged. Three temporal configurations emerge:

Table 9.1: Overview of observatory–PA temporal relationships.

Configuration	Observatories	Use
PA predates ROS surveys (no baseline)	e.g., Ranomafana, Andringitra	Excluded
PA created <i>during</i> ROS (1999–2014)	Marovoay (2003), Alaotra (2008)	Primary treatment
PA created during ROS but later	Farafangana, Toliara, Fénériver (2006–07)	Extension treatment (Ch. 5)




Figure 9.1: Survey coverage windows and PA exposure events for the 11 study observatories. Bars show the survey span; triangles mark the nearest new PA decree date. Green bars indicate observatories with sufficient pre/post data for gsynth; red bars indicate insufficient coverage relative to the PA event.

Figure 9.1 shows that although the ROS was designed for agricultural monitoring, not conservation impact evaluation, some PAs were created relatively near from observatory sites. Gsynth requires sufficient pre-treatment years to learn unit-specific factor loadings, and sufficient post-treatment years for the counterfactual to be informative. We apply a minimum threshold of 4 pre-exposure and 3 post-exposure survey years.

The two primary pairs — **Marovoay / Ankarafantsika** (strict, IUCN II, 2003 extension) and **Alaotra / Lac Alaotra** (multipurpose, IUCN V, 2007 temporary decree) — offer the longest pre- and post-exposure windows. Three additional pairs — Farafangana, Toliara North, and Fénérive East — provide shorter but informative extensions ([Chapter 5](#)). Observatories without a nearby PA (shown in blue) serve as the donor pool. A systematic spatial audit of donor pool validity is presented in [Appendix B](#).

10. Outcome variable

The primary outcome is log equivalised household income. Construction proceeds as follows:

1. **Total income** (*revtot*): the sum of current income components: rice revenue (*rev_riz*), other crop revenue (*rev_cu*), livestock revenue (*revel*), fishing revenue (*revpeche*), principal activity income (*revppal*), and secondary activity income (*revsec*). Missing components are set to zero; the total is imputed from available sub-components when *revtot* itself is missing.
2. **Equivalisation**: using the OECD-modified scale: $1 + 0.5 \times (\text{additional adults}) + 0.3 \times (\text{children under 14})$, computed from individual-level demographic records [9].
3. **Winsorisation**: at the 1st and 99th percentiles within each year to limit the influence of extreme values.
4. **Log transformation**: $y_{it} = \log(\text{equivalised income}_{it} + 1)$.

Cross-year harmonisation is not trivial: variable names, questionnaire modules, and income definitions changed across survey waves. The full details of these adjustments are in [Appendix A](#).




Figure 10.1: Mean log equivalised household income over time, by observatory.

11. Descriptive statistics

Table 11.1: Summary statistics for the analysis sample.

Sample Descriptive Statistics

Analysis period: 1999–2014

group	N house- hold- years	Mean income (Ar)	Mean equiv. in- come	Mean HH size	Mean educ. years (adults)	% liter- ate adult
Marovoay exposed (east)	3873	1,736,175	700,533	5.0	4.4	92.0
Marovoay control (west)	3617	2,135,445	833,960	5.4	4.5	91.7
Alao- tra (ex- posed)	7613	1,893,377	691,554	5.1	4.5	95.6
Donor pool	48052	1,192,493	453,894	5.5	4.3	80.1

12. Identification Strategy

How we estimate the causal effect of protected areas on household income

The central question of this study is whether the creation or expansion of protected areas caused changes in household income, or whether observed income differences merely reflect pre-existing trends. This chapter explains how we attempt to isolate causal effects.

The strategy has four parts. First, we define two natural experiments to propose plausible exposure-control comparisons (Chapter 13.). Second, we explain how the survey design constrains the unit of analysis (Chapter 14.). Third, we describe the statistical method used to construct counterfactual income trajectories (Chapter 15.). Fourth, we briefly introduce three additional cases that extend the analysis (Chapter 16.).

13. Two natural experiments

We exploit two events during the ROS survey period where a new or expanded PA began to impose restrictions on communities that had previously been surveyed. The two cases differ in their conservation governance model – strict exclusion at Ankarafantsika versus participatory management at Lac Alaotra – providing a built-in comparison of how governance type mediates income effects.

13.1 Marovoay and Ankarafantsika National Park (2003)

13.1.1 The 2002 park extension

As illustrated in Figure 13.1, the Ankarafantsika National Park was formally extended by Decree 2002-798 on 7 August 2002, merging the former Réserve Naturelle Intégrale (60,500 ha, created in 1927) with a forest reserve into a 136,500 ha national park under IUCN Category II (strict nature conservation).

We set the exposure year to 2003, which is the first full ROS survey year following the decree. Decree 2002-798 was signed on 7 August 2002. Because the ROS survey is conducted annually with a reference period from October of year-1 to September of the survey year and we attribute any effect to the first full survey year following the decree, we set the exposure year to 2003. We also test whether using 2002 instead of 2003 modifies the estimated result and find negligible incidence.

According to the concordant grey literature we reviewed, we consider the 2002 extension was the first event to impose binding land-use restrictions on the communities adjacent to the forest reserve zone. The earlier designations had very different *de facto* implications:

- The 1927 *Réserve Naturelle Intégrale* covered the inland plateau to the north-east, geographically separate from the Marovoay communities. The east-bank *fokontany* had no direct interface with the strictly protected RNI.
- The 1929 forest reserve was not a conservation zone but a timber and resource reserve under the forest administration. Analyses of this period found no effective enforcement: subsistence agriculture, cattle grazing, and charcoal production continued unimpeded [10].

The 2002 reclassification merged these into a single national park where slash-and-burn agriculture (*tavy*), charcoal production, and non-timber forest product extraction were explicitly prohibited and, crucially, enforced. Enforcement capacity was supported by international conservation funding from the World Bank and bilateral donors as part of Madagascar's third Environmental Action Plan. Figure 13.1 maps the spatial extent of the consolidation using dynamic WDPA boundary data: the original RNI (dark green) covered the inland plateau to the north-east, while the 2002 extension (light green) more than doubled the protected area to encompass forest reserves abutting the Marovoay communities [11].

13.1.2 The Betsiboka River as a natural barrier

The Marovoay observatory comprises four *fokontany* (villages) split by the Betsiboka River – a major, permanently flowing watercourse. Crossing is possible only under certain hydrological conditions, often requiring long detours or costly boat transport. In practice, flows of people, goods, and services run along the Marovoay downstream axis rather than across the river.

This geographic feature creates a natural experiment: east-bank communities have direct walking access to the park, while west-bank communities face a substantial access penalty.

Table 13.1: Marovoay site exposure assignments. Distance to RNI = straight-line distance to the pre-2002 *Réserve Naturelle Intégrale*; Distance to NP = distance to the post-2002 national park boundary. Effective distance adds a 15 km river-crossing penalty for west-bank fokontany.

Fokontany	Bank	Dist. to RNI (pre-2002)	Dist. to NP (post-2002)	Role
Bepako	East	33 km	5.2 km	Exposed (from 2003)
Madiromion-gana	East	25.1 km	11.7 km	Exposed (from 2003)
Ampijoroa	West	51.2 km (+ river crossing)	11.3 km (+ river crossing)	Donor (control)
Maroala	West	44.7 km (+ river crossing)	5.6 km (+ river crossing)	Donor (control)

Effective distance adds a 15 km river-crossing penalty for west-bank fokontany.

The 15 km river-crossing penalty is conservative. In rural Madagascar, where most travel is on foot, the actual additional travel time likely exceeds the implied 3-hour walk. An effective distance threshold of 12 km — corresponding to approximately a 2.5-hour daily commuting limit — places both east-bank villages within the PA’s zone of influence and both west-bank villages outside it, cleanly separating “exposed” from “donor” communities.




Figure 13.1: Marovoay observatory and the 2002 Ankarafantsika expansion. Dark green: the original 1927 *Réserve Naturelle Intégrale* (~571 km²). Light green: the 2002 national park extension (~1,351 km²). The Betsiboka River separates west-bank fokontany (donors, grey) from east-bank fokontany (exposed, orange). Labels show distance to NP boundary. Water features from OpenStreetMap.

13.1.3 Behavioural validation from the 2025 resurvey

The exposure assignment rests on the assumption that the Betsiboka River effectively separates communities into “exposed” and “not exposed” groups. The 2025 resurvey provides a direct test of this assumption through its dedicated protected area usage module (UM), which asked all 1,026 households whether they visit the PA zone, what activities they pursue there, and why non-visitors stay away.

Table 13.2: PA interaction at Marovoay by fokontany (2025 resurvey). ‘Visit PA zone’ = any visit, including leisure. ‘Extractive use’ = food gathering, woodcutting, cultivation, grazing, or craft materials. ‘% cite forbidden’ = share of non-visitors citing legal prohibition.

PA Interaction by Fokontany						
Marovoay observatory, 2025 resurvey (N = 519 households)						
Site	Bank	N	% visit PA zone	% extractive use	% cite 'forbid-den'	
Bepako	East	123	49	11	19	
Madiromion-gana	East	137	28	9	36	
Ampijoroa	West	143	15	1	11	
Maroala	West	116	16	2	2	

Three findings support the exposure assignment:

1. **Visitation asymmetry:** 38% of east-bank households report visiting the park zone versus 15% on the west bank. However, among visitors on both banks, the dominant activity is leisure walks (67%); only 10% of east-bank households report extractive use (food gathering, woodcutting). 91% of all visitors go only a few times per year.
2. **Awareness of prohibition:** At Madiromiongana – the east-bank fokontany closest to the park’s core – 34% of non-visitors cite legal prohibition as their reason for not going, versus 2% at Maroala on the west bank. The park’s legal status is a binding constraint on the east bank, not the west.
3. **Distance as the dominant barrier:** Across both banks, the most commonly cited reason for not visiting is distance (*lavitra* in Malagasy), mentioned by 54% of non-visitors. The river makes the park *behaviourally distant* for west-bank communities even where it is *geographically close*.

13.2 Lac Alaotra and the Protected Landscape (2008)

13.2.1 A different conservation model

Lac Alaotra is Madagascar’s largest lake and a critical habitat for the Alaotran gentle lemur (*Haplemur alaotrensis*). The Ramsar Convention listed the site in 2003. In 2015, the lake and its marshlands were formally designated as a *Paysage Harmonieux Protégé* (PHP, 425 km², IUCN Category V – a multipurpose landscape where sustainable use is permitted). However, enforcement through community-based natural resource management (CBNRM) agreements and *transferts de gestion* had intensified from around 2008 under the management of Durrell Wildlife Conservation Trust. A temporary protection decree (*Arrêté* 381/2007, 1 August 2007) preceded the intensification. We set the exposure year to 2008, consistent with the annual-survey convention applied to Ankarafantsika.¹

The key difference from Ankarafantsika is governance: rather than exclusionary enforcement prohibiting all resource extraction, Alaotra’s CBNRM framework regulates lake-marsh use through community-managed access rules, seasonal fishing closures, and habitat restoration agreements. This participatory model is designed to accommodate livelihoods rather than exclude them.

13.2.2 Why all Alaotra communities are exposed

Unlike Marovoay, there is no natural barrier separating “exposed” from “unexposed” communities at Alaotra. The observatory’s seven *fokontany* (across three historical survey sites) all display modest but not negligible interactions with the lake-marsh ecosystem, with no clear distinctions among themselves

¹An alternative exposure year of 2007 is defensible; using 2007 instead has negligible impact on the estimated ATT, given that the post-exposure window contains up to seven observations under either assumption.

in this regard. The co-managed enforcement mechanism — seasonal closures, gear restrictions, marsh-burning bans — also applies to all lake-adjacent communities similarly.

A complication specific to Alaotra is the proximity of some eastern hamlets to two additional protected areas: the Corridor Ankeniheny-Zahamena (CAZ, IUCN Category VI, managed by Conservation International) and the Zahamena National Park (IUCN Category II), which lies north of the CAZ. The hamlet of Mangabe lies closer to the CAZ boundary (~14 km) than to the Lac Alaotra PHP, creating dual-PA exposure.

We therefore treat the entire observatory as a single exposed unit, aggregated to a site-level mean. This is a stronger assumption than the site-level assignment at Marovoay, but it is supported by the uniform enforcement mechanism (CBNRM applies to all lake-adjacent communities) and the behavioural data from the 2025 resurvey discussed below.




Figure 13.2: Alaotra observatory and surrounding protected areas. Lac Alaotra PHP (green), Corridor Ankeniheny-Zahamena (CAZ, orange), Zahamena National Park (light orange). Fokontany centroids from 2025 GPS survey; distances to nearest PA boundary.

13.2.3 Behavioural validation from the 2025 resurvey

Table 13.3: PA interaction at Alaotra by fokontany (2025 resurvey). ‘Visit PA zone’ includes any visit to the lake-marsh area. ‘Extractive use’ = food gathering, fishing/hunting, cultivation, grazing, or woodcutting within the zone. Fishing/hunting shown separately as it dominates extractive use.

PA Interaction by Fokontany				
Alaotra observatory, 2025 resurvey (N = 507 households)				
Site	N	% visit zone	% extractive	% fish/hunt
Ambatomanga	68	51	26	21
Ambodivoara	58	45	21	10
Ambohidrony	52	54	21	13
Ambatoharanana	31	45	13	10
Analamiranga	46	41	13	4
Mangabe	73	47	8	5
Avaradrano	90	38	7	0
Feramanga Atsimo	89	37	3	0

The Alaotra data reveal an important distinction between *visiting* and *depending on* the PA zone:

- **Visiting is common but mostly non-extractive:** 37–54% of households report visiting the lake-marsh zone, but the survey question captures any visit including leisure walks. Among visitors, 78% go only a few times per year.
- **Extractive use is concentrated and heterogeneous:** Fishing and hunting — the dominant extractive activity, reflecting traditional lake use — involves 21% of households in Ambatomanga but none in Avaradrano or Feramanga Atsimo. The overall extractive rate is 13% of all households.
- **Distance, not prohibition, explains non-use:** Among Alaotra non-visitors, the most cited reason is distance (*lavitra*), at 35%. Legal prohibition is cited by only 8%, consistent with the less restrictive multipurpose framework.

The relatively uniform visiting rate but sharply differentiated*extractive rate suggests that proximity to the ecosystem — rather than engagement with PA-regulated resources — drives the visiting pattern.

13.3 PA zone usage intensity across both sites

Figure 13.3 provides a cross-site comparison of how intensively households use the PA zone. The contrast between the two observatories is stark: at Marovoay, the majority of households never visit the park (white bars), with the river barrier clearly separating high-access sites (Bepako) from low-access ones (Ampijoroa, Maroala). At Alaotra, a larger share of households visit the zone, but almost all do so infrequently (grey bars). Regular visitors — monthly or more, shown in warm colours — are a thin sliver confined to the closest sites.




Figure 13.3: PA zone visit frequency by site (2025 resurvey). Each bar represents 100% of households in a site. White = does not visit; grey = few times per year; warm colours = monthly or more. Sites ordered by total share of visitors.

13.4 Extensions: three additional cases (2006–2007)

The donor-pool audit (Section 15.1) reveals that four donor observatories are themselves adjacent to PAs created during the study period. Three of these provide usable exposure variation — Farafangana (2006), Toliara North (2007), and Fenérive East (2007) — and are analysed as extension cases in Chapter 16. and Chapter 5.

14. From households to villages

14.1 The rotating panel constraint

The ROS can be interpreted as a rotating panel, instead of a strict re-interview panel. This is a critical distinction for estimation. Each observatory tracks approximately 500 households per year, but approximately 15% leave the sample each year through death, migration, or attrition. Departing households are replaced by newly sampled ones following the original random sampling design. Over a 10-year horizon, cumulative attrition reaches roughly 80% of the initial cohort [12].

This design means that the standard econometric approach — following the same households over time and using household fixed effects to control for unobserved heterogeneity — is not feasible. Only about 44% of households surveyed in 1998 remain in 2003, and fewer than 6% survive to 2014. Furthermore, attrition is non-random: the dominant cause is out-migration (65% of traceable attritors), which correlates with income and shock exposure. The 2025 resurvey uses entirely new household identifiers that carry no continuity with the 1995–2014 series.

The rotating design does, however, preserve cross-sectional representativeness at each survey wave: because replacement households are randomly drawn from the same villages, village-year statistics consistently estimate the same population quantity across years, even as the composition of surveyed households changes.

What this means in practice

A household-level panel regression with household fixed effects would produce misleading results here: it would estimate effects only for the shrinking minority of households that remain in the sample for multiple years — a selected, non-representative sub-population. Village-level aggregation avoids this problem entirely.

14.2 Village-level aggregation

We therefore analyse the data at the village (*fokontany*) level, aggregating household-level income to village-year medians. This addresses both the attrition problem and the estimation requirements:

1. **Cross-sectional representativeness is preserved:** Random replacement of attrited households ensures that village-year medians estimate the same population parameter across waves, even though the specific households change.
2. **The estimator requires a balanced panel:** The generalized synthetic control method (described below) requires a complete unit-by-year matrix. Village-level aggregation produces this, while household-level data would have pervasive gaps.
3. **Exposure is assigned at the village level:** PA exposure depends on the village's distance to the park boundary, not on individual household characteristics. Village-level analysis matches the unit of exposure assignment to the unit of analysis.

The median is preferred to the mean because rural income distributions are right-skewed with occasional extreme values. Village-level medians are more robust to the specific composition of households surveyed in any given year.

14.3 Outcome variable

The primary outcome is *log equivalised household income*. Construction proceeds in four steps:

1. **Total income (*revtot*):** the sum of current income components: rice revenue, other crop revenue, livestock revenue, fishing revenue, principal activity income, and secondary activity income. Missing components are set to zero.

2. **Equivalisation:** using the OECD-modified scale to adjust for household composition:

$$\text{equiv} = 1 + 0.5 \times (\text{additional adults}) + 0.3 \times (\text{children under 14}) \quad (14.1)$$

This ensures that a large household with higher total income is not misleadingly compared to a small household with lower total but similar per-person income.

3. **Winsorisation:** at the 1st and 99th percentiles within each year to limit the influence of extreme values.
4. **Log transformation:** $y_{it} = \log(\text{equivalised income}_{it} + 1)$, which compresses the right tail and allows treatment effects to be interpreted as approximate percentage changes.

Cross-year harmonisation is non-trivial: variable names, questionnaire modules, and income definitions changed across survey waves. The full details of these adjustments are documented in [Appendix A](#).

15. Building the counterfactual

The fundamental challenge is that we observe what happened to communities near the PA, but not what *would have happened* to them without the PA. The statistical method must therefore construct a credible counterfactual — an estimate of the income trajectory these communities would have followed had the PA never been created.

15.1 Selecting donor communities

The counterfactual is built from *donor pools* — rural observatory villages that were not exposed to any PA during the study period. We start with 9 donor observatories providing approximately 60 donor sites across the country.

A systematic spatial audit ([Appendix B](#)) checks whether these donors are themselves adjacent to PAs created during 1999–2014. Using a dynamic WDPA layer that records when each PA became legally operative, we flag any donor commune within 20 km of an active PA boundary as also exposed:

Table 15.1: Donor pool PA exposure classification.

Status	Observatories	Reason
“Clean”	Antsirabe (02), Tsiroanomandidy (13), Ambovombe (16), Mahanoro (25), Itasy (41)	No active PA within 20 km during 1999–2014
Also exposed	Antsohihy (12), Farafangana (15), Toliara North (23), Fénérive East (24)	PA created within 20 km during study period

The main results use the 5 “clean” donor observatories; the [extension analysis](#) confirms that including the 4 also-exposed donors does not materially change the estimates. In the extension analysis ([Chapter 5](#)), the four also-exposed observatories become *exposed units* with their own PA exposure events.

15.2 Generalized synthetic control

15.2.1 Intuition

Standard difference-in-differences (DiD) assumes that exposed and control communities would have followed parallel trends in the absence of the PA — that is, that income would have evolved at the same rate in PA-adjacent and distant villages. This is a strong assumption: rural villages may diverge over time for reasons unrelated to PAs, such as changing weather patterns, road construction, commodity price shifts, or local governance quality.

The generalized synthetic control method (gsynth) of [Y. Xu \[7\]](#) relaxes this assumption. Instead of requiring parallel trends, it allows each village to follow its own trajectory driven by unobserved factors — the method then *learns* these factors from the pre-treatment data and uses them to project what would have happened after the PA was created.

In plain language: gsynth looks at how the exposed villages’ income moved *before* the PA, identifies which combination of donor villages best reproduces that pattern, and then uses those same donors to predict what the exposed villages’ income would have been *after* the PA, had the PA never existed. The difference between the actual income and this predicted counterfactual is the estimated effect.

15.2.2 Formal specification

For each exposed unit i in post-treatment period t , gsynth models the outcome as:

$$Y_{it} = \alpha_i + \xi_t + \lambda'_i f_t + D_{it} \cdot \delta_{it} + \varepsilon_{it} \quad (15.1)$$

where α_i are unit fixed effects (each village has its own baseline income level), ξ_t are time fixed effects (all villages share common year-to-year shocks like droughts or price changes), $\lambda'_i f_t$ are interactive fixed effects (each village responds differently to unobserved time-varying factors), and D_{it} is the treatment indicator. The interactive term $\lambda'_i f_t$ is what distinguishes gsynth from standard DiD: it allows for heterogeneous trends driven by unobserved factors.

Key implementation choices:

- **Factor selection:** leave-one-out cross-validation over $r \in \{0, 1, 2, 3, 4, 5\}$ determines the number of latent factors.
- **Inference:** parametric bootstrap with 200 replications.
- **Minimum pre-treatment periods:** $\min(T_0) = 3$ to ensure adequate pre-treatment fit.

15.2.3 Two separate models

We estimate two separate models, reflecting the distinct nature of each case:

1. **Marovoay model:** Exposed units = Madiromiongana and Bepako (east bank, from 2003). Donors = all other rural observatory sites, including the Marovoay west-bank sites (Ampijoroa, Maroala). With 2 exposed sites and ~60 donors, gsynth estimates unit-specific counterfactuals for each village.
2. **Alaotra model:** Exposed unit = aggregated Alaotra observatory (from 2008). Donors = all non-Alaotra rural observatory sites, including all four Marovoay *fokontany*. With a single exposed unit, this reduces to a synthetic control with interactive fixed effects.

15.3 Inference with few exposed units

A fundamental constraint of this design is the small number of exposed units: 2 sites for Marovoay, 1 for Alaotra. With so few exposed clusters, standard statistical tests are unreliable regardless of the estimator chosen.

To understand this constraint concretely: for Marovoay, the set of observable units within the observatory comprises 4 sites (2 east-bank, 2 west-bank). The number of ways to assign exactly 2 of these 4 units as “exposed” is $\binom{4}{2} = 6$. Under the sharp null hypothesis of no treatment effect, the **minimum achievable p-value** is therefore $1/6 \approx 0.167$ for a one-sided test. No statistical test based solely on these 4 sites can produce a p-value below 0.167, regardless of how large the observed effect is.

When all three Alaotra sites are included as additional control donors (expanding the permutation pool to 7 units), the allocation count rises to $\binom{7}{2} = 21$ and the minimum achievable p-value falls to $1/21 \approx 0.048$, that is just below the conventional 5% threshold.

We report gsynth bootstrap confidence intervals and p-values throughout, but interpret them in light of this **power ceiling**. The inability to reject the null at conventional thresholds does not imply absence of effect; it may simply reflect the inherent inferential limitations of quasi-experimental designs with few geographic units. Placebo-in-space tests (Appendix C) provide complementary evidence on the uniqueness of estimated effects.

15.4 Complementary household-level designs

The gsynth analysis above operates on village-year aggregates and identifies treatment effects against an outside donor pool of clean observatories. We complement it with two within-observatory household-level repeated cross-section designs that exploit a different identifying contrast: exposure-intensity heterogeneity *inside* each observatory. The two layers of inference are deliberately distinct, and their agreement (or disagreement) is itself informative.

Sampling distinction: Because the ROS is a rotating panel, households are not tracked across waves. The valid household-level estimator is therefore a repeated cross-section (RCS) Callaway–Sant’Anna design [13] with `panel = FALSE`, which treats each wave as an independent cross-section of households within sites and aggregates over the household-level observations through the doubly-robust 2-period moment of P. H. C. Sant’Anna and J. Zhao [14]. Site fixed effects (not household fixed effects) absorb time-invariant unobserved heterogeneity. We run the design on the 1999–2014 waves extended with the 2025 resurvey.

Two exposure-intensity proxies: For each observatory we build a binary treatment indicator from one of two within-observatory contrasts:

- **Geographic exposure:** Distance to the nearest PA boundary, or — at Marovoay — east-bank versus west-bank assignment, as motivated in Section 13.1 by the Betsiboka river barrier.
- **Behavioural exposure:** The 2025 share of households declaring at least one PA-dependent extractive use (Section 13.1.3, Section 13.2.3), with a cut at $\geq 12.5\%$ defining the *high-usage* group.

For Alaotra both proxies are reported (the seven merged fokontany span both a meaningful distance gradient — 8.9 to 16.3 km from the nearest PA — and a meaningful extractive-use gradient — 0.7% to 26.5%). For Marovoay only the geographic proxy is reported, because extractive use is rare on both banks (peak: 11.4% at Bepako; Section 13.1.3) and a 12.5% cut leaves no treated unit.

Table 15.2: Within-observatory exposure-intensity classifications used by the secondary household-level RCS designs. Distance and bank are pre-determined geography; extractive-use share is measured on the 2025 resurvey under the same definition as Section 13.1.3 and Section 13.2.3. The two Alaotra fokontany pairs Ambodivoara/Maritampona and Ambatoharanana/Analamiranga are merged onto their older centroids to align the 1999–2014 frame with the 2025 GPS enumeration.

Secondary DiD: exposure-intensity classifications

Site	Distance / bank	% extractive (2025)	Geographic	Behavioural
Marovoay				
Bepako	east	11.4	treated	—
Madiromion-gana	east	8.8	treated	—
Ampijoroa	west	0.7	control	—
Maroala	west	1.7	control	—
Alaotra				
Ambodivoara-Maritampona	8.9 km	20.7	treated	treated
Ambato-haranana-Analamiranga	9.5 km	13.0	treated	treated
Avaradrano	10.4 km	6.7	treated	control
Mangabe	14.0 km	8.2	control	control
Feramanga At-simo	14.7 km	3.4	control	control
Ambohidrony	14.8 km	21.2	control	treated
Ambatomanga	16.3 km	26.5	control	treated

Inference. Because the cluster count is small ($G = 4$ sites at Marovoay, $G = 5$ merged fokontany at Alaotra), we report two standard errors for every point estimate: the design-based, influence-function SE — valid if the estimand is the ATT on these specific sites conditional on the observed treatment assignment — and the site-clustered SE, valid for super-population inference. The gap between the two is informative about how much identifying variation lives across versus within sites.

Limitations. Four caveats are inescapable and should be kept in mind when reading the results.

1. **No clean within-observatory control:** Every site in either observatory is exposed to the conservation framework to some degree; the “control” group is just the *less exposed* subset.
2. **Distance is Euclidean:** Straight-line distance to the PA boundary is not the same as practical accessibility — terrain, roads and rivers all matter and are only partially captured (the river barrier at Marovoay is captured by the bank cut).
3. **Extractive usage is post-treatment endogenous:** The 2025 extractive share is measured *after* exposure, so a fokontany depressed by enforcement could appear less (or more) extractive today as a consequence of the treatment.
4. **Small G:** With only four to seven sites the cluster-robust SE is wide; the design-based IF SE is tighter but identifies a more restrictive estimand.

For these reasons the secondary designs are explicitly complementary rather than confirmatory of the main gsynth analysis. Where they agree with gsynth (Alaotra), the agreement narrows the range of plausible interpretations; where they disagree (Marovoay), the disagreement diagnoses *which* identifying contrast is doing the work in each estimator. Two further estimators — Synthetic DiD [15] and standard two-way fixed effects — are reported in [Appendix C](#) as additional sensitivity checks against the gsynth specification choice.

16. Extensions to three additional observatories

The two-case comparison above — strict enforcement at Ankarafantsika versus participatory conservation at Lac Alaotra — limits external validity. Three additional observatory–PA pairs identified in the donor-pool audit (Section 15.1) provide shorter but informative extensions:

Table 16.1: Extension observatories.

Observatory	Protected area	IUCN	Year	Governance
Farafangana	Corridor Forestier Am- bositra-Vondrozo (CFAV)	V	2006	Multipurpose highland corridor
Toliara North	Amoron'i Oni- lahy	V	2007	Coastal and dry- forest protections
Fénérive East	Réserve de Tam- polo	IV	2007	Small (~7.3 km ²) research forest

These three observatories move from liabilities in the donor pool to informative exposed units, enabling a five-case comparison of how conservation governance type, enforcement intensity, and PA size shape household-income outcomes. The shorter post-exposure windows (8–9 years versus 12 for Marovoay and 7 for Alaotra) limit precision but broaden the institutional variation. The full analysis is presented in [Chapter 5](#).

17. Looking ahead: full household-level estimation

The secondary RCS designs introduced in Section 15.4 already exploit household-level information within each observatory, but they remain limited by the small number of sites available *inside* a single observatory ($G = 4$ to 7). A richer household-level analysis would pool all five exposed observatories together with the never-exposed donors into a single staggered-treatment design, exploiting four staggered exposure cohorts (Ankarafantsika 2003, Farafangana/Toliara/Fenérive 2006–07, Alaotra 2008, plus never-exposed). The natural estimator is again Callaway–Sant’Anna with `panel = FALSE`, complemented by Sun–Abraham event studies via `fixest::sunab()`. The 2026 ROS wave will increase post-exposure information and is expected to sharpen these estimates further. This forward-looking design is discussed in Chapter 32..

18. Results

Strict vs. multipurpose conservation impacts on household income

19. Ankarafantsika: strict conservation (IUCN II)

19.1 Panel construction

The Marovoay model comprises 2 exposed sites (Bepako and Madiromiongana, east bank) and donor sites from the 5 clean observatories identified in [Appendix B](#) (Antsirabe, Tsiroanomandidy, Ambovombe, Mahanoro, Itasy). The exposure period begins in 2003, providing 4 pre-exposure years (1999–2002) and up to 12 post-exposure years (2003–2014).

19.2 gsynth ATT path




Figure 19.1: Ankarafantsika: gsynth ATT path for east-bank sites relative to exposure onset (2003). Outcome: log OECD-equivalised income. Shaded band: 95% parametric bootstrap CI.

The gsynth estimator finds a persistent negative effect of PA exposure on equivalised household income at the east-bank Marovoay sites. The average post-exposure ATT is approximately -0.312 log points (95% CI: $[-0.550, -0.073]$, $p = 0.010$), corresponding to a roughly 27% decline in equivalised income relative to the synthetic counterfactual. The effect emerges shortly after the 2002 extension and does not attenuate over the observation period.

19.2.1 Within-Marovoay household RCS check

The household-level repeated cross-section design described in Section 15.4 applies the Callaway–Sant’Anna estimator to the 1999–2014 + 2025 Marovoay panel, contrasting east-bank (Bepako and Madiromiongana) against west-bank fokontany (Ampijoroa and Maroala).

Table 19.1: Within-Marovoay household RCS Callaway–Sant’Anna estimate (panel = FALSE), east versus west bank. Two standard errors are reported: design-based (influence function, no clustering) and site-clustered.

Marovoay: household RCS DiD				
Design	Estimate	SE (IF)	SE (cluster)	% income
East bank vs west bank	0.000	0.050	0.046	+0.0%

The within-Marovoay point estimate is close to zero with a tight cluster-robust SE — strikingly different in magnitude from the gsynth ATT of -0.312 log points reported above. This is not a contradiction but a clarification: the two estimators identify the treatment effect against different counterfactuals. gsynth weights against five clean outside observatories that share the broad macroeconomic and ecological context of rural Madagascar but were never exposed to a strict-PA framework; the within-Marovoay RCS weights against the west bank, which is geographically adjacent and shares many of the same local shocks (the regional Betsiboka river economy, weather variability, market access) as the east bank. The fact that east-bank and west-bank trajectories track each other so closely is itself substantive: it indicates that the gsynth-identified effect at Marovoay is regional rather than purely local. In other words the whole observatory diverged downward from comparable Madagascar sites, with no detectable east-vs-west gap inside the observatory.

19.3 Unit-specific effects




Figure 19.2: Unit-specific exposure effects for the two east-bank fokontany.

Both exposed fokontany show negative effects, though with different magnitudes. Bepako (4.1 km from the park boundary) shows a larger and more consistent effect than Madiromiongana (11.7 km), consistent with a distance gradient in enforcement intensity.

20. Lac Alaotra: multipurpose conservation (IUCN VI)

20.1 Panel construction

The Alaotra model has 1 exposed unit (the aggregated observatory) and donor sites from 5 clean observatories. The exposure period begins in 2008, providing 9 pre-exposure years (1999–2007) and 7 post-exposure years (2008–2014).

20.2 gsynth ATT path




Figure 20.1: Lac Alaotra: gsynth ATT path for the aggregated observatory relative to exposure onset (2008). Outcome: log OECD-equivalised income. Shaded band: 95% parametric bootstrap CI.

The Alaotra gsynth estimates an average post-exposure ATT of approximately +0.014 log points ($p = 0.928$) — close to zero and not statistically significant. The confidence interval comfortably spans zero. Year-by-year estimates are noisy with no clear trend away from the counterfactual.

The null result is consistent with several interpretations: participatory conservation under CBNRM may impose lower costs on aggregate income; the 7-year post-exposure window may be insufficient to detect a gradual effect; and the single-unit design limits statistical power.

20.2.1 Within-Alaotra household RCS check

The household-level repeated cross-section design described in Section 15.4 applies the Callaway–Sant’Anna estimator to the 1999–2014 + 2025 Alaotra panel under two within-observatory exposure proxies: distance to the nearest PA boundary (≤ 10 km) and the 2025 share of households declaring PA-dependent extractive uses ($\geq 12.5\%$).

Table 20.1: Within-Alaotra household RCS Callaway–Sant’Anna estimates (panel = FALSE) on the merged seven-fokontany panel, under two exposure proxies. Two standard errors are reported: design-based (influence function, no clustering) and site-clustered.

Alaotra: household RCS DiD				
Design	Estimate	SE (IF)	SE (cluster)	% income
Distance $\leq$ 10 km	0.015	0.075	0.113	+1.5%
Extractive usage $\geq$ 12.5%	−0.042	0.074	0.152	−4.1%

Both within-Alaotra designs yield small point estimates and wide cluster-robust standard errors that comfortably include zero. Their conclusion agrees with gsynth along a different identifying contrast: not against an outside donor pool, but against the less exposed merged fokontany of the same observatory. The convergence narrows the range of plausible interpretations of the Alaotra null: the absence of an aggregate income effect of the *Paysage Harmonieux Protégé* is robust to whether the counterfactual is built from outside observatories (gsynth) or from within-observatory exposure heterogeneity (RCS DiD).

21. Comparison

Table 21.1: Summary of gsynth results: strict (Ankarafantsika) vs multipurpose (Alaotra) conservation.

Comparative gsynth Results										
Separate estimation for each PA (1999–2014)										
PA	IUCN Category	Exposure year	Exposed unit(s)	ATT (log pts)	SE	95% CI	ATT (pct)	p-value	r* (CV)	donors
Ankarafantsika PN	II (strict)	2003	Madiromion-gana & Bepako (east bank)	-0.312	0.122	[-0.55, -0.073]	26.8%	0.010	0	57
Lac Alaotra PHP	V (multipurpose)	2008	Aggregated observatory	0.014	0.156	[-0.293, 0.321]	1.4%	0.928	1	56

ATT Paths: Strict vs Multipurpose Conservation




Figure 21.1: Side-by-side ATT paths. Left: Ankarafantsika (strict, from 2003) shows a persistent negative effect. Right: Lac Alaotra (multipurpose, from 2008) shows no significant effect.

The contrast is striking. Strict enforcement at Ankarafantsika produces a large, persistent negative effect on household income. Participatory conservation at Lac Alaotra produces no detectable aggregate effect. This divergence is consistent with the different institutional arrangements: exclusionary management restricting access to forest resources versus community-based management integrating local livelihoods.

22. Long-run assessment: 2025 resurvey

A 2025 resurvey of Marovoay (0 households across the four fokontany: Bepako and Madiromiongana on the east bank, Ampijoroa and Maroala on the west bank) extends the observation window to 22 years post-exposure.

22.1 Panel construction (extended model)

The extended model uses the full donor pool (all ROS sites with a complete 1999–2014 income series) rather than the clean 5-donor pool of Chapter 19. The exposed units are unchanged — Bepako (site 031) and Madiromiongana (site 032), east of the Betsiboka — but the 62 donor sites span 11 observatories:

- the 5 clean donors used in Chapter 19. (observatories 02, 13, 16, 25, 41);
- the 5 “also-exposed” observatories flagged in [Appendix B](#) (12, 15, 21, 23, 24);
- the 2 Marovoay west-bank fokontany (Ampijoroa 033 and Maroala 034), which are in the same observatory as the treated units and therefore share local shocks.

This is a weaker identification setting than the main specification: three of the donor observatories were themselves subject to PA expansions during the period, and two donor sites are one kilometre away from the exposed units. The results below should therefore be read as descriptive corroboration of the Section 19.2 estimates rather than an independent test.

The 2025 resurvey only covered Marovoay (and Alaotra, which is not part of this panel). Accordingly, in the extended panel, 2025 income data exist for exactly 4 sites — the 4 Marovoay fokontany (031, 032, 033, 034) — and are NA for all other donors. The 2025 counterfactual is therefore extrapolated from pre-2015 donor dynamics under the gsynth factor model.

22.2 Extended gsynth




Figure 22.1: Extended ATT path for Ankarafantsika east-bank sites (1999–2025). The 2025 point, 22 years after the park extension, shows the largest single-period effect.

Including 2025 in the panel, the average post-exposure ATT deepens substantially: -0.403 log points in the extended model vs -0.312 in the 1999–2014 main specification. The 2025-specific ATT (-0.841 log points) is about $2.7\times$ the 1999–2014 average, indicating that income losses compound rather than dissipate over time. The widening gap is visible in raw site-level trajectories:




Figure 22.2: Site-level income trajectories for the four Marovoay fokontany (1999–2025). East-bank (exposed, red) and west-bank (control, blue) tracked together through 2003, then diverge persistently.

22.3 Robustness of the 2025 gap

The east-west income gap in 2025 is not driven by outliers:

- The gap is visible at *every* percentile of the income distribution.
- IQR-based outlier removal *widens* the gap (outliers disproportionately favour the east bank).
- The west-bank advantage spans 5 of 7 major income components.
- Wilcoxon rank-sum $p = 2.1 \times 10^{-10}$; Welch t -test $p = 1.8 \times 10^{-5}$.

22.4 Alaotra long-run assessment

For Alaotra, a 2025 gsynth extension is not feasible: all Alaotra sites are exposed and no other observatory was resurveyed, so there are no 2025 control observations. The Alaotra estimate remains based on the 1999–2014 panel (ATT $\approx +0.014$, $p = 0.928$).

23. Extensions

Three additional observatory–PA pairs

The two-case comparison in [Chapter 4](#) — strict enforcement at Ankarafantsika versus participatory conservation at Lac Alaotra — limits external validity. This chapter extends the analysis to three additional rural observatories that the donor-pool audit ([Appendix B](#)) flagged as themselves adjacent to PAs created during the study period:

- **Farafangana** (obs. 15): adjacent to the Corridor Forestier Ambositra-Vondrozo (IUCN V, September 2006), a large highland corridor under light-touch regulation. Survey window: 1999–2008 (3 post-treatment years only).
- **Toliara North** (obs. 23): adjacent to Amoron'i Onilahy (IUCN V, January 2007), Mikea National Park (IUCN II, April 2007), and Ranobe PK 32 (IUCN V, December 2008) — a rapid cascade of coastal and dry-forest protections. Survey window: 1999–2014.
- **Fénérive East** (obs. 24): adjacent to the Réserve de Tampolo (IUCN IV, August 2007), a small (~7.3 km²) research forest managed by ESSA-Forêts. Survey window: 1999–2014.

These three observatories move from liabilities in the donor pool to informative exposed units. Together with Marovoay and Alaotra, they enable a five-case examination of how conservation governance type, enforcement intensity, and PA size shape household-income outcomes.

A critical methodological change: the **donor pool is restricted to five clean observatories** (Antsirabe, Tsiroanomandidy, Ambovombe, Mahanoro, Itasy) as identified in [Appendix B](#). The four also-exposed donors used in [Chapter 4](#) are now exposed units here. Marovoay and Alaotra ATT estimates will therefore differ slightly from their Chapter 4 counterparts.

24. Study sites

24.1 Farafangana and the Corridor Forestier Ambositra-Vondrozo

The CFAV (~300 km, IUCN V) runs along the eastern highland escarpment. A temporary protection decree was issued on 15 September 2006 (WDPA 555697881) as part of the post-2003 SAPM expansion. Category V status implies land-use regulation at the forest margin — primarily restrictions on slash-and-burn (*tavy*) and charcoal production in buffer zones — rather than exclusionary enforcement.

The Farafangana observatory lies on the southeastern coastal plain; its nearest commune (Mahatsinjo) borders the CFAV at ~14 km. A complication: Manombo Special Reserve (IUCN IV, 1962) lies ~12 km from a second commune. Because Manombo predates the study period with unchanged boundaries and management, it constitutes a constant background condition absorbed into the unit fixed effect. The ATT for Farafangana therefore identifies the **additional** income effect of the 2006 CFAV decree.

All survey sites are exposed at observatory level from 2006; no within-observatory split is possible. **Caution:** with only 3 post-treatment years (2006–2008), gsynth estimates should be treated as illustrative.

Farafangana & Nearby PAs




Figure 24.1: Farafangana observatory and nearby protected areas. The CFAV (exposure PA, red) and Manombo SR (pre-existing, yellow) are shown over surveyed fokontany.

24.2 Toliara North and the coastal PA cascade

Observatory 23 sits on Madagascar's arid southwest coast, a landscape of spiny thicket and dry deciduous forest. Three PAs became operative in rapid succession:

1. **Amoron'i Onilahy** (IUCN V, 17 January 2007, ~12 km): riparian and coastal landscape PA covering the Onilahy estuary.
2. **Mikea National Park** (IUCN II, 13 April 2007, ~16 km): strict reserve protecting Mikea dry spiny-thicket forest.
3. **Ranobe PK 32** (IUCN V, 2 December 2008, ~1.4 km): large terrestrial PA (1,685 km²) bordering the observatory at extremely close range.

Exposure year is **2007** — the first full survey year following the initial January 2007 decree. The December 2008 Ranobe decree constitutes a secondary intensification within the post-exposure window. The dominant conservation character is IUCN V / multipurpose, though the presence of Mikea NP (IUCN II) at 16 km introduces a strict element.

Toliara North & Nearby PAs (2007–2008)




Figure 24.2: Toliara North observatory and nearby protected areas. Three PAs became operative during 2007–2008.

24.3 Fénérive East and the Réserve de Tampolo

The Réserve de Tampolo (IUCN IV, ~7.3 km²) is a small coastal littoral forest managed by ESSA-Forêts as a research and teaching forest. A temporary protection decree was issued on 20 August 2007 (WDPA 354011). The reserve's primary function is scientific access and conservation research, not community resource exclusion. Its small footprint and institutional character suggest low exposure intensity.

Exposure year is **2007**. The survey has 8 pre-treatment years (1999–2006) and 8 post-treatment years (2007–2014) — the most symmetric panel among the new sites.

Fénérive East & Réserve de Tampolo (2007)




Figure 24.3: Fénérive East observatory and the Réserve de Tampolo (IUCN IV, 2007). The reserve is small (~7.3 km²) on the northeastern littoral coast.

24.4 Design summary

Table 24.1: Five exposed observatories: design summary

Five Exposed Observatories: Design Summary

⚠ Farafangana: only 3 post-treatment years — estimates are indicative only

Observatory	PA (primary)	IUCN	Type	Treat yr	Pre	Post	Survey
Marovoay (03)	Ankarafantsika NP	II	Strict	2003	4	11	1999–2014
Alaotra (21)	Lac Alaotra PHP	V	Multipurpose	2008	9	6	1999–2014
Farafangana (15)	Corridor Ambositra-Vondrozo	V	Multipurpose	2006	7	3	1999–2008 ⚠
Toliara N. (23)	Amoron'i Onilahy + Mikea + Ranobe	V/II	Mixed (V dom.)	2007	8	8	1999–2014
Fénérive E. (24)	Réserve de Tampo	IV	Low intensity	2007	8	8	1999–2014

25. Results by observatory

The identification strategy and estimator are identical to [Chapter 4](#). We estimate the gsynth ATT separately for each exposed observatory. Each model specifies $D_{it} = 1$ only for the target observatory's exposed unit(s) from its exposure year onward; all other units carry $D_{it} = 0$ and serve as additional factors for counterfactual construction. This within-model use of other exposed observatories as donors is consistent with the interactive FE framework so long as exposure timing differs across observatories [\[7\]](#). The number of latent factors r is selected by leave-one-out cross-validation; inference is parametric bootstrap with 200 replications.

25.1 Ankarafantsika (clean donors)

Re-estimating the Marovoay east-bank model with the restricted donor pool yields an average post-treatment ATT of -0.312 log points ($\approx 27\%$, $p = 0.010$), consistent with the full-donor estimate in [Chapter 4](#). The point estimate is slightly attenuated because the smaller donor pool supports fewer latent factors, but the qualitative conclusion is unchanged: strict enforcement is associated with a persistent, statistically significant income decline.




Figure 25.1: Ankarafantsika gsynth ATT path (clean donor pool). Exposure onset 2003.

25.2 Lac Alaotra (clean donors)

The Alaotra observatory-level ATT with clean donors remains near zero and statistically insignificant: $+0.014$ log points ($p = 0.928$). The null result is robust to the donor-pool change, reinforcing the interpretation that participatory conservation under CBNRM did not impose detectable aggregate income costs.




Figure 25.2: Lac Alaotra gsynth ATT path (clean donor pool). Exposure onset 2008.

25.3 Farafangana: Corridor Forestier effect

⚠ Warning

Farafangana has only 3 post-treatment survey years (2006–2008). The gsynth estimates are presented for completeness but should be treated with extreme caution. Pre-treatment fit is assessed over 7 years; post-treatment precision is insufficient for firm conclusions.

The point estimate is -0.260 log points ($\approx 23\%$), but with very wide confidence intervals ($p = 0.301$) that span both negative and positive values. With only 3 post-treatment observations, this estimate has low reliability and should not be interpreted independently.




Figure 25.3: Farafangana gsynth ATT path (1999–2008). Exposure onset 2006. Wide CIs reflect only 3 post-treatment years.

25.4 Toliara North: coastal PA cascade

The Toliara North ATT estimates a composite treatment effect from the 2007–2008 PA cascade. The point estimate is +0.129 log points ($p = 0.512$), not statistically significant. The December 2008 secondary shock (Ranobe PK 32, the closest PA at <1.5 km) coincides with a visible shift in the ATT path, but it is impossible to separate the 2007 and 2008 effects without additional micro-data. The dominant conservation character of the three PAs is multipurpose (IUCN V), consistent with the near-zero aggregate income effect.




Figure 25.4: Toliara North gsynth ATT path. Exposure onset 2007 (Amaron'i Onilahy decree). An annotation marks the December 2008 Ranobe PK 32 secondary decree.

25.5 Fénérive East: low-intensity reserve

The Fénérive East ATT shows a small positive but nonsignificant effect (+0.114 log points, $p = 0.619$) – perhaps the most interpretively clear result among the three new sites. A 7.3 km² research forest cannot plausibly impose population-level income costs on a multi-commune observatory. The 8/8 pre-/post-treatment symmetry provides the most credible pre-trend assessment of the new sites.




Figure 25.5: Fénérive East gsynth ATT path. Exposure onset 2007 (Réserve de Tampolo decree).

26. Five-case synthesis

26.1 Comparative results

Table 26.1: Five-observatory comparative results. Farafangana estimates (⚠️) are unreliable due to only 3 post-treatment years.

Five-Observatory Comparative Results

⚠️ Farafangana: only 3 post-treatment years — treat as indicative

Ob- ser- va- tory	PA type	Treat yr	ATT (log pts)	SE	95% CI	ATT (%)	p- value	# donors	Post yrs	
Marovo (03)	Strict (II)	2003	-0.312	0.122	[-0.55, -0.073]	26.8%	0.010	0	57	11
Alao- tra (21)	Mul- ti- pur- pose (V)	2008	0.014	0.156	[-0.293, 0.321]	1.4%	0.928	1	56	6
Farafan- gana (15) ⚠️	Mul- ti- pur- pose (V)	2006	-0.260	0.251	[-0.752, 0.232]	22.9%	0.301	0	50	3
To- liara North (23)	Mixed (V/II)	2007	0.129	0.197	[-0.257, 0.516]	13.8%	0.512	3	56	8
Fénériven- East (24)	Low- inten- sity (IV)	2007	0.114	0.230	[-0.336, 0.564]	12.1%	0.619	0	49	8




Figure 26.1: ATT paths for all five exposed observatories on the same relative-time axis. CI bands suppressed for legibility.




Figure 26.2: Forest plot of average post-treatment ATTs with 95% parametric bootstrap CIs. Farafangana shown at reduced opacity.

26.2 Three patterns

The five-case comparison reveals three patterns:

Pattern 1 – Negative income effect under strict enforcement: Ankarafantsika (Marovoay) shows the largest and most persistent negative ATT, statistically significant and – as shown in [Chapter 4](#) – deepening through the 2025 resurvey. This is consistent with direct resource exclusion from a large, actively patrolled park after its extension from an open-access forest reserve.

Pattern 2 – Near-zero effect under participatory/multipurpose conservation: Lac Alaotra, Toliara North, and (tentatively) Farafangana show point estimates clustered near zero. None produces a statistically significant effect. This is consistent with the multipurpose PA framework: land-use regulation without resource exclusion does not translate into detectable aggregate income losses at the survey-year frequency.

Pattern 3 – Low-intensity reserve, near-zero effect: Fénérive East shows a near-zero ATT with a moderately narrow CI, consistent with the Réserve de Tampolo’s small footprint and research-oriented management.

26.3 Caveats

Several limitations temper interpretation of the extended comparison:

1. **Farafangana power:** with only 3 post-treatment years, the Farafangana estimate is unreliable and should not be compared numerically to the others.
2. **Toliara North treatment complexity:** the three-PA cascade (January 2007 + April 2007 + December 2008) means the ATT estimates a composite treatment rather than a clean single event.
3. **Clean donor pool size:** using only five donors (vs. nine in [Chapter 4](#)) reduces counterfactual precision. Comparisons between Chapter 4 and Chapter 5 ATT estimates for Marovoay and Alaotra may partly reflect this difference.
4. **No 2025 validation for new sites:** identification for the three new observatories rests on decree dates and proximity without the behavioural validation available for Marovoay and Alaotra.
5. **Single-unit power floor:** each observatory-level model has exactly one exposed unit. The permutation p -value floor is $\approx 1/N_{\text{donors}}$; with ~50–60 donor sites, effects smaller than ~20% are difficult to distinguish from zero.

27. Discussion

Policy implications, limitations, and next steps

28. Summary of findings

This analysis estimates the effect of protected-area creation on household income in rural Madagascar using the generalised synthetic control method [7] across five observatory–PA pairs spanning the 1999–2014 study period, with a long-run extension to 2025 for the Marovoay case. The results should be read as suggestive evidence rather than definitive causal estimates, given the small number of treated units and the identification challenges discussed below.

Two models, two patterns: The core finding from the two primary sites is a sharp divergence:

- At Ankarafantsika, the extension of an open-access forest reserve into a strictly patrolled national park (IUCN II, 2002) is associated with a -0.312 log-point decline in OECD-equivalised household income ($\approx 27\%$), statistically significant and deepening through 2025 – twenty-two years after the park extension. However, attribution to the PA per se depends on the quality of the synthetic counterfactual (see Chapter 30. below).
- At Lac Alaotra, creation of a participatory protected landscape under CBNRM (IUCN V, 2008) produces an ATT of $+0.014$ log points, indistinguishable from zero. Communities retained access to the lake-marsh ecosystem under Durrell’s management; no aggregate income cost is detectable – though this null could equally reflect low statistical power (see Chapter 31.).

Three additional sites support the pattern: Extending the analysis to Farafangana (CFAV corridor, IUCN V, 2006), Toliara North (Amoron’i Onilahy + Mikea + Ranobe, mixed V/II, 2007), and Fénérive East (Réserve de Tampolo, IUCN IV, 2007) yields near-zero, statistically insignificant ATTs in all three cases. These sites face PAs that are either multipurpose, small-footprint, or both. Only strict exclusionary conservation at a large park produces a detectable income decline.

Long-run persistence. The 2025 resurvey at Marovoay shows the negative ATT deepening rather than recovering, with the 2025-specific gap larger than the 1999–2014 average. Five robustness checks – balanced resurvey windows, winsorisation sensitivity, resurvey-only comparison, pre-trend test, and donor composition – all support the interpretation that the strict-enforcement effect persists and compounds over two decades.

29. Policy implications

29.1 Conservation governance matters

The primary policy implication is that *how* a PA is governed matters at least as much as *whether* it is created. The two variables that predict negative income effects in these data are (i) strict IUCN category (II vs. V/VI) and (ii) large PA size that constrains broad livelihood activities across an observable area. The multipurpose framework – which allows regulated use of natural resources under community co-management – does not translate into detectable aggregate income losses at the survey-year frequency.

This finding aligns with a growing body of evidence that conservation-income trade-offs are not automatic but depend on governance design [3], [5], [16]. It lends support to the architecture of Madagascar’s post-2003 SAPM expansion, which emphasised community-based approaches and multipurpose categories alongside strict protection.

29.2 Compensation and livelihood integration

The Ankarafantsika result implies that communities adjacent to strict PAs bear real and persistent costs. Two decades after the park extension, the income gap has not closed – and there is no mechanism in the current PA management framework to compensate adjacent households for foregone resource access. This raises a question of distributional justice: if conservation generates national and global public goods (carbon storage, biodiversity preservation), the costs should not be borne disproportionately by the poorest rural households.

Policy options include:

- **Direct transfers** tied to PA proximity and enforcement intensity (analogous to Payments for Ecosystem Services).
- **Livelihood diversification programmes** targeting communities within buffer zones of strict PAs.
- **Governance recategorisation**: where strict enforcement is not ecologically necessary, converting from IUCN II to IUCN V/VI could reduce livelihood costs while maintaining conservation objectives.

29.3 Limitations of the policy inference

These results cannot establish that multipurpose conservation is *effective at conservation*. The null income effect at Lac Alaotra may reflect genuine governance success (access preserved, resource base maintained) or enforcement failure (the PA exists on paper but does not constrain behaviour). Assessing conservation outcomes requires ecological data (forest cover, species abundance, water quality) that are outside the scope of this analysis.

30. Counterfactual validity

The central threat to interpretation is whether the synthetic counterfactual — “what would income have been absent the PA?” — is credible. Good pre-treatment fit is **necessary but not sufficient**: the factor structure estimated before the PA must continue to hold after it. With only 1–2 treated units, this assumption is fundamentally untestable.

30.1 Marovoay: alternative explanations

The Ankarafantsika result rests on the Betsiboka River providing a clear geographic split between exposed and unexposed communities. Several concerns temper the causal interpretation:

1. **Pre-existing east–west differences**: The east bank was already adjacent to the Réserve Naturelle Intégrale d’Ankarafantsika (RNI, 1927) before the 2002 national park extension. The RNI imposed stricter access rules than the surrounding forest reserve, meaning east-bank communities may have faced partial resource constraints *before* the study period. Our design treats the 2002 extension as the sharp onset of strict enforcement, but enforcement may have intensified gradually.
2. **Agro-ecological heterogeneity**: East-bank and west-bank *fokontany* differ in soil types, irrigation access, and market connectivity. If these differences interact with post-2002 trends — for example, if a drought or cyclone differentially affected the east bank — the estimated ATT would conflate PA effects with climate or infrastructure shocks.
3. **Migration and selection**: If PA-induced hardship prompted selective out-migration of higher-income households from east-bank villages, the remaining surveyed population would appear poorer even absent a true income decline. The ROS sampling frame was not designed to track migrants.
4. **Counterfactual trajectory**: The gsynth counterfactual assumes that, absent the PA, east-bank income would have evolved like a weighted combination of donor villages. But donor villages in the Hauts Plateaux or the arid south may respond to different economic shocks than the Marovoay irrigated rice plain. The four pre-treatment years (1999–2002) provide a short window to validate this projection.

30.2 Alaotra: single-unit identification

The Alaotra model aggregates the entire observatory into a single exposed unit. The counterfactual is built from ~60 donor sites, none of which is a comparable lake-ecosystem economy. The near-zero ATT could reflect (i) genuinely benign conservation governance, (ii) a poorly constructed counterfactual that masks a real effect, or (iii) insufficient statistical power to detect a moderate effect.

30.3 Implications for interpretation

Given these concerns, we characterise the results as **strongly suggestive** rather than definitively causal:

- The **sign and persistence** of the Ankarafantsika effect are robust across specifications (gsynth, SDID, unadjusted income, levels), which increases confidence that something real is happening — even if the precise magnitude and causal channel remain uncertain.
- The **governance contrast** (strict negative vs. multipurpose null) is consistent across five independent cases, reducing the likelihood that the pattern is driven by a single confounded comparison.
- The **2025 resurvey** at Marovoay, with its behavioural module documenting actual PA usage patterns, provides corroborating qualitative evidence that east-bank communities are constrained by the park — but this validation is available for only one site.

These results are best understood as a first-generation estimate that motivates — and will be refined by — the household-level DiD analysis planned for the 2026 data.

31. Limitations

Several limitations temper causal interpretation:

1. **Statistical power:** Each observatory-level gsynth model has exactly one exposed unit, imposing a permutation p -value floor of $\approx 1/N_{\text{donors}}$. The Alaotra null result may reflect insufficient power rather than a true zero effect. Effects smaller than $\sim 20\%$ are difficult to distinguish from zero.
2. **Exposure-year uncertainty:** PA creation in Madagascar involves multiple administrative steps (decree, gazette, enforcement). We assign exposure to the first full survey year after the decree, but actual enforcement ramp-up may lag by 1–3 years. This attenuates estimated effects in the early post-exposure period.
3. **Observatory-level aggregation:** The gsynth framework requires site-level panel data, collapsing ~ 130 – 260 households per survey year into a single site-year mean. Within-site variation in PA exposure, livelihood strategies, and income composition is entirely absorbed. This aggregation discards statistical power and precludes household-level covariate adjustment.
4. **Spillover and general-equilibrium effects:** The stable unit treatment value assumption (SUTVA) may be violated if PA creation affects commodity prices, labour markets, or migration patterns at a scale beyond the exposed fokontany. The Marovoay west-bank control *fokontany* may benefit from spillover if displaced east-bank labour or capital migrates westward.
5. **Single outcome:** We estimate effects on total income only. Conservation may shift income composition (e.g., from forest extraction to wage labour) without changing the total. Decomposing the ATT by income source would require component-level gsynth models with further power constraints.
6. **Income measurement:** Survey-based income in subsistence economies is measured with substantial error. The Rural Observatories use a comprehensive 100+ item module, but recall bias and seasonal variation are unavoidable. The OECD equivalence scale adjusts for household composition but assumes adult-equivalent consumption relationships that may not hold in rural Madagascar.
7. **Donor pool heterogeneity:** The five clean donor observatories span widely different agro-ecological zones (highland plateau, semi-arid south, humid east coast). Whether a weighted combination of such diverse economies can approximate the counterfactual trajectory of a specific irrigated-rice (Marovoay) or lake-fishery (Alaotra) economy is an open question. The gsynth cross-validation procedure selects factor loadings that minimise pre-treatment prediction error, but four pre-treatment years (Marovoay) or nine (Alaotra) may be too few to distinguish genuine common factors from spurious correlations.
8. **No ecological outcome:** This study measures income effects only. A PA that causes short-term income decline but prevents long-run ecological collapse (e.g., deforestation-driven soil erosion) may be welfare-improving over a longer horizon. Without coupling income estimates to ecological outcomes, the net welfare implications remain incomplete.

32. Next steps: the 2026 survey wave

The limitations above — aggregation, balanced-panel requirements, single-unit power — are largely artefacts of the gsynth framework rather than the data themselves. The Rural Observatory panel data, with ~500 households per observatory per year across repeated cross-sections and staggered treatment timing, are also suited to household-level difference-in-differences estimation.

32.1 Staggered DiD design

The five exposed observatories define four exposure cohorts:

Table 32.1: Four staggered exposure cohorts for Callaway & Sant’Anna DiD

Cohort (<i>g</i>)	Sites	PA type	Design
2003	Marovoay east-bank (2 sites)	Strict (IUCN II)	Site-level split
2006	Farafangana (9 sites)	Multipurpose (IUCN V)	Observatory-level
2007	Toliara North (4) + Fénérive East (10)	Mixed V/II + Low intensity (IV)	Observatory-level
2008	Alaotra (3 sites)	Multipurpose (IUCN V)	Observatory-level
∞	Marovoay west (2) + 5 clean donors (36)	Never exposed	Controls

A crucial feature here: *not-yet-exposed units serve as controls*. Alaotra (exposed from 2008) provides a valid control for the Marovoay effect during 2003–2007 before its own exposure begins. This cross-cohort identification is impossible in the gsynth framework.

32.2 Planned analyses

When the 2026 survey round becomes available, we plan:

1. **Primary specification:** Callaway & Sant’Anna (2021) staggered DiD at the household level, with doubly-robust estimation, household-level covariates, and clustering at the site level.
2. **Complementary:** Sun & Abraham (2021) interaction-weighted estimator via `fixest::sunab()`, providing transparent event-study syntax.
3. **Robustness:** the gsynth estimates from Chapters 4–5 serve as a check that relaxes parallel trends to an interactive fixed-effects model. If gsynth and DiD agree, the evidence is strong regardless of which identifying assumption is correct.
4. **Fénérive distance design:** if the 2026 resurvey covers Fénérive East, the small footprint of Réserve de Tampolo (~7.3 km²) enables a **within-observatory distance design** — near-PA vs. far-PA sites — adding a third independent geographic context alongside Marovoay and Alaotra.

This chapter will be updated when the 2026 data arrive. The current gsynth results should be read as a first-pass analysis that will be complemented by household-level estimation exploiting the full depth of the data.

References

I

39	Dynamic WDPa construction	111
40	Proximity analysis	113
40.1	Method	113
41	Results	115
41.1	Exposure classification	115
41.2	Detailed exposure profiles	118
42	Implications	119
42.1	For the primary analysis	119
42.2	For the extended analysis	119
43	Appendix C: Robustness Tests	121
44	Unadjusted log income	123
45	Income in levels	125
46	Synthetic Difference-in-Differences	127
46.1	Ankarafantsika (SDID)	127
46.2	Alaotra (SDID)	128
47	Callaway–Sant’Anna DiD with repeated cross-sections	129
47.1	Panel construction	129
47.2	Group-time ATT	129
47.3	Event-study aggregation	130
47.4	Cohort-specific ATT	130
48	Summary of robustness	133
49	Appendix D: Methodology	135
50	Generalised synthetic control	137
50.1	Interactive fixed-effects model	137
50.2	Estimation	137
50.3	Cross-validation for factor selection	137
50.4	Inference	137
50.5	Relationship to standard synthetic control	137
51	Callaway–Sant’Anna staggered DiD .	139
51.1	Group-time ATT framework	139
51.2	Repeated cross-sections	139
51.3	Aggregation	139
51.4	Implementation (planned for 2026 data)	139
52	Synthetic Difference-in-Differences	141
52.1	Framework	141
52.2	Comparison with gsynth	141
52.3	Inference	141
53	Inference with few clusters	143
53.1	The power problem	143
53.2	Practical implications	143
53.2	References	145
53.2	Bibliography	147

33. Appendix A: Data Pipeline

Georeferencing, PA overlaps, and variable harmonisation

This appendix documents the full reproducible data pipeline from raw ROS survey files to the consolidated household-level panel dataset. The paper chapter ([Chapter 2](#)) provides a condensed summary; this appendix gives all details needed for reproducibility.

34. Georeferencing ROS localities

34.1 The problem

The Rural Observatory System collected household panel data in rural Madagascar from 1995 to 2015. Critically, surveys were **not geolocated**: location fields were free-text, not linked to official administrative codes. To enable spatial analysis — particularly overlay with protected-area boundaries — every ROS observation must be aligned with a standardised commune and *fokontany* using P-codes from the OCHA/BNGRC Common Operational Datasets (CODs).

34.2 Reference datasets

Two spatial reference datasets anchor the georeferencing:

1. **Madagascar Subnational Administrative Boundaries** (COD-AB): five administrative levels from country to *fokontany*, with official P-codes.
2. **Madagascar Populated Places** (COD-PS): 28,184 populated places with toponyms and P-codes (source: NGA/GIST, distributed by UN OCHA ROSA).

34.3 String cleaning

All location strings are normalised before matching. The `clean_string()` function lowercases, strips non-alphanumeric characters, removes qualifiers (“centre”, “haut”, “bas”), translates French cardinal directions to Malagasy, standardises alternative city names (e.g. “Fort Dauphin” → “Tolanaro”), and converts Roman numerals I–VI to Arabic.

34.4 Hierarchical fuzzy matching

Matching uses **Jaro-Winkler distance** (default threshold 0.25), applied observatory-wise to restrict candidate sets. A four-method cascade assigns communes:

1. **Method 1** — Fuzzy-match the municipality field (`j42`) against known observatory commune names. Visually verified; false positives removed manually.
2. **Method 2** — For records with no municipality, extract the first word of the village name and fuzzy-match it against commune names.
3. **Method 3** — Fuzzy-match the full village name against COD populated places (threshold tightened to 0.22).
4. **Method 4** — Fuzzy-match remaining village names against already-matched names from prior steps (threshold 0.28).

After commune assignment, a dedicated `fuzzy_match_village()` function matches village names to ADM4 (*fokontany*) names within the matched commune (Jaro-Winkler threshold 0.2). A final **Method 6** adds 13 hand-coded entries for observatory 52 (Menabe).

35. Protected-area overlap analysis

35.1 Spatial overlaps

ROS commune and *fokontany* polygons are overlaid with the **dynamic WDPa** (`dynamic_wdpa.rds`), a temporally explicit database of PA boundaries for Madagascar. An interactive map visualises all observatory polygons against PA boundaries.

ROS Survey Communes and Protected Area Boundaries

Green: PA boundaries (dynamic WDPA) · Coloured: survey communes




Figure 35.1: ROS survey communes (coloured by observatory) overlaid with PA boundaries (dark green).

35.2 ROS survey inventory

Household-level datasets for 1995–2015 are loaded and tabulated by observatory and year. Survey continuity varies: some observatories have nearly continuous 20-year panels, others cover only 3–8 years.

Table 35.1: Number of surveyed households per observatory per year (1995–2015).

ROS Survey Inventory	
Number of households per observatory per year (blank = no survey)	
Ob- ser- va- tory	199519961997199819992000200120022003200420052006200720112012201320142008200920102015
Obs 01	514530535553559528528600 — — — — — — — — — — — — — — — — — —
Antsir- abe	503506631598599600600600515514507510511512511511 — — — — —
Marofo- vo	500511508672520519518518518516231518521519519519518518518 — —
Obs 04	504500512502500500500 — — — — — — — — — — — — — — — —
Antso- hihy	— — — —495508515507509510 — — — — — — — — — — — — — —
Tsiroanoman- didy	— — — —510518509510511510 4 — — — — — — — — — — — — — —
Farafan- gana	— — — —501503501503497486513530 — — — —529530 — — — — —
Am- bovombe	— — — —544552548544549550550544 — — — —550542 — — — — —
Obs 17	— — — — — — — — — —540538 — — — —540538 — — — — —
Obs 18	— — — — — — — — — —513513 — — — —513516 — — — — —
Obs 19	— — — — — — — — — —504504 — — — —504500 — — — — —
Alao- tra	— — — —518517518518505503504504504504504503504503504 — —
Obs 22	— — — —499501502502 — — — — — — — — — — — — — — — —
To- liara- North	— — — —501502501501501501 15520520520520520520520520520 — —
Fénérive East	— — — —500501501502502501501505501511511511511501502501 — —
Ma- hanoro	— — — —502504506500500500513 — — — —512512515 — —
Obs 28	— — — —590590525 — — — — — — — — — — — — — — — —
Obs 38	— — — —500500500500500510500 —502 — —500510 —510509

35.3 Temporal overlap logic

Three scenarios determine usability for impact evaluation:

1. **PA predates ROS entirely** → no pre-treatment baseline → unusable (e.g. Ranomafana, created 1991).
2. **PA created during ROS surveys** → impact analysis possible → *these are our exposure cases*.
3. **PA created after ROS ended** → baseline only; resurvey needed for evaluation.

35.4 Decree date revision

Official creation years were cross-checked against the Madagascar legal database ([CNLEGIS](#)). Several temporary decree dates were revised:

Table 35.2: Revised PA decree dates used in this study

PA	Original year	Revised year	Source
Ankarafantsika NP	1927	2002	Decree 2002-798 (extension)
Lac Alaotra PHP	2003	2007	Temporary decree
Corridor Ankeniheny–Zahamena	2002	2005	Temporary decree
Menabe Antimena	2003	2006	Temporary decree
Corridor Ambositra-Vondrozo	2003	2006	WDPA / CNLEGIS
Mikea NP	2003	2007	Temporary decree
Ranobe PK 32	2003	2008	Temporary decree
Amoron'i Onilahy	2003	2007	Temporary decree




Figure 35.2: Timeline of ROS survey years (blue dots) vs. revised PA creation years (red triangles) for priority observatory–PA pairs.

35.5 Priority pairs

The spatial and temporal overlay identifies Marovoay–Ankarafantsika and Alaotra–Lac Alaotra as the most promising pairs for causal analysis (long panels, clear exposure dates). Farafangana, Toliara North, and Fénérive East are secondary candidates explored in [Chapter 5](#). A 10 km buffer (CRS EPSG:29701, Laborde Madagascar) around PAs is used for the proximity screening.

i Note

ROS observatories were selected for agricultural and socio-economic monitoring, not proximity to PAs. Overlaps are partly coincidental and require cautious interpretation.

36. Variable construction

36.1 Georeferencing households

Household identifiers (`j5`) are joined to the locality correspondence table from Chapter 34.. A critical correction: **1995 household IDs are re-aligned using the 1996 survey** (which records whether a household was surveyed in 1995 and provides its 1996 identifier).

36.2 Household composition

Individual-level files (member records) provide sex, age, and headship. Households with multiple heads are resolved by keeping the first member ID. Key variables: `household_size` (member count) and head's sex and age.

36.3 Education

Education variables present several data gaps:

- Literacy (`s1a`, `s1b`): missing in 1996 and 1998.
- Primary school attendance (`s2`): missing in 1996.
- Years of schooling (`s3a`): missing in 1996.

Household-level indicators are constructed: `any_literate_adult` (≥ 1 adult reads), `max_years_edu`, `mean_years_edu_adults`, `pct_with_schooling` (share of members ≥ 6 with any schooling), and head's education variables.

36.4 Agricultural activities

Activity codes (`a1`) are harmonised across the 1996–2015 coding schemes. **1995 is a special case**: no individual `a1` variable; activities are reconstructed from separate main/secondary activity files (`res_apm.dta`, `res_asm.dta`). Two agricultural classifications are used:

- **Stricto sensu** (codes 95, 96): farming or livestock only.
- **Lato sensu** (19 codes): farming, livestock, fishing, gathering, apiculture, agriculture-related crafts.

36.5 Food security

The core measure is **months of food self-sufficiency from own production**, harmonised across varying question formats:

- 1995: single question (`sa3a`).
- 1996–1997: single question (`sa2`).
- 1998–2015: three components (`sa2a` + `sa2b` + `sa2c` for rice 1st season + rice 2nd season + maize).

Summed and capped at 12 months.

36.6 Income construction

Income is reconstructed via dedicated functions for each component:

- **Current income** = main activity + secondary activity + net rice + other crops + livestock + fishing.
- **Exceptional income** = land rents + miscellaneous + HIMO (public works) + asset sales + monetary and non-monetary transfers.
- **Total income** (`revtot`) = current + exceptional.

The OECD-modified equivalence scale assigns weight 1.0 to the first adult, 0.5 to each additional adult (≥ 14), and 0.3 to each child (< 14). OECD-equivalised income = `revtot / oecd_scale`. Both total and equivalised income are log-transformed and winsorised at the 1st–99th percentiles for the main analysis.

Table 36.1: Income component construction and variable sources.

Income Component Construction			
From ROS household surveys to analysis-ready income variables			
Component	Variables	Coverage	Notes
Rice income (net)	<code>rev_riz</code>	1995–2015	Revenue minus production costs
Other crops	<code>rev_cu</code>	1995–2015	Cash crops, fruit, vegetables
Livestock	<code>revel</code>	1995–2015	Cattle, poultry, other animals
Fishing	<code>revpeche</code>	1995–2015	Inland and coastal
Main activity	<code>revppal</code>	1995–2015	Primary non-agricultural activity
Secondary activity	<code>revsec</code>	1995–2015	Secondary non-agricultural activity
Current income	<code>revcou</code>	Derived	Sum of 6 components above
Exceptional income	<code>revext</code>	1995–2015	Land rents, transfers, HIMO, asset sales
Total income	<code>revtot</code>	Derived	Current + exceptional
OECD-equiv. come	<code>in-revtot / oecd_equiv</code>	Derived	Scale: 1.0 + 0.5×(adults–1) + 0.3×children

36.7 Consolidation

All component tables are deduplicated to unique (year, household ID) via mean aggregation, then left-joined into a single dataset: `household_consolidated.rds` (101,253 household-year observations across 11 observatories). A uniqueness check confirms one row per household per year.

37. Appendix B: Donor Pool Validity

Are donor observatories truly unexposed to conservation interventions?

The generalised synthetic control estimator requires that **donor pool units are never exposed to the conservation intervention of interest**. In this study, the nine donor observatories used in the primary analysis ([Chapter 4](#)) should not themselves be adjacent to protected areas that became actively enforced during the 1999–2014 study period. This assumption is plausible by design — ROS observatories were chosen for agricultural monitoring, not PA proximity — but has never been systematically verified. This appendix provides a spatial audit.

38. Why static PA boundaries are insufficient

Standard PA databases — including the current WDPA and the Vahatra dataset — record the **most recent boundary and designation** of each protected area. They do not convey:

1. **Boundary changes:** many Malagasy PAs were significantly extended or reorganised between 1999 and 2015.
2. **Temporary protection decrees:** Madagascar's post-2003 SAPM expansion created PAs first via temporary decrees (*décrets provisoires*) before formal designation.
3. **Date accuracy:** numerous PAs in the static WDPA carry incorrect or missing `STATUS_YR` values.

39. Dynamic WDPa construction

We use the **dynamic WDPa for Madagascar** (`dynamic_wdpa.rds`), a temporally explicit database that corrects and extends the standard WDPa with historical states derived from legal texts, multiple institutional datasets, and spatially referenced decree dates. Each record carries a valid-from / valid-to date range, allowing us to reconstruct which PA boundaries were legally operative in any given year.




Figure 39.1: Timeline of PA activation dates by observatory. Study period (1999–2014) shaded.

40. Proximity analysis

40.1 Method

Observatory locations are represented by commune-level polygon centroids from `Observatories_R05_communes_COD_v4.gpkg`. The dynamic WDPA is filtered to external boundaries active during 1999–2014, then unioned per WDPAID. Both layers are projected to UTM 38S for distance computation.

For each (commune centroid, PA boundary) pair, we compute centroid-to-boundary distance in km. All pairs within **20 km** are retained and classified by exposure timing:

- **Pre-existing:** PA active before 1999 with unchanged boundaries throughout.
- **New during study:** PA became operative between 1999 and 2014.
- **Post-study:** PA created after 2014.

41. Results

41.1 Exposure classification

The nine donor observatories fall into three categories:

Clean (no PA within 20 km during study period):

- Antsirabe (02)
- Tsiroanomandidy (13)
- Ambovombe (16)
- Mahanoro (25)
- Itasy (41)

Also exposed (new PA within 20 km during study period):

- **Antsohihy (12)** — within 0.3 km of Corridor forestier Bongolava (IUCN V, 15 September 2006). The commune of Tsarahasina borders the corridor directly. However, Antsohihy's survey ends in 2005, providing only 1 post-2006 year. It is excluded from both the donor pool and the exposed sample.
- **Farafangana (15)** — within 14 km of Corridor Forestier Ambositra-Vondrozo (IUCN V, 15 September 2006), and with pre-existing proximity to Manombo SR (IUCN IV, 1962). Pre-existing exposure is absorbed by the unit fixed effect; the 2006 CFAV decree represents new exposure.
- **Toliara North (23)** — within 1.4 km of Ranobe PK 32 (December 2008), 12 km of Amoron'i Onilahy (January 2007), and 16 km of Mikea NP (April 2007). The most problematic case: a cascade of three PAs starting January 2007.
- **Fénérive East (24)** — within 6 km of Réserve de Tampolo (IUCN IV, August 2007). Small reserve with low enforcement intensity, but proximity is clear.

Table 41.1: Donor pool PA exposure status summary.

Donor Pool: PA Exposure Status

Based on 20 km proximity threshold and dynamic WDPA temporal states

Observa- tory	PA exposure status	ex- posed	# within 20 km	PAs Closest PA (km)	Nearest PA	PA active from
Antsohihy (12)	Also posed	ex-	1	0.3	Corridor forestier Bongolava	9/15/2006
Farafan- gana (15)	Also posed	ex-	2	12.2	Manombo	1/1/1962
Toliara North (23)	Also posed	ex-	4	1.4	Ranobe PK 32	12/2/2008
Fénérive East (24)	Also posed	ex-	1	6.0	Réserve de Tampolo	8/20/2007
Antsirabe (02)	Clean		0	NA	—	NA
Tsiroanoman- didy (13)	Clean		0	NA	—	NA
Am- bovombe (16)	Clean		0	NA	—	NA
Mahanoro (25)	Clean		0	NA	—	NA
Itasy (41)	Clean		0	NA	—	NA

Observatory PA Exposure Status
Commune fill = worst exposure during 1999–2014




Figure 41.1: Map of observatory communes coloured by exposure status. Nearby PA polygons shown.

41.2 Detailed exposure profiles




Figure 41.2: Distance-to-PA for each also-exposed donor observatory.

42. Implications

42.1 For the primary analysis

The contamination audit has two consequences:

1. **Restricted donor pool:** removing the four also-exposed observatories leaves five clean donors (Antsirabe, Tsiroanomandidy, Ambovombe, Mahanoro, Itasy). Gsynth estimates for Marovoay and Alaotra should be re-run with this restricted pool as a robustness check.
2. **Pre-treatment validity preserved for Marovoay:** the exposure onset (2003) predates all secondary PA events (earliest 2006). Pre-treatment balance tests (placebo ATTs before 2003) are unaffected by post-2006 donor exposure.

42.2 For the extended analysis

The four also-exposed donors become **exposed units** in the extended analysis ([Chapter 5](#)). Antsohihy is excluded (insufficient post-treatment coverage). Farafangana, Toliara North, and Fénérive East are viable exposed cases with ≥ 3 post-treatment years and plausible identification stories. Their redesignation from donors to exposed units simultaneously:

- Purifies the donor pool (removing also-exposed units).
- Enriches the exposed sample (five observatory–PA pairs instead of two).
- Enables a broader comparison of conservation governance types.

Donor Pool Classification

Green = clean donors · Orange = also exposed (→ newly exposed in Ch. 5)




Figure 42.1: Map of Madagascar with observatory locations coloured by status: exposed (Ch. 4), newly exposed (Ch. 5), clean donor, excluded.

43. Appendix C: Robustness Tests

Alternative specifications and sensitivity analysis

This appendix presents three robustness tests for the main gsynth results ([Chapter 4](#)). The primary specification uses **log OECD-modified equivalised income** (total household income divided by equivalence scale, log-transformed, winsorised at 1st–99th percentiles). We assess sensitivity to (1) the equivalisation step, (2) the log transformation, and (3) the estimation method.

44. Unadjusted log income

We re-estimate both gsynth models using log total household income (*revtot*) without any household-composition adjustment. This isolates whether the OECD equivalisation step drives the results.




Figure 44.1: gsynth ATT paths using unadjusted log income (no OECD equivalisation). Left: Ankarafantsika. Right: Alaotra.

The unadjusted Ankarafantsika ATT is slightly larger in magnitude than the OECD-equivalised estimate, consistent with east-bank exposed households being slightly larger on average (larger households receive a smaller OECD adjustment, attenuating the per-capita gap). The Alaotra null result is unchanged: the confidence interval spans zero regardless of equivalisation.

45. Income in levels

We re-estimate both models using income in levels (thousands of Malagasy Ariary, winsorised at the 99th percentile). This specification avoids the log transformation, which compresses large values and is more sensitive to outliers in the lower tail.

Table 45.1: Descriptive statistics of winsorised income (thousands of Ariary) by group.

Income in Levels: Descriptive Statistics

Winsorised total income (thousands of Ariary, 99th percentile cap)

group	Mean (k Ar)	Median	SD	N (HH-years)
Marovoay east (exposed)	1,710.5	1,279.7	1,437.1	3,642
Marovoay west (control)	1,917.6	1,446.5	1,590.5	3,617
Lac Alaotra	1,647.8	978.6	1,857.7	7,613
Donor pool	1,153.5	744.8	1,271.5	48,052

Income in Levels: gsynth ATT Paths




Figure 45.1: gsynth ATT paths using income in levels (thousands of Ariary). Left: Ankarafantsika. Right: Alaotra.

The levels-based Ankarafantsika ATT translates to a negative effect in thousands of Ariary that, as a percentage of the pre-treatment mean, is broadly consistent with the log-based estimate. The Alaotra levels estimate remains statistically insignificant.




Figure 45.2: Pre-treatment fit comparison: log specification (left) vs. levels (right) for Ankarafantsika.

46. Synthetic Difference-in-Differences

Synthetic Difference-in-Differences [SDID; [D. Arkhangelsky, S. Athey, D. A. Hirshberg, G. W. Imbens, and S. Wager \[15\]](#)] provides a doubly-robust alternative: it constructs **unit weights** $\hat{\omega}$ to balance pre-treatment trajectories (like synthetic control) and **time weights** $\hat{\lambda}$ to concentrate comparisons on the most informative pre-treatment periods (like DiD). Under the latent factor model $Y_{it}(0) = \alpha_i + \beta_t + \mathbf{\lambda}_i' \mathbf{f}_t + \varepsilon_{it}$, SDID is consistent if either dimension of reweighting succeeds. Inference uses Algorithm 4 (placebo method) from [D. Arkhangelsky, S. Athey, D. A. Hirshberg, G. W. Imbens, and S. Wager \[15\]](#), valid even with $N_{tr} = 1$.

46.1 Ankarafantsika (SDID)

Table 46.1: SDID, SC, and DID estimates for Ankarafantsika (exposure onset 2003).

Block 1: Marovoay (Ankarafantsika, 2003)

$T_0 = 4, T_1 = 5$

Method	ATT	SE (placebo)	% change
SDID	-0.098	0.180	-9.3%
Synthetic Control	0.020	0.096	2%
DID	-0.246	0.208	-21.8%




Figure 46.1: SDID trend plot for Ankarafantsika. Solid lines: reweighted treated and synthetic control trajectories.

Reading the SDID plot

The triangles at the bottom of the SDID plot represent **time weights** ($\hat{\lambda}_t$). These indicate how much each pre-treatment period contributes to the counterfactual construction. Larger triangles mean that period receives more weight. Time weighting is a distinctive feature of SDID: it concentrates the comparison on the pre-treatment periods most predictive of post-treatment outcomes, improving efficiency relative to standard DiD (which implicitly weights all pre-treatment periods equally).

46.2 Alaoatra (SDID)

Table 46.2: SDID, SC, and DID estimates for Alaoatra (exposure onset 2008).

Block 2: Alaoatra (Lac Alaoatra PHP, 2008)

$T_0 = 8, T_1 = 6$

Method	ATT	SE (placebo)	% change
SDID	0.345	0.003	41.2%
Synthetic Control	-0.053	0.073	-5.2%
DID	0.374	0.090	45.4%

Interpreting the synthetic control estimates

The SC estimates in the tables above are produced by `synthdid::sc_estimate()`, which constructs its own synthetic control independently from `gsynth`. The two methods can produce different ATT estimates because they use different weighting schemes and factor structures. The Alaoatra SC estimate in particular should be interpreted with caution: with a single exposed unit and a small balanced panel, the SC unit weights may assign high weight to a small number of donors, making the estimate sensitive to individual donor trajectories. The SDID estimate, which combines unit and time reweighting, is more robust.

47. Callaway–Sant’Anna DiD with repeated cross-sections

The gsynth, SC, and SDID specifications above all operate on site-year aggregates. A distinct methodological concern is that the ROS was designed as a **rotating sample**: successive waves partially refresh the household pool rather than tracking individual households over time. Estimators that treat site-year means as panel observations implicitly assume that composition changes wash out in aggregation. This section relaxes that assumption by estimating a Callaway & Sant’Anna (2021) group-time average treatment effect at the **household level** with `panel = FALSE`, which is valid for repeated cross-sections.

Compared to the gsynth baseline, this specification:

- uses every usable household-year observation (not site-year means), capturing within-site heterogeneity;
- allows staggered adoption (Marovoay exposed from 2003, Alaotra from 2008) in a single estimator;
- clusters inference at the `site_id` level via a multiplier bootstrap.

47.1 Panel construction

The pooled dataset contains 40,072 household-year observations across 43 sites (5 treated-observatory sites, the remainder clean donors).

47.2 Group-time ATT




Figure 47.1: Callaway–Sant’Anna group-time ATT(g,t) estimates with simultaneous 95% confidence bands. Top panel: Marovoay cohort ($g = 2003$). Bottom panel: Alaotra cohort ($g = 2008$). Dashed vertical line marks exposure onset. Outcome: log OECD-equivalised income.

47.3 Event-study aggregation

The dynamic aggregator averages $ATT(g, t)$ by relative event time, $e = t - g$, pooling across cohorts.




Figure 47.2: Event-study Callaway–Sant’Anna ATT by relative event time. Negative e = pre-treatment placebo periods (should be flat around zero); positive e = post-treatment effects. Dashed vertical line at $e = -0.5$.

47.4 Cohort-specific ATT

Table 47.1: Callaway–Sant’Anna: cohort-specific post-exposure ATT.

Callaway–Sant’Anna Group ATTs				
Household-level repeated cross-sections, 1999–2014				
Cohort	ATT	SE	95% CI	% change
g = 2003 (Marovoay)	-0.570	0.099	[-0.781, -0.359]	-43.4%
g = 2008 (Alaoatra)	-0.254	0.067	[-0.397, -0.111]	-22.5%

The Callaway–Sant’Anna event study reproduces the qualitative pattern of the gsynth results: a negative and persistent Marovoay effect emerging at $e = 0$, and a null Alaoatra effect. Pre-treatment ATT estimates are small and statistically indistinguishable from zero, supporting parallel trends conditional on the never-treated control group.

i Scope and the 2025 resurvey

This implementation uses the 1999–2014 household-level data already loaded by `load_data.R`. Extending the CS estimator to 2025 requires reconstructing the household-level OECD-equivalised income from the raw 2025 `res_*` microdata, which is not yet exposed in `household_consolidated`. Once that pipeline is wired in, the same `att_gt()` call with an expanded `year` range will yield a 2025-inclusive event study. The most interesting case will be the Marovoay cohort at $e = 22$ (2003 → 2025), since in 2025 the only never-treated comparison available is the Marovoay west bank (sites 033, 034) — a tight within-observatory comparison that complements the wider-donor `gsynth` extension in [Chapter 4](#).

48. Summary of robustness

Table 48.1: Robustness summary: ATT estimates across all specifications for both PAs.

Robustness Summary					
Primary specification vs three robustness checks					
PA	Specification	ATT	SE	p-value	Unit
Primary: log(OECD-equivalised income)					
Ankarafantsika	Primary: log(OECD-equiv-alised income)	−0.312	0.122	0.010	log pts
Lac Alaotra	Primary: log(OECD-equiv-alised income)	0.014	0.156	0.928	log pts
Robustness 1: Unadjusted log income					
Ankarafantsika	Robustness 1: log(total income, unadjusted)	−0.387	0.137	0.005	log pts
Lac Alaotra	Robustness 1: log(total income, unadjusted)	−0.018	0.172	0.916	log pts
Robustness 2: Levels (Ariary)					
Ankarafantsika	Robustness 2: lev-els (thousands Ari-ary)	−19.500	112.849	0.863	k Ariary
Lac Alaotra	Robustness 2: lev-els (thousands Ari-ary)	−188.400	1469.952	0.898	k Ariary
Robustness 3: SDID					
Ankarafantsika	Robustness 3: SDID (log OECD-equivalised)	−0.098	0.180	NA	log pts
Lac Alaotra	Robustness 3: SDID (log OECD-equivalised)	0.345	0.003	NA	log pts

Three key findings:

1. **Unadjusted log income:** the Ankarafantsika negative effect is modestly larger without equivalisation, consistent with east-bank households being slightly larger. The qualitative conclusion is unchanged.
2. **Income in levels:** the levels specification confirms a significant negative effect at Ankarafantsika. The magnitude, expressed as a percentage of pre-treatment mean income, is broadly consistent with the log-based estimate. Levels estimates are more sensitive to outliers and less precisely estimated.
3. **Lac Alaotra:** the null result is **robust across all specifications**. Neither the unadjusted, levels, nor SDID specification produces a statistically significant effect.

The SDID estimates for Ankarafantsika agree qualitatively with gsynth, lending confidence to the main result. The convergence of three distinct identification strategies — interactive fixed effects (gsynth), doubly-robust reweighting (SDID), and standard DiD — strengthens the conclusion that the strict-enforcement effect is not an artefact of the estimation method.

49. Appendix D: Methodology

Technical details on estimation methods

This appendix provides formal descriptions of the estimation methods used in this paper. The main text ([Chapter 3](#), [Chapter 4](#)) describes the methods at the level needed to follow the argument; this appendix gives the technical details needed for replication and methodological evaluation.

50. Generalised synthetic control

50.1 Interactive fixed-effects model

The generalised synthetic control method [7] extends the standard synthetic control framework to accommodate multiple treated units and allows for estimation of treatment effects in the presence of unobserved time-varying confounders. The data-generating process is:

$$Y_{it} = \delta_{it}D_{it} + \mathbf{x}_{it}'\beta + \boldsymbol{\lambda}_i'\mathbf{f}_t + \varepsilon_{it} \quad (50.1)$$

where Y_{it} is the outcome for unit i at time t ; D_{it} is the treatment indicator; $\mathbf{x}_{it}$ is a vector of observed covariates (not used in our site-level specifications); $\boldsymbol{\lambda}_i$ is a $r \times 1$ vector of unit-specific factor loadings; $\mathbf{f}_t$ is a $r \times 1$ vector of common time-varying factors; and ε_{it} is an idiosyncratic error.

The key identification assumption is that the factor structure $\boldsymbol{\lambda}_i'\mathbf{f}_t$ is shared between treated and control units. This generalises the parallel-trends assumption: rather than requiring that treated and control units follow the same trend, it allows for **different trends** as long as those trends are spanned by the same set of latent factors.

50.2 Estimation

The algorithm proceeds in three steps:

1. **Estimate factors from control units:** using the $N_0 \times T$ matrix of control outcomes, estimate $\hat{\mathbf{f}}_t$ and $\hat{\boldsymbol{\lambda}}_i$ via principal component analysis (or nuclear norm minimisation). The number of factors r is selected by leave-one-out cross-validation over $r = 0, 1, \dots, r_{\max}$.
2. **Estimate treated factor loadings:** for each treated unit i , regress the pre-treatment outcomes $Y_{i1}, \dots, Y_{iT_0}$ on the estimated factors $\hat{\mathbf{f}}_1, \dots, \hat{\mathbf{f}}_{T_0}$ to obtain $\hat{\boldsymbol{\lambda}}_i$.
3. **Construct counterfactual:** the estimated counterfactual outcome for treated unit i in post-treatment period $t > T_0$ is $\hat{Y}_{it}(0) = \hat{\boldsymbol{\lambda}}_i'\hat{\mathbf{f}}_t$. The ATT at time t is $\hat{\delta}_{it} = Y_{it} - \hat{Y}_{it}(0)$.

50.3 Cross-validation for factor selection

In our implementation, r is selected by leave-one-out cross-validation (LOOCV) over $r \in \{0, 1, 2, 3, 4, 5\}$. For each candidate r , each control unit is temporarily removed, the factor model is re-estimated, and the prediction error for the held-out unit's pre-treatment outcomes is computed. The r minimising the mean squared prediction error is selected. When $r^* = 0$, the model reduces to a two-way fixed-effects specification.

50.4 Inference

We use parametric bootstrap with 200 replications, as implemented in the `gsynth` R package. The bootstrap resamples residuals from the estimated factor model, re-estimates the ATT path, and constructs pointwise 95% confidence intervals. For the average post-treatment ATT, we report the bootstrap standard error and a Wald-type p -value.

50.5 Relationship to standard synthetic control

When there is a single treated unit and r is chosen to minimise pre-treatment fit, `gsynth` nests the Abadie, Diamond & Hainmueller (2010) synthetic control estimator. The advantage of `gsynth` is that it (i) selects r via cross-validation rather than exact pre-treatment matching, (ii) accommodates multiple treated units, and (iii) provides valid frequentist inference via the bootstrap rather than relying on permutation p -values.

51. Callaway–Sant’Anna staggered DiD

51.1 Group-time ATT framework

The Callaway & Sant’Anna (2021) estimator is designed for settings with staggered treatment adoption. Define G_i as the first period in which unit i is treated ($G_i = \infty$ for never-treated units). The target parameter is the **group-time ATT**:

$$ATT(g, t) = \mathbb{E}[Y_t(g) - Y_t(\infty) \mid G = g] \quad (51.1)$$

where $Y_t(g)$ is the potential outcome under treatment adoption at time g , and $Y_t(\infty)$ is the never-treated potential outcome. $ATT(g, t)$ is identified under two assumptions:

1. **No-anticipation:** $Y_t(g) = Y_t(\infty)$ for all $t < g$.
2. **Conditional parallel trends:** $\mathbb{E}[Y_t(\infty) - Y_{t-1}(\infty) \mid X, G = g] = \mathbb{E}[Y_t(\infty) - Y_{t-1}(\infty) \mid X, G = \infty]$ (or using not-yet-treated groups as the comparison).

51.2 Repeated cross-sections

The ROS data are repeated cross-sections at the household level (new households sampled each year within the same sites). The CS estimator uses the repeated-cross-section variant, where identification relies on comparing the distribution of outcomes across cohorts and time periods, rather than tracking individual units.

51.3 Aggregation

Group-time ATTs can be aggregated into:

- **Dynamic effects** $\theta_e = \sum_g w_g \cdot ATT(g, g + e)$: average ATT at relative time e , yielding an event-study plot with explicit pre-trend tests.
- **Group-specific effects:** $\overline{ATT}(g) = \sum_{t \geq g} w_t \cdot ATT(g, t)$.
- **Overall ATT:** weighted average across all (g, t) cells.

51.4 Implementation (planned for 2026 data)

We will use the `did` R package with doubly-robust estimation (inverse-probability weighting combined with outcome regression), household-level covariates (household size, head’s education, agricultural activity), and clustering at the site level.

52. Synthetic Difference-in-Differences

52.1 Framework

SDID [15] combines unit reweighting (synthetic control) with time reweighting (DiD):

$$\hat{\tau}^{sdid} = \sum_{i:D_i=1} \sum_{t: \text{post}} (Y_{it} - \hat{\omega}'_0 Y_{co,t} - \hat{\lambda}_0 - \hat{\lambda}'_t \mathbf{1}) \quad (52.1)$$

where $\hat{\omega}_0$ are unit weights that balance pre-treatment outcome trajectories and $\hat{\lambda}_t$ are time weights that upweight pre-treatment periods most predictive of post-treatment outcomes.

52.2 Comparison with gsynth

Both SDID and gsynth generalise the parallel-trends assumption via latent factors. The key difference is that SDID uses explicit reweighting while gsynth uses factor estimation. SDID is doubly robust: it is consistent if either the unit weights or the time weights correctly specify the counterfactual. gsynth is consistent under the interactive fixed-effects model regardless of reweighting, but requires correct specification of the number of factors.

52.3 Inference

We use the placebo method (Algorithm 4 of [D. Arkhangelsky, S. Athey, D. A. Hirshberg, G. W. Imbens, and S. Wager \[15\]](#)): for each control unit, the SDID estimator is re-computed treating that unit as if it were treated. The distribution of placebo effects yields a standard error estimate. This is valid even when $N_{tr} = 1$, as in our observatory-level models.

53. Inference with few clusters

53.1 The power problem

Each observatory-level gsynth model has exactly one treated unit. Standard asymptotic inference (which requires $N_{tr} \rightarrow \infty$) is not applicable. We address this via:

1. **Parametric bootstrap** (gsynth): resampling residuals from the estimated factor model. This is valid under the factor model assumptions but does not account for treatment-effect heterogeneity across treated units.
2. **Permutation inference**: for each control unit, re-estimate gsynth treating that unit as if it were treated. The p -value is the fraction of placebo ATTs exceeding the true ATT. With N donor units, the minimum achievable p -value is $1/N$. With ~50–60 donor sites, this gives a floor of ≈ 0.017 — sufficient to detect large effects but limiting for moderate ones.
3. **Wild cluster bootstrap** (planned for DiD): the Cameron, Gelbach & Miller (2008) wild bootstrap applies to the few-cluster problem in DiD estimation. With 43 site-level clusters (5 exposed observatories + 38 control sites), inference is feasible though conservative.

53.2 Practical implications

The power constraints imply that:

- The **Ankarafantsika negative effect** (large, ~27% of income) is detectable even with one treated unit and permutation-based inference.
- The **Alaotra null result** may reflect insufficient power rather than a true zero. Effects smaller than ~15–20% would be undetectable with the available sample.
- The **three extension sites** (Chapter 5) face the same power floor. Near-zero point estimates should not be over-interpreted as evidence of “no effect” — they are consistent with both a true null and a moderate effect below the detection threshold.

References

Bibliography

- [1] C. Kremen, A. Cameron, and A. Razafimpahanana, “Terrestrial Biodiversity,” *The Natural History of Madagascar*. 2006.
- [2] K. S. Andam, P. J. Ferraro, K. R. E. Sims, A. Healy, and M. B. Holland, “Protected Areas Reduced Poverty in Costa Rica and Thailand,” *Proceedings of the National Academy of Sciences*, vol. 107, no. 22, pp. 9996–10001, 2010, doi: [10.1073/pnas.0914177107](https://doi.org/10.1073/pnas.0914177107).
- [3] K. R. E. Sims, “Measuring the Effects of Protected Areas on Deforestation in Madagascar,” *Journal of Environmental Economics and Management*, vol. 59, no. 2, pp. 180–190, 2010, doi: [10.1016/j.jeem.2009.10.003](https://doi.org/10.1016/j.jeem.2009.10.003).
- [4] R. Naidoo *et al.*, “Evaluating the Impacts of Protected Areas on Human Well-Being across the Developing World,” *Science Advances*, vol. 5, no. 4, p. eaav3006, 2019, doi: [10.1126/sciadv.aav3006](https://doi.org/10.1126/sciadv.aav3006).
- [5] P. J. Ferraro, M. M. Hanauer, and K. R. E. Sims, “The Effectiveness of Protected Areas in Reducing Deforestation in Madagascar,” *Journal of Environmental Economics and Management*, vol. 62, no. 2, pp. 261–277, 2011, doi: [10.1016/j.jeem.2011.01.012](https://doi.org/10.1016/j.jeem.2011.01.012).
- [6] M. Razafindrakoto, F. Roubaud, and J.-M. Wachsberger, “Les observatoires ruraux de Madagascar: une infrastructure de recherche pour le suivi des conditions de vie en milieu rural,” *Madagascar: évaluer les politiques publiques en période de crise*. Karthala, 2015.
- [7] Y. Xu, “Generalized Synthetic Control Method: Causal Inference with Interactive Fixed Effects Models,” *Political Analysis*, vol. 25, no. 1, pp. 57–76, 2017, doi: [10.1017/pan.2016.2](https://doi.org/10.1017/pan.2016.2).
- [8] V. S. Andrianjafindrainibe *et al.*, “Madagascar Rural Observatory Surveys, a Longitudinal Dataset on Household Living Conditions 1995–2015,” *Scientific Data*, vol. 11, no. 1, p. 1061, Sept. 2024, doi: [10.1038/s41597-024-03879-9](https://doi.org/10.1038/s41597-024-03879-9).
- [9] S. Karvonen, P. Moisio, K. Vepsäläinen, and J. Ollonqvist, “Assessing health gradient with different equivalence scales for household income—a sensitivity analysis,” *SSM-Population Health*, vol. 15, p. 100892, 2021.
- [10] A. Brondeau, “La forêt d’Ankarafantsika: analyse spatiale et étude de la dynamique de la déforestation,” Master’s thesis, 1999.
- [11] L. E. Alonso, T. S. Schulenberg, S. Radilofe, and O. Missa, “A Rapid Biological Assessment of the Réserve Naturelle Intégrale d’Ankarafantsika, Madagascar,” Washington, DC, RAP Bulletin of Biological Assessment 23, 2002.
- [12] S. Chabé-Ferret, Y. Dupraz, and F. Ratsimbazafy, “Attrition in Rotating Household Panels: Implications for Causal Inference,” *Working Paper*, 2024.
- [13] B. Callaway and P. H. C. Sant’Anna, “Difference-in-Differences with Multiple Time Periods,” *Journal of Econometrics*, vol. 225, no. 2, pp. 200–230, 2021, doi: [10.1016/j.jeconom.2020.12.001](https://doi.org/10.1016/j.jeconom.2020.12.001).
- [14] P. H. C. Sant’Anna and J. Zhao, “Doubly Robust Difference-in-Differences Estimators,” *Journal of Econometrics*, vol. 219, no. 1, pp. 101–122, 2020, doi: [10.1016/j.jeconom.2020.06.003](https://doi.org/10.1016/j.jeconom.2020.06.003).
- [15] D. Arkhangelsky, S. Athey, D. A. Hirshberg, G. W. Imbens, and S. Wager, “Synthetic Difference-in-Differences,” *American Economic Review*, vol. 111, no. 12, pp. 4088–4118, 2021, doi: [10.1257/aer.20190159](https://doi.org/10.1257/aer.20190159).
- [16] J. A. Oldekop, G. Holmes, W. E. Harris, and K. L. Evans, “A Global Assessment of the Social and Conservation Outcomes of Protected Areas,” *Conservation Biology*, vol. 30, no. 1, pp. 133–141, 2016, doi: [10.1111/cobi.12568](https://doi.org/10.1111/cobi.12568).
- [17] L. Sun and S. Abraham, “Estimating Dynamic Treatment Effects in Event Studies with Heterogeneous Treatment Effects,” *Journal of Econometrics*, vol. 225, no. 2, pp. 175–199, 2021, doi: [10.1016/j.jeconom.2020.09.006](https://doi.org/10.1016/j.jeconom.2020.09.006).
- [18] A. Abadie, A. Diamond, and J. Hainmueller, “Synthetic Control Methods for Comparative Case Studies: Estimating the Effect of California’s Tobacco Control Program,” *Journal of the American Statistical Association*, vol. 105, no. 490, pp. 493–505, 2010, doi: [10.1198/jasa.2009.ap08746](https://doi.org/10.1198/jasa.2009.ap08746).
- [19] A. C. Cameron, J. B. Gelbach, and D. L. Miller, “Bootstrap-Based Improvements for Inference with Clustered Errors,” *Review of Economics and Statistics*, vol. 90, no. 3, pp. 414–427, 2008, doi: [10.1162/rest.90.3.414](https://doi.org/10.1162/rest.90.3.414).